

## Cardiovascular Effects of Carbon Dioxide in Man

David J. Cullen, M.D.,\* and Edmond I. Eger, II, M.D.†

Circulatory responses to administration of carbon dioxide were determined in 41 awake human volunteers during controlled or spontaneous respiration. Compared with spontaneous respiration ( $P_{aCO_2}$ , 37 torr), controlled respiration ( $P_{aCO_2}$ , 36 torr) was associated with a significantly lower cardiac index and stroke index but with an increased total peripheral resistance which maintained constant arterial blood pressure. When exogenous  $CO_2$  was administered during either controlled or spontaneous respiration (range 39 → 50 torr), cardiac index, heart rate, stroke index, indices of myocardial contractility, and forearm blood flow all increased significantly, while total peripheral resistance decreased significantly. These data serve as reference points for measurements of cardiovascular function in normal man, and for studying the modification of the circulatory response to  $CO_2$ . Anesthetic agents, drugs, and disease states which alter autonomic tone in either direction may modify the response. (Key words: Carbon dioxide: cardiovascular response; Heart: carbon dioxide; Arteries: carbon dioxide; Blood pressure: carbon dioxide.)

CARBON DIOXIDE induces direct and indirect circulatory effects in man and experimental animals. Acting directly,  $CO_2$  dilates peripheral arterioles and depresses myocardial contractility.<sup>1-5</sup> It also stimulates the central nervous system at several levels directly and at the vasomotor level indirectly by initiating afferent impulses from the peripheral chemoreceptors. Activation of the central nervous system evokes sympatho-

adrenal responses, resulting in increased myocardial contractility, tachycardia, and hypertension.<sup>3,5,6,7</sup> This paper reports the sum of these effects by detailing the circulatory responses of 41 awake human volunteers to administration of  $CO_2$  under carefully controlled circumstances. Control measurements (without added  $CO_2$ ) during both controlled and spontaneous respiration also serve as reference data for circulatory measurements in man.

### Methods

Forty-one healthy, young, adult, male volunteers were interviewed and informed consent was obtained. The procedures and consent form had been approved by the University of California and Stanford University Committees on Human Experimentation. All subjects fasted overnight and the studies were conducted the following morning.

Using local anesthesia, an arterial catheter was inserted percutaneously into the brachial or radial artery and a right atrial catheter was inserted through a 14-gauge needle into the basilic vein. In most cases, the catheter was advanced until a right ventricular pressure trace appeared and was then withdrawn until an atrial pressure wave was observed. A peripheral venous catheter was inserted in the forearm. Whitney strain gauges (mercury-filled plastic tubing) were placed about the fleshy portion of the forearm. Venous occlusion cuffs were placed at the base of the wrist and on the arm. A lead II electrocardiogram was used for recording heart rate. In 34 subjects, a precordial phonocardiogram and carotid pulse monitor were attached for determining intervals of electrical and mechanical systole. Each subject lay on an inflated air mattress which rested on an ultra-low-frequency air-bearing ballistocardiogram bed. The air mat-

\* Associate Professor, Department of Anesthesia, Massachusetts General Hospital and Harvard Medical School, Boston, MA 02114.

† Professor, Department of Anesthesia, University of California Medical Center, San Francisco, CA 94122.

Received from the Department of Anesthesia, University of California Medical Center, San Francisco, CA, and the Anesthesia Laboratories, Massachusetts General Hospital, Boston, MA 02114. Accepted for publication February 12, 1974. Supported in part by USPHS grant 5T1 GM00063-14 and USPHS 5P01 GM15571-05.

TABLE 1. Control Values (Means  $\pm$  1 SE)

	Controlled Respiration (n = 15)	Spontaneous Respiration (n = 26)
Age (years)	24.2 $\pm$ 0.8	24 $\pm$ 0.6
Height (cm)	175 $\pm$ 2	174 $\pm$ 1.4
Weight (kg)	70 $\pm$ 2.2	70.4 $\pm$ 1.4
P <sub>A</sub> O <sub>2</sub> (torr)	492* $\pm$ 13.6	525 $\pm$ 8.1
P <sub>A</sub> CO <sub>2</sub> (torr)	35.7 $\pm$ 1	37 $\pm$ 0.8
pH	7.39 $\pm$ 0.01	7.40 $\pm$ 0.001
Base excess (mEq/l)	-3.1* $\pm$ 0.5	-1.6 $\pm$ 0.3
Cardiac output (l/min)	4.64* $\pm$ 0.2	5.91 $\pm$ 0.2
Cardiac index (l/min/m <sup>2</sup> )	2.48* $\pm$ 0.1	3.15 $\pm$ 0.1
Stroke volume (ml)	78.7* $\pm$ 4	95.3 $\pm$ 3.6
Stroke index (ml/m <sup>2</sup> )	42.1* $\pm$ 1.6	51.1 $\pm$ 1.8
Heart rate (beats/min)	60 $\pm$ 1.7	62 $\pm$ 1.5
Mean arterial pressure (torr)	90 $\pm$ 2.6	88.2 $\pm$ 1.6
Mean right atrial pressure (torr)	3.6 $\pm$ 0.8	4.4 $\pm$ 0.5
Total peripheral resistance (dynes/sec/cm <sup>-5</sup> )	1,530* $\pm$ 78	1,174 $\pm$ 52
Ejection time (msec)	300 $\pm$ 7	289 $\pm$ 9
Mean rate ventricular ejection (ml/sec)	0.25 $\pm$ 0.01	0.32 $\pm$ 0.01
Left ventricular stroke work (kg-m)	0.10 $\pm$ 0.007	0.11 $\pm$ 0.005
Left ventricular work (kg-m/min)	6.04 $\pm$ 0.4	7.07 $\pm$ 0.4
Tension-time Index†	26.9 $\pm$ 0.1	24.9 $\pm$ 1.2
Ballistocardiogram IJ-wave amplitude	21.2 $\pm$ 2.9	14.6 $\pm$ 1.4
Forearm blood flow (ml/100 g tissue/min)	3.1 $\pm$ 0.3	3.0 $\pm$ 0.3
Forearm vascular resistance (dynes/sec/cm <sup>-4</sup> )	30.9 $\pm$ 3	34.3 $\pm$ 4
Forearm venous compliance (ml/torr)	0.16 $\pm$ 0.03	0.13 $\pm$ 0.02

\*  $P < 0.05$ , controlled respiration vs. spontaneous respiration.† AP  $\times$  ET  $\times$  0.001.

tress was deflated during ballistocardiographic and other cardiovascular measurements.

Mean arterial pressure (MAP), mean right atrial pressure (MRAP), and peripheral venous pressure (PVP) were transduced with Statham strain gauges. Duplicate cardiac outputs were obtained by dye dilution with indocyanine green using a Beckman Camo diodensitometer. Arterial P<sub>O<sub>2</sub></sub>, P<sub>CO<sub>2</sub></sub>, and pH were measured with blood-gas electrodes. Forearm blood flow was obtained by venous occlusion plethysmography with the Whitmer strain gauge.<sup>8</sup> The amplitude of the IJ wave of the ballistocardiogram was recorded. Oral and skin temperatures were measured with thermistors.

Calculated variables included stroke volume (SV), stroke index (SI), cardiac index (CI), total peripheral resistance (TPR), ejection time (ET), mean rate of left ventricular ejection (MRLVE), left ventricular stroke work (LVSF), left ventricular work (LVWF), forearm venous compliance (FVC), forearm vascular resistance (FVR), and base excess (BE).

Control measurements were obtained with the subject breathing oxygen from an anesthetic circle system via a mouthpiece. The nose was occluded by nose clips. After the subjects had become accustomed to the breathing system, ventilation of 15 of the 41 subjects was changed from spontaneous to controlled with a volume-limited ventilator. There was no sign that the subjects resisted ventilation over the range of CO<sub>2</sub> studied. End-tidal P<sub>CO<sub>2</sub></sub> was monitored with an infrared CO<sub>2</sub> analyzer and maintained at normal levels. The control (no added CO<sub>2</sub>) data for the 15 subjects whose ventilation was controlled were compared with the data from the 26 subjects whose ventilation was spontaneous at the same P<sub>ACO<sub>2</sub></sub>, using unpaired t tests.  $P < 0.05$  was accepted as statistically significant.

After control measurements had been obtained, CO<sub>2</sub> was added in incremental amounts to the breathing system to produce 2-4-torr stepwise increases in P<sub>ACO<sub>2</sub></sub>. All measurements were repeated after equilibrating for 6 minutes at each level of P<sub>ACO<sub>2</sub></sub>. At the end of 6 minutes, an arterial sample was

TABLE 2. Circulatory Responses to CO<sub>2</sub> (Means  $\pm$  1 SE)\*

	Controlled Respiration		Spontaneous Respiration	
	Regression Equation	Per Cent Change with 10-torr Increase in PaCO <sub>2</sub>	Regression Equation	Per Cent Change with 10-torr Increase in PaCO <sub>2</sub>
Cardiac output (l/min)	$y = 0.17x - 2.2$	39 $\pm$ 5	$y = 0.17x - 1.3$	32 $\pm$ 4
Cardiac index (l/min/m <sup>2</sup> )	$y = 0.09x - 0.8$	39 $\pm$ 5	$y = 0.10x - 0.6$	32 $\pm$ 4
Stroke volume (ml)	$y = 0.62x + 55.6$	10 $\pm$ 2	$y = 0.34x + 83.3$	11 $\pm$ 2
Stroke index (ml/m <sup>2</sup> )	$y = 0.34x + 29.7$	11 $\pm$ 2	$y = 0.23x + 44.4$	11 $\pm$ 2
Heart rate (beats/min)	$y = 1.74x - 4.8$	28 $\pm$ 4	$y = 1.7x - 1.8$	26 $\pm$ 3
Mean arterial pressure (torr)	$y = 0.82x + 60.9$	9 $\pm$ 1	$y = 0.89x + 50.4$	10 $\pm$ 2
Mean right atrial pressure (torr)	$y = -0.05x + 5.3$	0	$y = -0.08x + 7.4$	0
Total peripheral resistance (dynes/cm <sup>2</sup> )	$y = -28.7x + 2,540$	-17 $\pm$ 4	$y = -16.7x + 1,800$	-14 $\pm$ 3
Ejection time (msec)	$y = -0.25x + 309.1$	-0.54 $\pm$ 0.77	$y = -1.16x + 336$	-3 $\pm$ 2
Mean rate ventricular ejection (ml/sec)	$y = 0.0008x + 0.3$	3 $\pm$ 4	$y = 0.0025x + 0.2$	8 $\pm$ 3
Left ventricular stroke work (kg-m)	$y = 0.001x + 0.06$	12 $\pm$ 3	$y = 0.002x + 0.04$	25 $\pm$ 10
Left ventricular work (kg-m/min)	$y = 0.33x - 6.4$	54 $\pm$ 11	$y = 0.31x - 4$	46 $\pm$ 5
Tension-time index	$y = 0.25x + 18$	10 $\pm$ 2	$y = 0.11x + 22$	65 $\pm$ 2
Forearm blood flow (ml/100 g tissue/min)	$y = 0.1x - 0.1$	30 $\pm$ 11	$y = 0.12x - 1.1$	44 $\pm$ 10
Forearm vascular resistance (dynes/cm <sup>2</sup> )	$y = -0.13x + 37.5$	-6 $\pm$ 9	$y = -0.71x + 64$	-10 $\pm$ 10
Forearm venous compliance (ml/torr)	$y = 0.0009x + 0.14$	0	$y = 0.0014x + 0.08$	0

\* There was no significant difference comparing the circulatory responses to CO<sub>2</sub> during spontaneous respiration with those during controlled respiration.

obtained and the PaCO<sub>2</sub> value was used in the regression analyses described below. Measurements at three separate levels of PaCO<sub>2</sub> were obtained for each subject. The cardiovascular response to CO<sub>2</sub> was analyzed by linear regression, again comparing the 15 subjects during controlled respiration with the 26 subjects during spontaneous respiration. A regression line was calculated for each individual subject, with PaCO<sub>2</sub> always serving as the independent variable. The individual slopes and y intercepts were combined into two groups, controlled respiration and spontaneous respiration. The mean and standard error were calculated for the slope and y intercept for each group. The mean slope values for controlled respirations were compared statistically with the mean slope values for spontaneous respiration by unpaired t tests. Additionally, for each individual subject, the percentage change (mean and standard error) for each measured value in response to a 10-torr increase in PaCO<sub>2</sub> was calculated (table 2). The initial value measured prior to CO<sub>2</sub> challenge served as the baseline value.

## Results

### CONTROL DATA (TABLE 1)

The control measurements during both controlled respiration (n = 15) and spontaneous respiration (n = 26) showed that the subjects were equivalent with respect to age, height, weight, PaCO<sub>2</sub>, and pH. However, cardiac output and index, stroke volume and index, and PaO<sub>2</sub> were significantly lower, while TPR was significantly higher, during controlled respiration than during spontaneous respiration. A slight but statistically significant metabolic acidosis was present during controlled respiration.

### CIRCULATORY RESPONSES TO CO<sub>2</sub> (TABLE 2)

The mean PaCO<sub>2</sub> at which the regression lines began was 38.7  $\pm$  0.3 torr (1 SE). The mean PaCO<sub>2</sub> at which the last measurement was obtained was 50.2  $\pm$  0.9 torr. The mean increase in PaCO<sub>2</sub> was 11.5  $\pm$  0.9 torr. Given this hypercarbia, several circulatory variables (cardiac output, cardiac index, stroke volume,

TABLE 3. Effects of anesthetics on the Cardiac-index Response to CO<sub>2</sub> in Man\*

	Response (Δ ml/min Increase in Paco <sub>2</sub> )
Conscious	95
Cyclopropane, 15–20 per cent (alveolar) <sup>16</sup>	76
Cyclopropane, 25–30 per cent (alveolar) <sup>16</sup>	51
Halothane, 0.8 per cent (alveolar) <sup>17</sup>	34
Halothane, 1 per cent (inspired) <sup>16</sup>	23
N <sub>2</sub> O–meperidine–d-tubocurarine <sup>18</sup>	22
Fluroxene, 5 per cent (alveolar) <sup>20</sup>	59
Fluroxene, 9 per cent (alveolar) <sup>20</sup>	65
Isoflurane, 1.2 per cent (alveolar) <sup>21</sup>	75
Isoflurane, 1.8 per cent (alveolar) <sup>21</sup>	48
Enflurane, 1.9 per cent (alveolar) <sup>22</sup>	104
Spinal anesthesia (T1 level)†	24

\* Calculated milliliter increase in cardiac index per torr increase in P<sub>a</sub>CO<sub>2</sub>.

† Bonica, J. J.: Personal communication.

stroke index, mean arterial pressure, left ventricular stroke work, left ventricular work, forearm blood flow and heart rate) were significantly increased, while one (peripheral resistance) decreased.

The slopes of the various CO<sub>2</sub> response curves were similar during controlled and spontaneous respiration. For example, although the control cardiac output was lower during controlled respiration, the responsiveness of cardiac output to CO<sub>2</sub> administration was not different.

## Discussion

Administration of CO<sub>2</sub> during either controlled or spontaneous respiration significantly increased cardiac output and indices of myocardial contractility, while peripheral vascular resistance decreased. Hypercarbia acts through central nervous system stimulation to increase sympathetic tone, thereby increasing myocardial contractility and vascular tone.<sup>2,5,6,7,9,10</sup> Peripherally, however, the indirect effects of vasoconstriction are overcome by the direct peripheral vasodilating effect of CO<sub>2</sub>, so that total peripheral resistance decreases. Our data

quantitate these well-known effects of CO<sub>2</sub> in young healthy human subjects.

Just as anesthetics, narcotics, and disease states may modify the ventilatory response to CO<sub>2</sub>, so also may the circulatory responses to CO<sub>2</sub> be modified. For example, anesthetics depress the normal increase in cardiac index produced by elevation of P<sub>a</sub>CO<sub>2</sub> (table 3). The ability of various anesthetics to modify this response probably relates to suppression of sympathetic activity imposed by the anesthetics. For example, halothane<sup>11</sup> reduces peripheral ganglionic sympathetic activity in the intact cat and greatly attenuates the circulatory response to CO<sub>2</sub>. Conversely, cyclopropane, and fluroxene<sup>13</sup> induce central sympathetic activity, and both agents are associated with vigorous, albeit slightly blunted, cardiovascular response to increased CO<sub>2</sub> levels. Where the other anesthetics fit into this scheme is not well established.

Sympathetic blockade should reduce the circulatory response to CO<sub>2</sub>. During total spinal anesthesia in volunteers, the cardiac index response to exogenously administered CO<sub>2</sub> was markedly diminished, although slight response did occur,† possibly through inhibition of vagal activity. Undoubtedly, as other drugs and disease states are studied, further interference with normal circulatory responses to CO<sub>2</sub> will be demonstrated.

The differences between controlled respiration and spontaneous respiration for the control baseline data support previous findings.<sup>14,15</sup> Impedance of venous return during controlled respiration results in increased peripheral vascular tone and maintenance of blood pressure in the face of reduced cardiac output.

The authors gratefully acknowledge the technical assistance of Linda Ferrara, R.N.

## References

1. Blackburn JP, Conway CM, Leigh JH, et al: Myocardial carbon dioxide response curves. *Br J Anaesth* 42:599, 1970
2. Butcher RL, Bitensky L, Chayen J: Left ventricular performance in the conscious dog with chronically denervated heart. *Cardiovasc Res* 6:457–475, 1972

† Bonica, J. J.: Personal communication.

3. Tenney SM, Lamb TW: Physiological consequences of hypoventilation and hyperventilation, *Handbook of Physiology*. Section 3, Respiration, volume II. Edited by WO Fenn, H Rahn. Washington D.C., American Physiological Society, 1965, chapter 37
4. Nejad NS, Ogden E: Effect of blood pH and CO<sub>2</sub> tension on performance of the heart lung preparation. *Proc Soc Exp Biol Med* 126:771-776, 1967
5. Price HL: Effect of carbon dioxide on the cardiovascular system. *ANESTHESIOLOGY* 21:652-662, 1960
6. Cross BA, Silver IA: Central activation of sympathoadrenal system by hypoxia and hypercapnia. *J Endocrinol* 24:91-103, 1962
7. Downing SE, Siegal JH: Baroreceptor and chemoreceptor influences on sympathetic discharge to the heart. *Am J Physiol* 204:471-479, 1963
8. Whitney RJ: The measurement of volume changes in the human limb. *J Physiol* 121:1, 1953
9. Millar RA, Biscoe TJ: Postganglionic sympathetic discharge and the effect of inhalation anesthetics. *Br J Anaesth* 38:92-114, 1966
10. Norman J, Atkinson SA: The effect of cardiac sympathetic blockade on the relationship between cardiac output and carbon dioxide tension in anesthetized dogs. *Br J Anaesth* 42:592-602, 1970
11. Skovsted T, Price ML, Price HL: The effects of halothane on arterial pressure, pre-ganglionic sympathetic activity, and barostatic reflexes. *ANESTHESIOLOGY* 31:507-514, 1969
12. Price HL, Warden JC, Cooperman LH, et al: Central sympathetic excitation caused by cyclopropane. *ANESTHESIOLOGY* 30:426-438, 1969
13. Skovsted P, Price HL: Central sympathetic excitation caused by fluorene. *ANESTHESIOLOGY* 32:210-217, 1970
14. Feisal KA, Abboud FM, Eckstein J: Effects of adrenergic blockade on cardiovascular responses to airway pressure. *Am J Physiol* 213:127-133, 1967
15. Morgan BC, Martin WE, Hornbein TF, et al: Hemodynamic effects of intermittent positive-pressure respiration. *ANESTHESIOLOGY* 27:584-590, 1966
16. Cullen DJ, Eger EI II, Gregory GA: The cardiovascular effects of carbon dioxide in man, conscious and during cyclopropane anesthesia. *ANESTHESIOLOGY* 31:407-413, 1969
17. Hornbein TF, Martin WE, Bonica JJ, et al: Nitrous oxide effects of the circulatory and ventilatory responses to halothane. *ANESTHESIOLOGY* 31:250-260, 1969
18. Prys-Roberts C, Kelman GR, Kain C, et al: Cardiac output and carbon dioxide levels during halothane anesthesia in man. *Br J Anaesth* 39:687-688, 1967
19. Prys-Roberts C, Kelman GR, Kain C, et al: Circulatory influences of artificial ventilation during nitrous oxide anesthesia in man. II. Results: The relative influences of mean intrathoracic pressure and arterial carbon dioxide tension. *Br J Anaesth* 39:533-547, 1967
20. Cullen BF, Eger EI II, Smith NT, et al: The circulatory response to hypercapnia during fluorene anesthesia in man. *ANESTHESIOLOGY* 34:415-420, 1971
21. Cromwell TH, Stevens WC, Eger EI II, et al: The cardiovascular effect of compound 469 (Forane) during spontaneous ventilation and CO<sub>2</sub> challenge in man. *ANESTHESIOLOGY* 35:17-25, 1971
22. Marshall BJ, Klingenstein CH, Neigh JL, et al: Some pulmonary and cardiovascular effects of enflurane (Ethrane) anesthesia with varying PaCO<sub>2</sub> in man. *Br J Anaesth* 43:996-1002, 1971

## Drugs and Their Actions

**SCOPOLAMINE DELIRIUM** The effects of intravenously administered physostigmine on scopolamine-induced confusion and amnesia were studied in the parturient patient. Physostigmine completely reversed the central depression and amnesia produced by scopolamine. All patients became cooperative and oriented within 3 to 5 minutes and had good antegrade memory after administration of the drug. The effectiveness of physostigmine in this preliminary study warrants more definitive studies of the maternal and fetal effects of the drug. (Smiler, B., Bartholomew, E.G., Sivak, B.J., and others: *Physostigmine Reversal of Scopolamine Delirium in Obstetric Patients*. *Am J Obstet Gynecol* 115: 326-329, 1973.)