

```

%root_path = fileparts(mfilename("fullpath"));
%root_path = pwd;

% retrieve selected cuts used for further fitting
[GH_FFCutsFits,CutFitFileGH] =
recover_selected_cuts('GH_FFCutsFits','e:\SHARE\Fe\Data\analysis\07_fit_with_multicu
t_all\sel_cuts');

```

```

fn = fieldnames(GH_FFCutsFits);
%nf = 11
%for nf = 2:numel(fn)
%selectCuts = fn{nf};
selectCuts = 'Ei200_off110GHmaxFF';
if ismember(selectCuts(1:3),{'msk','dat'})||...
    contains(selectCuts,'all_fit_par')
    %continue
    error('Can not fit field: %s',selectCuts);
end

fprintf('*****\n');

```

```

*****

```

```

fprintf('Processing cuts: %s \n', selectCuts);

```

```

Processing cuts: Ei200_off110GHmaxFF

```

```

fprintf('*****\n');

```

```

*****

```


```

cuts_list = GH_FFCutsFits.(selectCuts);
if ~iscell(cuts_list)
    cuts_list = {cuts_list};
end
if contains(selectCuts,'Ei200')
    min_Ei = 30;
    mi = maps_instrument(200,600,'S');
end
if contains(selectCuts,'Ei400')
    min_Ei = 40;
    mi = maps_instrument(401,600,'S');
end
if contains(selectCuts,'Ei800')
    min_Ei = 70;
    mi = maps_instrument(786,600,'S');
end
sample=IX_sample(true,[1,0,0],[0,1,0],'cuboid',[0.04,0.03,0.02]);
for j=1:numel(cuts_list)

```

```
cuts_list{j} = cuts_list{j}.set_instrument(mi);
cuts_list{j} = cuts_list{j}.set_sample(sample);
plot(cuts_list{j}); lz 0 1; keep_figure;
```

end



```
dir_name = "GH";

dE_step = 2; %original energy transfer step data were binned to. No point in going
finer
half_dE = 5; % half width of data binning


en_mm = cuts_list{1}.data.img_range(:,4);
en_mm(1) = max(min_Ei,en_mm(1));
%en_mm(1) = 100;
%en_mm(2) = 170;
cut_en = en_mm(1)+half_dE:2*half_dE:(en_mm(2)-half_dE);

cut_dimensions = 1;
GH_FFCutsFits.(selectCuts) = cuts_list;

GH_FFCutsFits = fit_multicutsFF_along_direction(...
    GH_FFCutsFits,selectCuts,cut_dimensions,cut_en,dE_step,half_dE,true);
```


```
*****
**** En = 35±5 going to 115
*****
```

Fe_ei200_align_sym3.sqw
 $-0.1 \leq \xi \leq 0.1$ in $[1-\xi, 1, 0]$, $-0.1 \leq \eta \leq 0.1$ in $[1, 1, \eta]$, $30 \leq E \leq 40$ eV
 $\zeta = -0.005:0.01:0.605$ in $[1, 1+\zeta, 0]$




 **** En = 45±5 going to 115

Fe_ei200_align_sym3.sqw
 $S=0.82 \pm 0.038$; $J_0= 25 \pm 0.5$; $J_1= 0 \pm 0$; $\gamma=6.36957 \pm 0.286903$
 $-0.1 \leq \xi \leq 0.1$ in $[1-\xi, 1, 0]$, $-0.1 \leq \eta \leq 0.1$ in $[1, 1, \eta]$, $40 \leq E \leq 50$ eV
 $\zeta = -0.005:0.01:0.605$ in $[1, 1+\zeta, 0]$




Fe_ei200_align_sym3.sqw
 $-0.1 \leq \xi \leq 0.1$ in $[1-\xi, 1, 0]$, $-0.1 \leq \eta \leq 0.1$ in $[1, 1, \eta]$, $40 \leq E \leq 50$ eV
 $\zeta = -0.005:0.01:0.605$ in $[1, 1+\zeta, 0]$



 **** En = 55±5 going to 115


Fe_ei200_align_sym3.sqw

S=0.79±0.037; J0= 25±0.55; J1= 0± 0; gamma=12.3001±0.504865
 $-0.1 \leq \xi \leq 0.1$ in $[1-\xi, 1, 0]$, $-0.1 \leq \eta \leq 0.1$ in $[1, 1, \eta]$, $50 \leq E \leq 100$ eV
 $\zeta = -0.005:0.01:0.605$ in $[1, 1+\zeta, 0]$



Fe_ei200_align_sym3.sqw


S=0.79±0.037; J0= 25±0.55; J1= 0± 0; gamma=12.3001±0.504865
 $-0.1 \leq \xi \leq 0.1$ in $[1-\xi, 1, 0]$, $-0.1 \leq \eta \leq 0.1$ in $[1, 1, \eta]$, $50 \leq E \leq 100$ eV
 $\zeta = -0.005:0.01:0.605$ in $[1, 1+\zeta, 0]$




 **** En = 65±5 going to 115

Fe_ei200_align_sym3.sqw

S=0.97±0.091; J0= 26±0.57; J1= 0± 0; gamma=22.6668±3.28226
 $-0.1 \leq \xi \leq 0.1$ in $[1-\xi, 1, 0]$, $-0.1 \leq \eta \leq 0.1$ in $[1, 1, \eta]$, $60 \leq E \leq 100$ eV
 $\zeta = -0.005:0.01:0.605$ in $[1, 1+\zeta, 0]$




Fe_ei200_align_sym3.sqw
 $-0.1 \leq \xi \leq 0.1$ in $[1-\xi, 1, 0]$, $-0.1 \leq \eta \leq 0.1$ in $[1, 1, \eta]$, $60 \leq E \leq 70$ in $[1, 1+\zeta, 0]$
 $\zeta = -0.005:0.01:0.605$ in $[1, 1+\zeta, 0]$




 **** En = 75±5 going to 115

Fe_ei200_align_sym3.sqw
 $S=0.85 \pm 0.034$; $J_0= 28 \pm 0.38$; $J_1= 0 \pm 0$; $\gamma=21.6565 \pm 0.814554$
 $-0.1 \leq \xi \leq 0.1$ in $[1-\xi, 1, 0]$, $-0.1 \leq \eta \leq 0.1$ in $[1, 1, \eta]$, $70 \leq E \leq 85$ in $[1, 1+\zeta, 0]$
 $\zeta = -0.005:0.01:0.605$ in $[1, 1+\zeta, 0]$




Fe_ei200_align_sym3.sqw
 $-0.1 \leq \xi \leq 0.1$ in $[1-\xi, 1, 0]$, $-0.1 \leq \eta \leq 0.1$ in $[1, 1, \eta]$, $70 \leq E \leq 85$ in $[1, 1+\zeta, 0]$
 $\zeta = -0.005:0.01:0.605$ in $[1, 1+\zeta, 0]$



 **** En = 85±5 going to 115


Fe_ei200_align_sym3.sqw

S= 1.1 ± 0.088 ; J0= 29 ± 0.57 ; J1= 0 ± 0 ; gamma= 40.0281 ± 4.38424
 $-0.1 \leq \xi \leq 0.1$ in $[1-\xi, 1, 0]$, $-0.1 \leq \eta \leq 0.1$ in $[1, 1, \eta]$, $80 \leq E \leq 100$ eV
 $\zeta = -0.005:0.01:0.605$ in $[1, 1+\zeta, 0]$



Fe_ei200_align_sym3.sqw


$-0.1 \leq \xi \leq 0.1$ in $[1-\xi, 1, 0]$, $-0.1 \leq \eta \leq 0.1$ in $[1, 1, \eta]$, $80 \leq E \leq 100$ eV
 $\zeta = -0.005:0.01:0.605$ in $[1, 1+\zeta, 0]$




**** En = 95 ± 5 going to 115

Fe_ei200_align_sym3.sqw

S= 1 ± 0.079 ; J0= 30 ± 0.59 ; J1= 0 ± 0 ; gamma= 42.8955 ± 4.59022
 $-0.1 \leq \xi \leq 0.1$ in $[1-\xi, 1, 0]$, $-0.1 \leq \eta \leq 0.1$ in $[1, 1, \eta]$, $90 \leq E \leq 110$ eV
 $\zeta = -0.005:0.01:0.605$ in $[1, 1+\zeta, 0]$




Fe_ei200_align_sym3.sqw
 $-0.1 \leq \xi \leq 0.1$ in $[1-\xi, 1, 0]$, $-0.1 \leq \eta \leq 0.1$ in $[1, 1, \eta]$, $90 \leq E \leq 1$
 $\zeta = -0.005:0.01:0.605$ in $[1, 1+\zeta, 0]$




 **** En = 105±5 going to 115

Fe_ei200_align_sym3.sqw
 $S = 1.1 \pm 0.089$; $J_0 = 31 \pm 0.73$; $J_1 = 0 \pm 0$; $\gamma = 57.0849 \pm 6.48893$
 $0.1 \leq \xi \leq 0.1$ in $[1-\xi, 1, 0]$, $-0.1 \leq \eta \leq 0.1$ in $[1, 1, \eta]$, $100 \leq E \leq$
 $\zeta = -0.005:0.01:0.605$ in $[1, 1+\zeta, 0]$




Fe_ei200_align_sym3.sqw
 $0.1 \leq \xi \leq 0.1$ in $[1-\xi, 1, 0]$, $-0.1 \leq \eta \leq 0.1$ in $[1, 1, \eta]$, $100 \leq E \leq$
 $\zeta = -0.005:0.01:0.605$ in $[1, 1+\zeta, 0]$





 **** En = 115±5 going to 115



Warning: Dataset:2 points with zero, negative or undefined (infinite or NaN) error bars have been removed from fit, which is 3.2787

Fe_ei200_align_sym3.sqw
 $S = 1.1 \pm 0.13$; $J_0 = 32 \pm 0.85$; $J_1 = 0 \pm 0$; $\gamma = 64.1732 \pm 8.94883$
 $0.1 \leq \xi \leq 0.1$ in $[1-\xi, 1, 0]$, $-0.1 \leq \eta \leq 0.1$ in $[1, 1, \eta]$, $110 \leq E \leq$
 $\zeta = -0.005:0.01:0.605$ in $[1, 1+\zeta, 0]$



Fe_ei200_align_sym3.sqw
 $0.1 \leq \xi \leq 0.1$ in $[1-\xi, 1, 0]$, $-0.1 \leq \eta \leq 0.1$ in $[1, 1, \eta]$, $110 \leq E \leq$
 $\zeta = -0.005:0.01:0.605$ in $[1, 1+\zeta, 0]$






```

c1 = GH_FFCutsFits.(selectCuts);
GH_FFCutsFits.(selectCuts) = [c1{:}];


beep
beep
beep

all_fit_par = GH_FFCutsFits.([selectCuts,'_all_fit_par']);
bg_data = extract_bg_par(all_fit_par,[1,2]);
% original slope parameters
pd(bg_data(1));lx(all_fit_par(1).en,all_fit_par(end).en);keep_figure;

```



```
pd(bg_data(2));lx(all_fit_par(1).en,all_fit_par(end).en);keep_figure
```



```
%pd(bg_data(3));lx(fit_res.peak_fit_par(1).en,fit_res.peak_fit_par(end).en);keep_figure;
%pd(bg_data(4));lx(fit_res.peak_fit_par(1).en,fit_res.peak_fit_par(end).en);keep_figure
```

```
doc = matlab.desktop.editor.getActive;
doc.save;
thisFile = 'e:\SHARE\Fe\Data\analysis\06_fit_with_AQ0\fit_cutFe_GH.m';
targ_file = fullfile(fileparts(thisFile),['fit_cutFe_GH_',selectCuts,'.pdf']);
export(thisFile,targ_file);
```

```
fit_src_struct = GH_FFCutsFits;
save(CutFitFileGH,"fit_src_struct",-v7.3');
close all;
%end
```