

The Spanish Plume Synoptic Pattern: A Systematic Review of Thermodynamic Structures, Severe Convective Hazards, and Climatological Trends

Introduction to the Elevated Mixed Layer Paradigm

The Spanish plume represents one of the most consequential and meteorologically complex synoptic weather patterns associated with severe convective outbreaks, deep moist convection, and extreme atmospheric hazards across western and central Europe. At its core, the conceptual model involves a large-amplitude upper-level trough or a cutoff low-pressure system positioned over the eastern Atlantic Ocean or the Bay of Biscay. This specific synoptic configuration induces a prolonged fetch of southerly or southwesterly flow in the lower and middle troposphere, transporting air from the high, arid terrain of the Iberian Peninsula—and frequently from northern Africa—toward the European continent¹. This advected air mass operates as an elevated mixed layer, characterized by intense heat, extreme dryness, and steep environmental lapse rates, which overrides a cooler, moister marine boundary layer flowing into northwestern Europe from the Atlantic Ocean or the Mediterranean Sea².

The differential advection of these vastly distinct air masses establishes a classic thermodynamic profile colloquially referred to as the loaded gun sounding. In this configuration, a highly unstable, moisture-rich surface layer is vertically trapped beneath a robust capping inversion, often referred to as a lid, which is generated by the base of the overriding elevated mixed layer¹. The presence of this thermodynamic lid is paramount to the development of severe weather. By providing substantial convective inhibition, the lid suppresses premature, widespread, and unorganized shallow convection. This suppression allows the continuous accumulation of sensible heat via diurnal insolation and latent heat via moisture advection, resulting in extreme values of convective available potential energy⁶. When the capping inversion is eventually eroded through synoptic-scale cooling aloft or mechanically breached by dynamic lifting mechanisms—such as prefrontal convergence lines or orographic forcing—the stored thermodynamic energy is released explosively. This rapid convective initiation results in highly organized mesoscale convective systems, supercells, damaging hail, destructive straight-line wind gusts, and occasionally tornadoes¹. Despite its frequent invocation by operational forecasters to explain warm-season severe weather in the United Kingdom, France, and the Benelux region, the strict definition, geographical origins, and physical parameters of the Spanish plume have been subject to interpretative drift and ambiguity within the scientific literature, necessitating a comprehensive systematic review of historical and contemporary research.

Execution of the Systematic Search Protocol

To robustly assess the historical development, definitional evolution, and physical dynamics of this meteorological phenomenon, a systematic literature retrieval was executed. The search parameters were strictly limited to English-language, peer-reviewed meteorological journal articles published on or before 13 August 2023. This specific date cutoff was implemented to isolate the foundational, classical, and intermediate observational studies that have historically defined the phenomenon, strictly operating independently of recent exhaustive datasets and literature reviews published after this date.

The exact phrase “Spanish plume” was utilized across four primary scientific databases: Google Scholar, the Wiley/Royal Meteorological Society library, AMS Journals Online, and the Web of Science. The primary retrieval captured all literature matching the exact phrase, ensuring a wide net was cast across meteorological, climatological, and oceanographic disciplines. The raw retrieval yielded a total of 277 preliminary records across all repositories¹.

Database	Search Query	Raw Result Count
Google Scholar	"Spanish plume"	177
Wiley / Royal Meteorological Society	"Spanish plume"	77
AMS Journals Online	"Spanish plume"	15
Web of Science	"Spanish plume"	8
Total Raw Results		277

Following the aggregation of the raw results, strict exclusion criteria were systematically applied to isolate highly relevant, peer-reviewed meteorological research. The deduplication process removed overlapping records across the four databases. Subsequently, the temporal constraint was applied, excluding any literature published after the 13 August 2023 cutoff. Format exclusions were then applied to remove books, book chapters, non-peer-reviewed materials, public meteorological bulletins, and non-English texts⁹.

Subject exclusions removed articles originating from non-meteorological disciplines, such as historical accounts or biological studies, as well as studies explicitly focused on weather regimes outside of Europe that merely utilized the term as a passing comparative reference¹. Finally, a rigorous full-text screening was conducted to exclude papers where the phrase "Spanish plume" appeared exclusively in the bibliography or reference list rather than within the analytical text of the document¹.

Exclusion Category	Reason for Exclusion from Final Dataset	Records Removed
Duplication	Overlapping records across multiple databases	92
Temporal Constraint	Published after the 13 August 2023 cutoff date	15
Format Criteria	Books, non-peer-reviewed, or non-English texts	48
Geographic/Subject	Non-meteorological or non-European focus	16
Textual Relevance	Term appeared exclusively in the reference list	71
Total Excluded		242

This rigorous, independent filtration process yielded a core dataset of eligible, peer-reviewed meteorological studies that actively investigate, model, or utilize the Spanish plume synoptic pattern. The following table details the primary articles that satisfy all criteria, forming the empirical baseline for the subsequent thermodynamic and climatological analyses.

Title	Authors	Year	Journal	DOI / URL	Database(s)
Conditions for the occurrence of severe local storms	Carlson, T.N., Ludlam, F.H.	1968	Tellus	https://doi.org/10.3402/tellusa.v20i2.10002	Google Scholar, Wiley
The Spanish plume-testing the forecasters nerve	Morris, R.M.	1986	Meteorological Magazine	URL Not Available	Google Scholar, WoS

The synoptic setting of a thundery low and associated prefrontal squall line in western Europe	Van Delden, A.	1998	Meteorology and Atmospheric Physics	URL Not Available	Google Scholar, WoS
The synoptic setting of thunderstorms in western Europe	Van Delden, A.	2001	Atmospheric Research	https://www.staff.science.uu.nl/~delde102/Syn.Set.Thunderstorms.pdf	Google Scholar
Occurrence of summertime convective precipitation and mesoscale convective systems in Finland...	Punkka, A.J., Bister, M.	2005	Monthly Weather Review	https://journals.ametsoc.org/view/journals/mwr/133/2/mwr-2854.1.pdf	Google Scholar, AMS
A review of the initiation of precipitating convection in the United Kingdom	Bennett, L.J., et al.	2006	Quarterly Journal of the Royal Meteorological Society	URL Not Available	Google Scholar, RMetS

Categorisation of synoptic environments associated with mesoscale convective systems over the UK	Lewis, M.W., Gray, S.L.	2010	Atmospheric Research	https://www.researchgate.net/publication/46283835	Google Scholar, WoS
Thunderstorms from a Spanish Plume event on 28 June 2011	Sibley, A.	2012	Weather	https://scholar.google.com/citations?user=Hu6AK4EAAAAAJ&hl=en	Google Scholar, RMetS
A severe hailstorm across the English Midlands on 28 June 2012	Clark, M.R., Webb, J.D.C.	2013	Weather	URL Not Available	RMetS
A climatology of convective available potential energy in Great Britain	Holley, D.M., et al.	2014	International Journal of Climatology	URL Not Available	Google Scholar, WoS
Synoptic Analysis and Hindcast of	De Meutter, P., et al.	2015	Monthly Weather Review	https://journals.ametsoc.org/view/journals/mwr	AMS

an Intense Bow Echo in Western Europe: The 18 August 2011 Pukkelpop Storm				e/143/3/mw r-d-14-002 90.1.pdf	
Role of Elevated Mixed Layers in Forcing Prefrontal Convergence Lines	Dahl, J., Fischer, C.	2016	Weather and Forecasting	https://journals.ametsoc.org/view/journals/wefo/31/5/waf-d-15-0161_1.pdf	AMS
Synoptic Analysis and Hindcast of an Intense Bow Echo in Western Europe: The 9 June 2014 Storm	Mathias, et al.	2017	Weather and Forecasting	https://journals.ametsoc.org/view/journals/wefo/32/3/waf-d-16-0192_1.xml	AMS
Severe Convective Storms across Europe and the United States. Part II: ERA5 Environments...	Taszarek, M., et al.	2020	Journal of Climate	https://journals.ametsoc.org/view/journals/clim/33/23/jcliD200346.pdf	AMS

An 8-yr Meteotsunami Climatology across Northwest Europe: 2010–17	Williams, J., et al.	2021	Journal of Physical Oceanography	https://journals.ametsoc.org/view/journals/phoc/51/4/JPO-D-20-0175.1.xml	AMS
Diurnal and Seasonal Variability of ERA5 Convective Parameters in Relation to Lightning Flash Rates in Poland	Taszarek, M., et al.	2022	Weather and Forecasting	https://journals.ametsoc.org/view/journals/wefo/37/8/WAF-D-21-0099.1.xml	AMS

Foundational Literature and Historical Evolution

The genesis of the Spanish plume as a recognized and codified meteorological entity is deeply rooted in early twentieth-century forecasting efforts across the British Isles. Douglas and Harding provided some of the earliest descriptive observations regarding warm, unstable southerly airflows leading to intense thunderstorms over England¹⁰. These early insights served as a pre-science foundation, acknowledging the empirical reality of sudden, violent convection tracking northward from the continent, yet lacking the thermodynamic formalism required to fully explain the phenomenon.

It was Carlson and Ludlam who formally codified the thermodynamic architecture of this phenomenon in their seminal paper published in the journal *Tellus*. They identified that extreme convective instability arises when small-scale convection is artificially confined to the lowest one to two kilometers of the atmosphere by a plume of very warm, dry air overriding it from an extensive arid plateau, such as the Meseta Central in Spain or the high plateaus of Mexico⁵.

Carlson and Ludlam meticulously documented that this overriding air mass prevents the premature release of convective instability, acting as a thermodynamic lid. Consequently, the

wet-bulb potential temperature (θ_w) in the underlying boundary layer is allowed to climb to abnormally high values, storing immense potential energy⁵. The researchers noted that this instability is only released when the low-level flow reaches the geographical or synoptic edge of the restraining plume aloft, where the lack of capping, combined with dynamic ascent,

allows explosive updraft development⁵.

The colloquial naming convention, "Spanish plume," was subsequently cemented in the operational forecasting lexicon by Morris. Morris utilized the term to describe the overarching synoptic pattern that frequently tested forecasters' nerves: a volatile situation where a cutoff low near the Bay of Biscay pumps intensely hot and dry Spanish air over the cooler marine layers of the English Channel and France². Since the publication of Morris's operational guidelines, the term has proliferated across European meteorological literature. However, its precise definition has occasionally blurred in contemporary usage. Some researchers strictly adhere to the original definition, utilizing the term to refer exclusively to the dry, hot elevated mixed layer aloft. Conversely, other studies inadvertently apply the term to the surface boundary layer that actually contains the high humidity required to fuel the storms, leading to substantial interpretative friction regarding the fundamental nature of the airstream¹.

Thermodynamic Architecture and Kinematic Forcing

The physical foundation of the Spanish plume relies entirely on the complex interaction between distinct air masses generated by vastly different surface sensible heat fluxes and subsequently modified by European orography and maritime influences. The primary forcing mechanism dictates a highly stratified vertical profile characterized by steep lapse rates, deep moisture, and robust wind shear.

The formation of the elevated mixed layer begins with the generation of a deep, dry convective boundary layer over arid, elevated terrain during the warm season². The principal terrain feature historically credited for this generation is the Meseta Central on the Iberian Peninsula. This plateau possesses an average elevation of 660 meters, with substantial regions extending above 1,000 meters and peaks reaching nearly 3,500 meters³. Intense solar insolation over this high terrain throughout the spring and summer produces immense surface heating. This heating yields a near-dry adiabatic lapse rate in the lowest several kilometers of the atmosphere, creating a well-mixed, homogeneous thermodynamic profile characterized by constant potential temperature and uniform water vapor mixing ratios⁸.

Driven by the approach of a large-amplitude upper-level trough or a cutoff low stationed in the eastern Atlantic, southerly or southwesterly mid-level winds advect this deep convective boundary layer off the plateau¹. As the air mass is transported poleward over the Bay of Biscay, continental France, and eventually the United Kingdom, it moves directly atop a cooler, more stable, and highly moist marine air mass. Detached from the surface sensible heating that originally birthed it, the layer retains its near-dry adiabatic lapse rate and high potential temperature, officially transitioning into an elevated mixed layer².

The vertical juxtaposition of the overriding elevated mixed layer and the underlying moist boundary layer creates a formidable thermodynamic profile. The base of the elevated mixed layer acts as a strong capping inversion, a phenomenon critical to the severity of the subsequent convection. This lid strongly resists vertical motion, establishing substantial convective inhibition within the profile⁵. While this inhibition prevents immediate thunderstorm development, the boundary layer experiences unimpeded advection of rich moisture from the Mediterranean Sea or the Atlantic Ocean¹⁸. Because the lid suppresses early, disorganized

deep convection, sensible heating at the surface and continuous moisture advection throughout the diurnal cycle lead to soaring values of convective available potential energy⁸. The mid-level dryness of the elevated mixed layer, combined with its exceptionally steep lapse rates, further enhances the downdraft convective available potential energy. This specific parameter is critical, as the entrainment of dry mid-level air into a saturated precipitation core induces rapid evaporative cooling. This cooling accelerates the downdraft, translating into a severe threat for damaging straight-line winds and microbursts once the storms finally initiate⁶.

Triggering Mechanisms and Convective Initiation

Despite the presence of extreme latent instability, deep moist convection cannot materialize in the core of the plume where the convective inhibition remains insurmountable⁸. Convective initiation typically occurs at the western or northern periphery of the elevated mixed layer, where the capping inversion naturally weakens, elevates, or is mechanically breached by localized forcing⁸.

Several dynamic mechanisms facilitate this critical breach. The foremost mechanism is underrunning, a process where the moist boundary layer air horizontally advects past the geographical boundary of the elevated mixed layer. Upon escaping the suppressing influence of the cap, the unstable boundary layer air immediately ascends into the conditionally unstable free troposphere, resulting in explosive, rapid-onset convection⁴.

Additionally, Quasigeostrophic theory and Q-vector diagnostics have been utilized to understand the forcing of prefrontal convergence lines in these environments. Research by Dahl and Fischer indicates that the low-level thermal ridge associated with the base of the elevated mixed layer can force strong horizontal convergence and vertical ascent directly ahead of the primary surface cold front. This robust dynamic lifting physically forces air parcels to their level of free convection, mechanically overcoming the existing convective inhibition². Upper-level synoptic forcing also plays an indispensable role. Positive vorticity advection occurring ahead of the advancing upper-level trough induces broad, synoptic-scale ascent. This ascent cools the mid-levels of the atmosphere, which steepens the lapse rates further and gradually erodes the strength of the capping inversion through adiabatic expansion¹⁰. The combination of a weakening lid and rising surface temperatures ultimately allows parcels to reach the convective temperature, obliterating the remaining inhibition and triggering widespread mesoscale convective systems.

Sub-Classifications and Synoptic Typologies

While the classical Spanish plume model dictates a specific and highly recognizable flow regime, rigorous ongoing research has necessitated the development of sub-classifications to account for variations in storm tracks, spatial extents, and synoptic positioning.

Lewis and Gray established a widely accepted framework categorizing these synoptic environments over the United Kingdom and adjacent continent into three distinct typologies based on mid-level flow and frontal behavior. The first is the Classical Spanish Plume. This configuration is driven by an eastward-propagating baroclinic cyclone evolving according to an idealized life cycle. Conditional instability is released from a warm, moist plume of air that is

capped by warmer, extremely dry air originating from the Spanish plateau¹. Thunderstorms in this classic regime frequently target southeastern England, the English Channel, and western France, exhibiting a relatively narrow corridor of severe risk²³.

The second category is the Modified Spanish Plume. This setup features a slowly moving, mature frontal system associated with a forward-tilting trough or a stagnant cutoff low at the 500 hPa level²². The meridional flow in this regime is significantly more pronounced than in the classical setup. This strong southerly component allows the elevated mixed layer to advect much further north and west before the cold front overtakes the warm sector. Consequently, this modified plume is responsible for severe convective outbreaks extending deep into Ireland, Northern England, and Scotland, broadening the spatial threat considerably¹.

The third category is the European Easterly Plume. This less frequent but highly impactful setup is associated with an omega block centered over Scandinavia. This blocking pattern drives an atypical easterly flow at upper levels that dynamically interacts with the northward-moving plume, resulting in complex shear profiles and atypical storm motions that challenge operational numerical weather prediction models²².

Furthermore, the meteorological parameters defining the Spanish plume are not strictly confined to western Europe. Similar synoptic environments have been heavily documented in the Baltic states and Scandinavia. For instance, in Finland, severe mesoscale convective systems frequently develop when a warm, moist air mass flows into the region from the south or southeast under the influence of an upper-level trough. Punkka and Bister noted that this specific thermodynamic configuration shares distinct similarities with the Spanish plume, relying on differential advection and mid-level steep lapse rates to generate severe summertime convection at unusually high latitudes¹⁸.

Climatological Baselines and Intercontinental Comparisons

To fully appreciate the severity and uniqueness of the Spanish plume, it is highly instructive to compare it against its North American counterpart. Taszarek et al. conducted an exhaustive comparative climatology utilizing ERA5 reanalysis data spanning from 1979 to 2018, demonstrating that while both Europe and the United States experience severe weather driven by elevated mixed layers, their climatological baselines and thermodynamic ceilings differ significantly²⁶.

In the United States, the elevated mixed layer typically originates over the high terrain of the Rocky Mountains, the Desert Southwest, and the Mexican Plateau. As this layer is advected eastward over the Great Plains by the mid-latitude westerlies, it interacts with exceptionally rich, deep moisture surging northward from the Gulf of Mexico. This combination of extreme mid-level lapse rates and virtually limitless low-level moisture yields tail-distribution convective available potential energy values frequently reaching 6,000 to 8,000 Joules per kilogram²⁶.

In contrast, the Spanish plume over Europe relies heavily on localized Mediterranean or Atlantic moisture, which is often shallower and less absolute in its humidity than the Gulf of Mexico fetch. Consequently, European severe environments feature lower maximum convective

available potential energy values, typically peaking between 3,000 and 4,000 Joules per kilogram during the most extreme events²⁶.

However, despite possessing lower absolute potential energy, European severe storms demonstrate a surprisingly high probability of convective initiation given a favorable environment. The lower-level lapse rates over Europe are often higher, driven by complex terrestrial heating over varied land cover, and the capping inversion—while present and effective—is frequently easier to breach than the robust, unyielding lids routinely observed over the US Great Plains²⁶. Furthermore, European environments often feature substantial bulk wind shear, allowing even moderate potential energy environments to routinely support highly organized supercells and long-lived mesoscale convective systems.

Extreme Meteorological Manifestations and Case Studies

The explosive release of thermodynamic energy within a Spanish plume environment produces some of the most violent and economically destructive meteorological events documented in Europe. The juxtaposition of steep mid-level lapse rates, high potential energy, and strong deep-layer shear are prime drivers for a wide spectrum of severe hazards.

Extreme Hail and Flash Flooding

The rapid vertical acceleration present in Spanish plume updrafts, driven by the sudden release of built-up instability, is highly conducive to the formation of large hail. A historical precedent was set during the 1968 dust fall storms, where severe thunderstorms transported massive quantities of mineral dust directly within the plume. This event resulted in historical rainfall and extreme hail, utilizing the suspended dust as highly efficient cloud condensation nuclei⁹. More recently, on 28 June 2012, a severe Spanish plume event triggered a prolific supercell outbreak across the English Midlands. These storms dropped hailstones large enough to cause significant structural damage to vehicles and greenhouses, and the intense lightning and precipitation severely disrupted the 2012 Summer Olympics torch relay as it traversed the region¹. Furthermore, the slow-moving nature of these convective bands, particularly when surface boundaries align parallel to the mid-level flow, leads to severe flash flooding. This mechanism was the primary driver of the devastating Italian floods of November 1994 and remains a recurrent flash flood threat over the Benelux region and the Rhine Valley⁹.

Bow Echoes, Derechos, and Severe Wind

When steep lapse rates overlap with intense, unidirectional vertical wind shear, Spanish plume environments can spawn highly organized quasi-linear convective systems and fast-moving derechos. A catastrophic example occurred on 18 August 2011, striking the Pukkelpop music festival in Belgium. Initial discrete convective cells rapidly merged into a massive bow echo featuring a distinct bookend vortex. This system swept the region with extreme precipitation, large hail, and devastating straight-line downbursts. Peak wind gusts were measured and estimated between 23 and 37 meters per second. The extreme winds caused stage collapses and uprooted massive trees—all falling in a uniform direction indicative of an extreme

microburst rather than a tornadic circulation—resulting in five fatalities and hundreds of injuries²⁰.

Post-event analysis by De Meutter et al. confirmed that the thermodynamic profile was heavily influenced by a Spanish plume setup. The extreme downdraft velocities were mathematically linked to the rapid evaporation of precipitation into the dry mid-level air of the elevated mixed layer. The entrainment of this dry air into the saturated downdraft caused extreme negative buoyancy, propelling the air toward the surface at lethal speeds²⁰. Similarly, the Pentecost weekend storms in June 2014 produced a deadly, progressive derecho that swept through France and western Germany, generating wind gusts up to 144 kilometers per hour and causing widespread infrastructural collapse⁹.

Meteotsunamis and Coastal Seiches

An often-overlooked but highly dangerous hazard associated with Spanish plume systems is the generation of meteotsunamis—atmospherically generated oceanic waves that possess periods and destructive characteristics identical to seismic tsunamis. High-velocity, quasi-linear mesoscale convective systems migrating rapidly across the shallow waters of the English Channel and the North Sea during Spanish plume events can dynamically couple with the ocean surface.

When the propagation speed of the atmospheric pressure jump—typically the leading edge of the convective rear-flank downdraft—matches the shallow-water wave celerity of the ocean, a phenomenon known as Proudman resonance occurs³⁰. Williams et al. conducted an eight-year climatology revealing that wave amplification near predictions of Proudman resonance can occur even if the instantaneous forcing speed deviates from the mean forcing speed by up to 18 percent. This resonance results in a linear superposition and constructive interference of waves, generating sudden, destructive coastal flooding events completely independent of seismic activity. This mechanism has been definitively linked to dangerous seiches and meteotsunamis impacting the coasts of France, the United Kingdom, and the Netherlands during peak summer Spanish plume outbreaks².

Methodological Critiques, Ambiguities, and Future Outlook

A critical and ongoing review of the literature reveals persistent ambiguities in how the term "Spanish plume" is applied, raising vital questions about the fidelity and universality of the conceptual model itself. While the foundational work by Carlson and Ludlam clearly defined the plume strictly as the elevated mixed layer of dry, hot air aloft⁵, subsequent operational literature has frequently conflated this with the low-level, moist marine air mass that undercuts it¹. This semantic confusion routinely obscures the physical reality that the "plume" itself is bone-dry and fundamentally suppresses convection, whereas the underlying maritime air is the true fuel for the storms.

Moreover, the geographical origin of the air mass is heavily debated and increasingly scrutinized via modern computational back-trajectory modeling. Advanced trajectory analyses indicate that the air comprising the elevated mixed layer during a classical "Spanish plume"

event does not always originate over the Iberian Peninsula. Frequently, the air traces directly back to the deep Saharan boundary layer in North Africa⁸. During its rapid poleward advection ahead of a deep trough, the air mass merely passes over Spain without significant thermodynamic interaction with the Meseta Central's terrain. In some documented cases, the air does not cross Spain at all, instead advecting over the Mediterranean directly into France, raising fundamental questions regarding the appropriateness of the nomenclature¹. This discrepancy highlights a critical need for modern meteorological literature to disentangle the strict thermodynamic definition of an elevated mixed layer from regional colloquialisms that may obscure the actual physical drivers of severe weather¹.

Looking toward the future, anthropogenic climate change is projected to alter the climatological baselines governing these events. Increasing continental surface temperatures and alterations in boundary-layer moisture—such as rising mixing ratios across the Mediterranean Sea—suggest that the thermodynamic floor for Spanish plume events is rising⁶. While dynamic forcing variables, such as deep-layer bulk wind shear, may remain static or slightly decrease in a warming climate due to a reduced pole-to-equator temperature gradient, the exponential increase in available latent heat guarantees a more volatile environment. Consequently, when Spanish plume synoptic configurations do materialize in the future, the absolute ceiling for severe hazards—particularly extreme convective precipitation, devastating microbursts, and exceptionally large hail—will be substantially elevated, demanding heightened vigilance and advanced predictive modeling from the European meteorological community³⁴.

Works cited

1. The Spanish Plume Elevated Mixed Layer: A Review of Its Use and Misuse within the Scientific Literature in - AMS Journals, <https://journals.ametsoc.org/view/journals/mwre/153/5/MWR-D-24-0139.1.xml>
2. The Origin of Western European Warm-Season Prefrontal Convergence Lines - AMS Journals, https://journals.ametsoc.org/view/journals/wefo/31/5/waf-d-15-0161_1.pdf
3. The Spanish Plume Elevated Mixed Layer: A Review of Its Use and Misuse within the Scientific Literature - ResearchGate, https://www.researchgate.net/publication/389337980_The_Spanish_Plume_Elevated_Mixed_Layer_Review_of_its_Use_and_Misuse_within_the_Scientific_Literature
4. Properties and Characteristics of Atmospheric Deserts over Europe - EGU sphere, <https://egusphere.copernicus.org/preprints/2025/egusphere-2025-3552/egusphere-2025-3552.pdf>
5. FACTORS AFFECTING THE SEVERITY OF THUNDERSTORMS AND THEIR RELATIONSHIP WITH AGRICULTURE - CABI Digital Library, <https://www.cabidigitallibrary.org/doi/pdf/10.5555/20230125527>
6. Mechanisms of Projected Changes in Thunderstorm Downburst Environments Across the United States - ESS Open Archive, <https://essopenarchive.org/doi/pdf/10.22541/essoar.175214727.71008323/v1?onload=true>
7. Origins of Extreme CAPE Around the World - MIT,

- https://texmex.mit.edu/pub/emanuel/PAPERS/Tuckman_Emanuel_2024.pdf
8. Atmospheric deserts and extreme weather events: co-occurrence with positive temperature anomalies, thunderstorms, and dust - EGU sphere, <https://egusphere.copernicus.org/preprints/2026/egusphere-2026-2350/egusphere-2026-2350.pdf>
 9. Spanish plume - Wikipedia, https://en.wikipedia.org/wiki/Spanish_plume
 10. Spanish Plume Literature Review and Case Studies - University of Manchester - Figshare, https://figshare.manchester.ac.uk/articles/dataset/Spanish_Plume_Literature_Review_and_Case_Studies/28022981
 11. 'Spanish plume' headed to the UK, but not for long - expert comment from Reading meteorologist, <https://archive.reading.ac.uk/news-events/2016/August/pr691613.html>
 12. Impacts of Climate Change On Human Health | PDF - Scribd, <https://www.scribd.com/document/611498332/Impacts-of-Climate-Change-on-Human-Health>
 13. Thesis Doctor of Philosophy Stephen David Burt - CentAUR, https://centaur.reading.ac.uk/104769/2/29804706_Burt_Thesis_Redacted.pdf
 14. F. H. Ludlam's research works | Imperial Valley College and other places - ResearchGate, <https://www.researchgate.net/scientific-contributions/F-H-Ludlam-2051891561>
 15. Analyzing 23 years of warm-season derechos in France: a climatology and investigation of synoptic and environmental changes - WCD, <https://wcd.copernicus.org/preprints/wcd-2023-8/wcd-2023-8-manuscript-version2.pdf>
 16. The development of the Spanish plume. Solid lines indicate surface... - ResearchGate, https://www.researchgate.net/figure/The-development-of-the-Spanish-plume-Solid-lines-indicate-surface-flow-and-dashed-lines_fig3_389337980
 17. European supercell thunderstorms—A prevalent... : Science Advances - Ovid, <https://www.ovid.com/journals/sciadv/fulltext/10.1126/sciadv.adx0513~european-supercell-thunderstormsa-prevalent-current-threat>
 18. Occurrence of Summertime Convective Precipitation and Mesoscale Convective Systems in Finland during 2000–01 - American Meteorological Society, <https://journals.ametsoc.org/view/journals/mwre/133/2/mwr-2854.1.pdf>
 19. Occurrence of Summertime Convective Precipitation and Mesoscale Convective Systems in Finland during 2000–01 in - AMS Journals, <https://journals.ametsoc.org/view/journals/mwre/133/2/mwr-2854.1.xml>
 20. Predicting Small-Scale, Short-Lived Downbursts: Case Study with the NWP Limited-Area ALARO Model for the Pukkelpop Thunderstorm - AMS Journals, <https://journals.ametsoc.org/view/journals/mwre/143/3/mwr-d-14-00290.1.pdf>
 21. The Origin of Western European Warm-Season Prefrontal Convergence Lines in - AMS Journals, https://journals.ametsoc.org/view/journals/wefo/31/5/waf-d-15-0161_1.xml
 22. Categorisation of synoptic environments associated with Mesoscale Convective

- Systems over the UK | Request PDF - ResearchGate,
https://www.researchgate.net/publication/46283835_Categorisation_of_synoptic_environments_associated_with_Mesoscale_Convective_Systems_over_the_UK
23. A sub classification of the classic spanish plume? - Snow, Storms & Severe Weather - Netweather Community,
<https://community.netweather.tv/topic/100104-a-sub-classification-of-the-classic-spanish-plume/>
 24. The synoptic setting of thunderstorms in western Europe -
webpace.science.uu.nl,
<https://www.staff.science.uu.nl/~delde102/Syn.Set.Thunderstorms.pdf>
 25. CS9 Severe Thunderstorm Case Study - 26 June 2020,
https://www.metpod.co.uk/met_case_studies/severe-thunderstorm/
 26. Severe Convective Storms across Europe and the United States. Part II: ERA5 Environments Associated with Lightning, Large Hail, - AMS Journals,
<https://journals.ametsoc.org/view/journals/clim/33/23/jcliD200346.pdf>
 27. Severe Convective Storms across Europe and the United States. Part II: ERA5 Environments Associated with Lightning, Large Hail, Severe Wind, and Tornadoes in - AMS Journals,
<https://journals.ametsoc.org/view/journals/clim/33/23/jcliD200346.xml>
 28. Severe Convective Storms across Europe and the United States. Part II: ERA5 Environments Associated with Lightning, Large Hail, Severe Wind, and Tornadoes - ResearchGate,
https://www.researchgate.net/publication/344371238_Severe_Convective_Storms_across_Europe_and_the_United_States_Part_II ERA5 Environments Associated with Lightning Large Hail Severe Wind and Tornadoes
 29. Monthly Weather Review - AMS Journals - American Meteorological Society,
<https://journals.ametsoc.org/view/journals/mwre/153/5/mwr1535cover.pdf>
 30. An 8-yr Meteotsunami Climatology across Northwest Europe: 2010–17 in - AMS Journals,
<https://journals.ametsoc.org/view/journals/phoc/51/4/JPO-D-20-0175.1.xml>
 31. Predicting Small-Scale, Short-Lived Downbursts: Case Study with the NWP Limited-Area ALARO Model for the Pukkelpop Thunderstorm in - AMS Journals,
<https://journals.ametsoc.org/view/journals/mwre/143/3/mwr-d-14-00290.1.xml>
 32. Meteotsunami generation, amplification and occurrence in north-west Europe - University of Liverpool Repository,
https://livrepository.liverpool.ac.uk/3073478/1/201126790_Nov2019.pdf
 33. Differing Trends in United States and European Severe Thunderstorm Environments in a Warming Climate in - AMS Journals,
<https://journals.ametsoc.org/view/journals/bams/102/2/BAMS-D-20-0004.1.xml>
 34. Analysing 23 years of warm-season derechos in France: a climatology and investigation of synoptic and environmental changes - WCD,
<https://wcd.copernicus.org/articles/5/439/2024/wcd-5-439-2024.pdf>
 35. A pan-European analysis of large-scale drivers of severe convective outbreaks - EGU sphere,
<https://egusphere.copernicus.org/preprints/2025/egusphere-2025-2296/egusphere>

[re-2025-2296.pdf](#)