

MATH1061/1021 Cheatsheet (Calculus)

Semester 1, 2024

Abyan Majid

 [Source](#)

1	Sets, numbers, functions	2	3.10 Finding max & min on a closed interval . . .	7	
1.1	Union, intersection, difference, comple- ment, subset	2	3.11 Concavity	7	
1.2	Common number sets	2	3.12 Second-derivative test	7	
1.3	Interval notation	2	3.13 Sensible order to curve-sketching	8	
1.4	Modulus	2	3.14 Tangent line (local linear approximation)	8	
1.5	Injective, surjective, bijective functions . .	2	3.15 Taylor polynomials	8	
1.6	Composite, inverse, hyperbolic functions .	3	3.16 Lagrange’s form of remainder	8	
			3.17 Partial sum, infinite series, geometric series	9	
			3.18 Taylor series	9	
2	Limits	3	4	Integral calculus	9
2.1	Definition of a limit	3	4.1	Riemann sums	9
2.2	Limit laws	3	4.2	Riemann integral	9
2.3	One-sided limits, existence of limit	3	4.3	Properties of Riemann integral	10
2.4	Infinite limits	4	4.4	Fundamental Theorem of Calculus I	10
2.5	Squeeze law	4	4.5	Fundamental Theorem of Calculus II	10
2.6	Continuous functions	4	4.6	Leibinz integral rule	10
			4.7	Logarithms	10
			4.8	Properties of logarithms	10
			4.9	Properties of exponentials	11
			4.10	Integration	11
			4.11	Linearity property of integration	11
			4.12	Integration by substitution	11
			4.13	Integration by parts	12
			4.14	Integration by partial fractions	12
			4.15	Length of a curve	12
			4.16	Solid of revolution	12
			4.17	Methods for finding solid of revolution . .	13

1 Sets, numbers, functions


1.1 Union, intersection, difference, complement, subset

Operation	Notation	Description
Union	$A \cup B$	All elements that are in A or B or both
Intersection	$A \cap B$	All elements that are both in A and B
Difference	$A \setminus B$	All elements that are in A but not in B
Complement	\overline{A}	All elements in the universal set that are not in A
Subset	$A \subseteq B$	A is a subset of B ; every element in A is also in B
Proper subset	$A \subset B$	A is a subset of B and $A \neq B$

1.2 Common number sets

Name (Symbol)	Set
Natural numbers (\mathbb{N})	$\{0, 1, 2, 3, 4, \dots\}$
Integers (\mathbb{Z})	$\{\dots - 2, -1, 0, 1, 2, \dots\}$
Rational numbers (\mathbb{Q})	$\left\{\frac{1}{2}, -\frac{4}{3}, \frac{17}{12}, \dots\right\}$
Irrational numbers ($\mathbb{R} \setminus \mathbb{Q}$)	$\{\sqrt{2}, \pi, e, \sqrt{7}, \dots\}$
Real numbers (\mathbb{R})	All numbers

1.3 Interval notation

Interval notation
$D = [-1, 1)$  <p>Note: Infinity ∞ is always open-bracketed, i.e. (∞, ∞)</p>

1.4 Modulus

Modulus
Distance on the number line
$ x - y $

1.5 Injective, surjective, bijective functions

Function type	Definition
Injective	A function $f : X \rightarrow Y$ where for all $x \in X$, x maps to a different $y \in Y$, and $ X \leq Y $.
Surjective	A function $f : X \rightarrow Y$ where for all $y \in Y$, y is an image of some $x \in X$, and $ X \geq Y $.
Bijective	A function that is both injective and surjective where $ X = Y $.

1.6 Composite, inverse, hyperbolic functions

Composite functions
A function in the form $f \circ g(x) = f(g(x))$ where some function f takes another function g as input.

Inverse functions	
Definition	How to invert a function
There exists an inverse function f^{-1} if and only if f is injective. If $f(x) = y$, then $f^{-1}(y) = x$, which implies that if $f : A \rightarrow B$, then $f^{-1} : B \rightarrow A$.	1) Rewrite $f(x)$ as y 2) Swap x and y 3) Make y the subject 4) Rewrite y as $f^{-1}(x)$

Hyperbolic functions		
Function	Derivative	Inverse
$\cosh(x) = \frac{e^x + e^{-x}}{2}$	$\frac{d}{dx} \cosh(x) = \sinh(x)$	$\cosh^{-1}(x) = \ln(x + \sqrt{x^2 - 1})$
$\sinh(x) = \frac{e^x - e^{-x}}{2}$		$\sinh^{-1}(x) = \ln(x + \sqrt{x^2 + 1})$

2 Limits

2.1 Definition of a limit

Limit
$\lim_{x \rightarrow a} f(x) = L$ as x approaches a , but not reaching a , the output gets closer and closer to L .

2.2 Limit laws

if we have $\lim_{x \rightarrow c} f(x) = L, \lim_{x \rightarrow c} g(x) = M$	
No	Law
1	$\lim_{x \rightarrow c} [f(x) + g(x)] = L + M$
2	$\lim_{x \rightarrow c} [f(x) - g(x)] = L - M$
3	$\lim_{x \rightarrow c} [f(x)g(x)] = LM$
4	$\lim_{x \rightarrow c} \left[\frac{f(x)}{g(x)} \right] = \frac{L}{M}$
5	$\lim_{x \rightarrow c} [f(x)]^n = L^n$
6	$\lim_{x \rightarrow c} \sqrt[n]{f(x)} = \sqrt[n]{L}$

2.3 One-sided limits, existence of limit

One-sided limits
Limit from below (left) is denoted as $\lim_{x \rightarrow c^-} f(x)$ Limit from above (right) is denoted as $\lim_{x \rightarrow c^+} f(x)$

Existence of a limit
<p>A limit exists only if</p> $\lim_{x \rightarrow c^-} f(x) = \lim_{x \rightarrow c^+} f(x) = L$ <p>in which case</p> $\lim_{x \rightarrow c} f(x) = L$

2.4 Infinite limits

Infinite limits
<p>If we take x larger and larger, and $f(x)$ gets closer to L, we say</p> $\lim_{x \rightarrow \infty} f(x) = L$ <p>You can algebraically solve infinite limits by dividing by the highest power of x in the denominator or to by using conjugates.</p>

2.5 Squeeze law

Squeeze law
<p>Suppose $g(x) \leq f(x) \leq h(x)$ for x near a</p> <p>if $\lim_{x \rightarrow a} h(x) = L$ and $\lim_{x \rightarrow a} g(x) = L$, then $\lim_{x \rightarrow a} f(x) = L$</p> <p>we say "$f(x)$ is squeezed between $g(x)$ and $h(x)$"</p>

2.6 Continuous functions

Continuous functions
<p>A function is continuous at a point if the limit exists at the point, and is equal to the value at that point.</p> <p>ie. $f(x)$ is continuous at $x = c$ if and only if:</p> <ol style="list-style-type: none"> 1) $f(c)$ is defined 2) $\lim_{x \rightarrow c} f(x)$ exists and is finite 3) $\lim_{x \rightarrow c} f(x) = f(c)$

3 Differential Calculus

3.1 Limit definition of derivatives

Limit definition of derivatives
<p>The derivative of a function f at a point a is given by</p> $f'(a) = \lim_{h \rightarrow 0} \frac{f(a+h) - f(a)}{h}$

3.2 Common derivatives

Common derivatives	
$f(x)$	$f'(x)$
x^n where $n \neq 0$	nx^{n-1}
$\sin(x)$	$\cos(x)$
$\cos(x)$	$-\sin(x)$
e^x	e^x
$\ln(x)$	$\frac{1}{x}$
$\tan(x)$	$\sec^2(x)$
c^x	$c^x \ln(c)$
$\ln y$	$\frac{1}{y} \frac{dy}{dx}$

3.3 Differential rules

Differential rules		
No	Name	What to do
1	Power rule	$\frac{d}{dx} x^n = nx^{n-1}$
2	Constant rule	$\frac{d}{dx} c = 0$, where $c \in \mathbb{R}$
3	Sum rule	$\frac{d}{dx} [f(x) + g(x)] = f'(x) + g'(x)$
4	Difference rule	$\frac{d}{dx} [f(x) - g(x)] = f'(x) - g'(x)$
5	Product rule	$\frac{d}{dx} [f(x)g(x)] = f(x)g'(x) + g(x)f'(x)$
6	Quotient rule	$\frac{d}{dx} \left[\frac{f(x)}{g(x)} \right] = \frac{f'(x)g(x) - f(x)g'(x)}{[g(x)]^2}$
7	Chain rule	<p>We are to find $\frac{d}{dx} [f \circ g(x)]$</p> <p>Method 1: Let $u = g(x)$, $y = f(u)$, and so $y'(x)$ is given by $\frac{dy}{dx} = \frac{df}{du} \cdot \frac{du}{dx}$</p> <p>Method 2: If you can see f and g clearly, then $\frac{d}{dx} [f \circ g(x)] = f'(g(x))g'(x)$</p>

3.4 Implicit differentiation

Implicit differentiation
<p>Functions that can't (or aren't easy) to be written as $y = f(x)$ can instead be written implicitly in the form $F(x, y) = 0$.</p> <p>In which case, to get the derivative, find $\frac{d}{dx} [F(x, y)]$ applying chain rule for terms involving y, where $\frac{d}{dx} y \Rightarrow \frac{dy}{dx}$ and finally, make $\frac{dy}{dx}$ the subject.</p>

3.5 Useful algebra for logarithmic differentiation

No	Useful algebra
1	$a^x \Leftrightarrow e^{x \ln a}$
2	$\ln(e^{x \ln(a)}) = x \ln a$
3	$y \ln(a) = \ln(x) \Leftrightarrow y = \frac{\ln(x)}{\ln(a)}$
4	$\ln(\sqrt{x}) \Leftrightarrow \frac{1}{2} \ln(x)$
5	$\log_b(a) \Leftrightarrow \frac{\ln(a)}{\ln(b)}$

3.6 L'Hopital's Rule

L'Hopital's Rule
<p>If $\lim_{x \rightarrow c} \frac{f(x)}{g(x)}$ is an indeterminate form, i.e. can be written as $\frac{0}{0}$ or $\pm \frac{\infty}{\infty}$, then $\lim_{x \rightarrow c} \frac{f(x)}{g(x)} = \lim_{x \rightarrow c} \frac{f'(x)}{g'(x)}$ where c can be infinity.</p> <p>NOTE: You can keep applying L'Hopital's Rule until you get a definite form.</p>

3.7 Extrema

Extrema	
An "extremum" is a maximum or a minimum.	
Extremum	Definition
Local max	A local max at $x = c$ means $f(c) \geq f(x)$ for any x near c
Local min	A local min at $x = c$ means $f(c) \leq f(x)$ for any x near c
Global max	A global max on some domain A where $f : A \rightarrow \mathbb{R}$ means $f(c) \geq f(x)$ for all $x \in A$
Global min	A global min on some domain A where $f : A \rightarrow \mathbb{R}$ means $f(c) \leq f(x)$ for all $x \in A$

3.8 First-derivative test

First-derivative test		
If f is continuous and $f'(c) = 0$		
Case	Observation	Conclusion
1	<ul style="list-style-type: none"> $f'(x)$ is positive for $x < c$ $f'(x)$ is negative for $x > c$ 	Local max
2	<ul style="list-style-type: none"> $f'(x)$ is negative for $x < c$ $f'(x)$ is positive for $x > c$ 	Local min
3	$f'(x)$ doesn't change sign	Not a local extremum (neither max or min)

3.9 Critical points

No	Critical point type
1	$f'(c)$ does NOT exist
2	$f'(c) = 0$
NOTES: • If $f(x)$ is differentiable, then any local extrema must occur at points $x = c$ where $f'(c) = 0$ • $f'(c) = 0$ does NOT guarantee a local extremum! You must apply first-derivative test!	

3.10 Finding max & min on a closed interval

Extreme value theorem
If f is a continuous function from a closed interval A to \mathbb{R} , then f attains a global max & global min value in A .

Finding max & min on a closed interval
To find max & min values on a closed interval, simply check: 1) Critical points 2) End points of the interval

3.11 Concavity

Concavity
• Concave up: $f''(x) > 0$ • Concave down: $f''(x) < 0$ • Point of inflection: if $f''(x)$ changes sign at c then $x = c$ is a point of inflection.

3.12 Second-derivative test

Second-derivative test		
If f, f' are differentiable and $f'(c) = 0$		
Case	Observation	Conclusion
1	$f''(c) > 0$	Local minimum
2	$f''(c) < 0$	Local maximum
3	$f''(c) = 0$	No conclusion (might be local min, max, or neither.) What to do from here: Do first-derivative test

3.13 Sensible order to curve-sketching

Sensible order to curve-sketching	
Step	To work out
1	What is the domain?
2	What is the y -intercept?
3	What happens as $x \rightarrow \pm\infty$ (Limits)
4	Is there anywhere that $f(x) \rightarrow \pm\infty$ (is denominator zero?)
5	Where are the critical points? <ul style="list-style-type: none"> • $f'(x) = 0$ • $f'(x)$ does NOT exist
6	Where is $f(x)$ increasing/decreasing? <ul style="list-style-type: none"> • $f'(x) > 0$: increasing • $f'(x) < 0$: decreasing
7	Where is $f(x)$ concave up or down? <ul style="list-style-type: none"> • $f''(x) > 0$: concave up (slope decreasing) • $f''(x) < 0$: concave down (slope increasing)

3.14 Tangent line (local linear approximation)

Tangent line
The tangent line of $f(x)$ at point c is given by $y = f(c) + f'(c)(x - c)$ This is also called the local linear approximation.

3.15 Taylor polynomials

Taylor polynomials
The Taylor polynomial for $f(x)$ around $x = a$ is $p_n(x) = f(a) + f'(a)(x - a) + \frac{f''(a)}{2!}(x - a)^2 + \cdots + \frac{f^{(n)}(a)}{n!}(x - a)^n$ or, using sigma notation, $p_n(x) = \sum_{k=0}^n \frac{f^{(k)}(a)}{k!}(x - a)^k$

3.16 Lagrange's form of remainder

Lagrange's form of remainder
<p>If $p_n(x)$ is the Taylor polynomial of $f(x)$ around $x = a$, then the n-th order remainder is given by</p> $R_n(x) = f(x) - p_n(x),$ <p>which satisfies the limit condition</p> $\lim_{x \rightarrow a} \frac{R_n(x)}{(x - a)^n} = 0.$ <p>If $f(x)$ has continuous derivatives up to order $n + 1$, then $R_n(x)$ can be expressed in</p> <p>Lagrange's Form: There exists a number c, strictly between a and x, for each x, such that</p> $R_n(x) = \frac{f^{(n+1)}(c)}{(n + 1)!}(x - a)^{n+1}.$

3.17 Partial sum, infinite series, geometric series

Term	Definition
Partial sum	$S_n = a_0 + a_1 + a_2 + \dots + a_n = \sum_{k=0}^n a_k$
Infinite series	$S = \sum_{k=0}^{\infty} a_k = \lim_{n \rightarrow \infty} \sum_{k=0}^n a_k$ If this limit exists and is finite, the series converges; otherwise, it diverges.
Geometric series	$\sum_{k=0}^{\infty} r^k = \frac{1}{1-r}$ for $ r < 1$

3.18 Taylor series

Taylor Series
<p>The Taylor series of $f(x)$ near $x = a$ is given by</p> $p_n(x) = f(a) + f'(a)(x-a) + \frac{f''(a)}{2!}(x-a)^2 + \dots + \frac{f^{(n)}(a)}{n!}(x-a)^n$ <p>or, in short,</p> $\sum_{k=0}^n \frac{f^{(k)}(a)}{k!}(x-a)^k$ <p>If the limit as $n \rightarrow \infty$ converges for x, then</p> $f(x) = \lim_{n \rightarrow \infty} p_n(x) = \sum_{k=0}^{\infty} \frac{f^{(k)}(a)}{k!}(x-a)^k$

4 Integral calculus

4.1 Riemann sums

Riemann sums
<p>Given an interval split into n subdivisions from a to b, each of width $\Delta x = \frac{b-a}{n}$, the lower and upper Riemann sums L_n and U_n are given by</p> $L_n = \sum_{i=1}^n m_i \Delta x \text{ and } U_n = \sum_{i=1}^n M_i \Delta x$ <p>where m_i is the minimum value of $f(x)$ on $[x_{i-1}, x_i]$ and M_i is the maximum value of $f(x)$ on $[x_{i-1}, x_i]$.</p>

4.2 Riemann integral

Riemann integral
<p>If we take Δx smaller and smaller, then m_i and M_i will get closer and closer. Hence, if U_n and L_n converge to the same limit, then</p> $\text{Area} = \lim_{n \rightarrow \infty} L_n = \lim_{n \rightarrow \infty} U_n = \int_a^b f(x) dx$

4.3 Properties of Riemann integral

Properties of Riemann integral	
No	Property
1	$\int_a^b c f(x) dx = c \int_a^b f(x) dx$
2	$\int_a^b [f(x) + g(x)] dx = \int_a^b f(x) dx + \int_a^b g(x) dx$
3	$\int_a^b f(x) dx = - \int_b^a f(x) dx$

4.4 Fundamental Theorem of Calculus I

Fundamental Theorem of Calculus I
If $F(x) = \int_a^x f(t) dt$, then $F'(x) = f(x)$ i.e. with area, we can find antiderivative.

4.5 Fundamental Theorem of Calculus II

Fundamental Theorem of Calculus II
$\int_a^b f(x) dx = \int_a^b F'(x) dx = F(b) - F(a)$ i.e. with antiderivative, we can find area.

4.6 Leibinz integral rule

Leibinz integral rule
$\frac{d}{dx} \left(\int_{a(x)}^{b(x)} f(t) dt \right) = f(b(x)) \cdot b'(x) - f(a(x)) \cdot a'(x)$

4.7 Logarithms

Logarithms
If $y = a^x$ then $x = \log_a(y)$ which results in two main rules: $\log_a(x) = \frac{\ln(x)}{\ln(a)}$ and $\frac{d}{dx} [\log_a(x)] = \frac{1}{x \ln(a)}$

4.8 Properties of logarithms

Properties of logarithms	
No	Property
1	$\ln(ax) = \ln(a) + \ln(x)$
2	$\ln\left(\frac{x}{a}\right) = \ln(x) - \ln(a)$
3	$\ln(x^n) = \ln(x \cdot x \cdot \dots \cdot x) = n \ln(x)$

4.9 Properties of exponentials

Properties of exponentials	
No	Property
1	$\ln(e^x) = x$
2	$e^{a+x} = e^a e^x$
3	$(e^x)^n e^{nx}$
4	$\frac{d}{dx}[e^x] = e^x$

4.10 Integration

Integration		
No	$f(x)$	$\int f(x)dx$
1	$x^n \ (n \neq -1)$	$\frac{x^{n+1}}{n+1} + C$
2	e^x	$e^x + C$
3	$\frac{1}{x}$	$\ln(x) + C$
4	$\sin(x)$	$-\cos(x) + C$
5	$\cos(x)$	$\sin(x) + C$
6	$\sinh(x)$	$\cosh(x) + C$
7	$\cosh(x)$	$\sinh(x) + C$
8	$\tan(x)$	$\ln \sec(x) + C$

4.11 Linearity property of integration

Linearity property of integration
$\int f(x) + g(x)dx = \int f(x)dx + \int g(x)dx$ $\int k f(x)dx = k \int f(x)dx$

4.12 Integration by substitution

Integration by substitution
$\int h'(g(x))g'(x)dx = h(g(x)) + C$ <p>which can be made more neat up by letting $u = g(x)$</p> $\int h'(u)\frac{du}{dx}dx = h(u) + C$ <p>and even neater by letting $f(u) = h'(u)$</p> $\int f(u)\frac{du}{dx}dx = \int f(u)du$ <p>(this is usually easier to integrate)</p> <p>Note: For definite integrals, remember to change the bounds</p> $\int_{x=a}^{x=b} f(u)\frac{du}{dx}dx = \int_{u(a)}^{u(b)} f(u)du$

4.13 Integration by parts

Integration by parts
$\int u'v dx = uv - \int uv' dx$ <p>or, for definite integrals:</p> $\int_a^b u'v dx = [uv]_a^b - \int_a^b uv' dx$


4.14 Integration by partial fractions

Integration by partial fractions
<p>For a function f that can be written in the form</p> $f(x) = \frac{a + bx}{(x - \lambda)(x - \mu)}$ <p>for distinct constants λ and μ, we can express $f(x)$ as a sum of simpler fractions</p> $f(x) = \frac{A}{x - \lambda} + \frac{B}{x - \mu}$ <p>where A, B can be found by equating $A(x - \mu) + B(x - \lambda) = ax + b$ and choosing $x = \lambda$ and $x = \mu$ respectively</p> $A(\lambda - \mu) = a\lambda + b \Rightarrow A = \frac{a\lambda + b}{\lambda - \mu}$ $B(\mu - \lambda) = a\mu + b \Rightarrow B = \frac{a\mu + b}{\mu - \lambda}$ <p>The decomposed fractions should be easier to integrate</p> $\int f(x) dx = \int \left(\frac{A}{x - \lambda} + \frac{B}{x - \mu} \right) dx = \int \frac{A}{x - \lambda} dx + \int \frac{B}{x - \mu} dx$ <p>which, using natural logarithms, results to</p> $A \ln x - \lambda + B \ln x - \mu + C$

4.15 Length of a curve

Length of a curve
<p>If $f : [a, b] \rightarrow \mathbb{R}$ is differentiable, then the length of its curve is given by</p> $\int_a^b \sqrt{1 + [f'(x)]^2} dx$

4.16 Solid of revolution



4.17 Methods for finding solid of revolution

Disk method
Let R be the region bounded by $y = f(x) \geq 0, y = 0, x_1 = a, x_2 = b$. The volume V of the solid S formed by revolving R around The x -axis is given by $V = \pi \int_a^b (f(x))^2 dx$

Washer method (generalized disk method)
Let region $R = \{(x, y) \in \mathbb{R}^2 0 \leq f(x) \leq y \leq g(x), a \leq x \leq b\}$. The volume V of the solid S formed by revolving R around the x -axis is given by $V = \pi \int_a^b [g(x)]^2 - [f(x)]^2 dx$

Shell method
Let R be the region with $0 \leq a \leq x \leq b$ and $0 \leq y \leq f(x)$ The volume V of the solid S formed by revolving R around the y -axis is given by $V = 2\pi \int_a^b xf(x)dx$