

Regression

Data Science Dojo

Agenda

- Introduction
- Cost Function & Gradient Descent
 - Minimization
 - Implementation
- Hands-on Example
- Evaluating Regression Models
- Regularization

INTRODUCTION


Regression



Sales Forecasts



Housing Price
Predictions



Daily Temperature
Highs & Lows

Regression vs Classification

- Classification

- Target is discrete with finite value set
- **Examples:** survived/dead, face/non-face, fraud/non-fraud

- Regression

- Target is continuous
- **Examples:** price, weight, height, temperature,

Notation

Passenger Id	Survived	Pclass	Name	Sex	Age	SibSp	Parch	Ticket	Fare	Cabin	Embarked
1	0	3	Braund, Mr. Owen Harris	male	22	1	0	A/5 21171	7.25		S
2	1	1	Cumings, Mrs. John Bradley (Florence Briggs Thayer)	female	38	1	0	PC 17599	71.2833	C85	C
3	1	3	Heikkinen, Miss. Laina	female	26	0	0	STON/O2. 3101282	7.925		S
4	1	1	Futrelle, Mrs. Jacques Heath (Lily May Peel)	female	35	1	0	113803	53.1	C123	S
5	0	3	Allen, Mr. William Henry	male	35	0	0	373450	8.05		S

x_4^5

5: The passenger is in the 5th row

4: The passenger's name is the 4th column

Notation: Ozone Dataset

So how do we describe all the rows?

	ozone	radiation	temperature	wind
Row 1	41	190	67	7.4
Row 2	36	118	72	8.0
Row 3	12	149	74	12.6
	18	313	62	11.5
	23	299	65	8.6
	19	99	59	13.8

$$x^1 = [190, 67, 7.4]$$

$$x^2 = [118, 72, 8.0]$$

$$x^3 = [149, 74, 12.6]$$

Notation: Ozone Dataset

The ozone dataset uses radiation, temperature and wind to predict ozone levels.

		x_1	x_2	x_3	
	ozone	radiation	temperature	wind	
Y	41	190	67	7.4	X
	36	118	72	8.0	
	12	149	74	12.6	
	18	313	62	11.5	
	23	299	65	8.6	
	19	99	59	13.8	

Using this notation, we can describe all the columns of the dataset.

Notation Summary

x^i	– Each row of features	}	Features
x_i	– Each column of features		
X	– Set of all the feature columns		
y^i	– Each row of the target(s)	}	Target
Y	– Set of all the target columns		
n	– Number of rows in the dataset		
m	– Number of columns in the dataset		

COST FUNCTION AND GRADIENT DESCENT


Defining a line

How do we define a line in slope-intercept notation?

- $y = mx + b$


In θ notation

- $h_{\theta}(x) = \theta_0 + \theta_1 x$



What is a good regression line?


- Wind Speed=15 mph
- Ozone = ?
- Use the line that is **somewhere in the middle**
- How do we define "middle"?



$$h_{\theta}(x) = \theta_0 + \theta_1 x$$

Residuals

Difference between hypothesis $h_{\theta}(x)$ (predicted value) and true value (known target)




Cost Function

Minimize the 'cost' or 'loss' function – $J(\theta)$


- Smaller for lower error
- Larger for higher error

$$J(\theta) = \frac{1}{2n} \sum_{i=1}^n (h_{\theta}(x^i) - y^i)^2$$




Cost Function

$$h_{\theta}(x) = \theta_0 + \theta_1 x$$




$$J(\theta) = \frac{1}{2n} \sum_{i=1}^n (h_{\theta}(x^i) - y^i)^2$$



Cost function in two dimensions

$$h_{\theta}(x) = \theta_0 + \theta_1 x$$

$$J(\theta) = \frac{1}{2n} \sum_{i=1}^n (h_{\theta}(x^i) - y^i)^2$$



HOW DO WE FIND OUT THE MINIMUM OF THE COST FUNCTION

Maximum/Minimum Problem

Find two nonnegative numbers whose sum is 9 and so that the product of one number and the square of the other number is a maximum.

Solution

Sum of number is 9

$$9 = x + y$$

Product of two numbers is

$$\begin{aligned} P &= x y^2 \\ &= x (9-x)^2 \end{aligned}$$

Solution

$$\begin{aligned} P' &= x(2)(9-x)(-1) + (1)(9-x)^2 \\ &= (9-x)[-2x + (9-x)] \\ &= (9-x)[9-3x] \\ &= (9-x)(3)[3-x] \\ &= 0 \end{aligned}$$

$$x=9 \text{ or } x=3$$

Maximum Problem

There are 50 apple trees in an orchard. each tree produces 800 apples. For each additional tree planted in the orchard, the apple output per tree drops by 10 apples. how many additional trees should be planted in the existing orchard in order to maximize the apple output of the orchard?

Solution

$$A = (50+t)(800-10t) = 40,000 + 300t - 10t^2$$

Solve for A' and set to 0 to find maximum.

$$A' = -20t + 300 = 0$$

$$t = 15$$

Adding 15 trees will maximize apple production

Gradients

- **Derivative:** slope in one direction
- What about more features?
- **Gradient:** a multi-dimensional derivative

Gradient Descent

- Goal : minimize $J(\theta)$
- Start with some initial θ and then perform an update on each θ_j in turn:

$$\theta_j^{k+1} := \theta_j^k - \alpha \frac{\partial}{\partial \theta_j} J(\theta^k)$$


- Repeat until θ converges

Gradient Descent

$$\theta_j^{k+1} := \theta_j^k - \alpha \frac{\partial}{\partial \theta_j} J(\theta^k)$$

- α is known as the learning rate; set by user
- Each time the algorithm takes a step in the direction of the steepest descent and $J(\theta)$ decreases.
- α determines how quickly or slowly the algorithm will converge to a solution


Intuition



$$\theta_j^{k+1} := \theta_j^k - \alpha \frac{\partial}{\partial \theta_j} J(\theta^k)$$


Learning Rate Effects Large α

$$\theta_j^{k+1} := \theta_j^k - \alpha \frac{\partial}{\partial \theta_j} J(\theta^k)$$




Learning Rate Effects Small α

$$\theta_j^{k+1} := \theta_j^k - \alpha \frac{\partial}{\partial \theta_j} J(\theta^k)$$



Gradient Descent Implementation

When do we stop updating?



- When θ_j^{k+1} is close to θ_j^k
- When $J(\theta^{k+1})$ is close to $J(\theta^k)$ [Error does not change]

Batch Gradient Descent

- How do we incorporate all our data?
- Loop!

For j from 0 to m :

$$\theta_j^{k+1} := \theta_j^k + \alpha \sum_{i=1}^n (y^i - h_{\theta}(x^i)) x_j^i$$

- h_{θ} is updated only once the loop has completed
- Weaknesses?

Stochastic Gradient Descent

- Consider an alternative approach:

for i from 1 to n:

for j from 0 to m:

$$\theta_j^{k+1} := \theta_j^k + \alpha (y^i - h_\theta(x^i)) x_j^i$$

- h_θ is updated when inner loop is complete
- If the training set is big, converges quicker than batch
- May oscillate around a minimum of $J(\theta)$ and never converge

Batch vs. Stochastic

Which is the best to use? It depends.

	Batch Gradient Descent	Stochastic Gradient Descent
Function	Updates hypothesis by scanning whole dataset	Updates hypothesis by scanning one training sample at a time
Rate of convergence	Slowly	Quickly (but may oscillate at minimum)
Appropriate Dataset Size	Small	Large

EVALUATING REGRESSION MODELS

Evaluation metrics for regression

- Mean Absolute Error (MAE)
- Root-Mean-Square Error (RMSE)
 - Root-Mean-Square Deviation
- Coefficient of Determination (R^2)

Mean Absolute Error

$$MAE(\theta) = \frac{\sum_{i=1}^n |h_{\theta}(x^i) - y^i|}{n}$$

- Mean of residual values
- "Pure" measure of error

Root-Mean-Square Error

$$RMSE(\theta) = \sqrt{\frac{\sum_{i=1}^n (h_{\theta}(x^i) - y^i)^2}{n}}$$

- Square root of mean of squared residuals
- Penalizes large errors more than small
- Good when large errors particularly bad

Coefficient of Determination (R^2)

$$R^2 = 1 - \frac{SS_{res}}{SS_{tot}}$$

where

$$SS_{res} = \sum_{i=1}^n (h_{\theta}(x^i) - y^i)^2 \quad SS_{tot} = \sum_{i=1}^n (y^i - \bar{y})^2$$



SS_{res} – Sum of squared residuals (i.e. total squared error)

SS_{tot} – Sum of squared differences from mean (i.e. total variation in dataset)

Result: Measure of how well the model explains the data


- "Fraction of variation in data explained by model"

Coefficient of Determination



R² Example

- $R^2 = 0.277$
- Want a much better model for real application
- $R^2 = 0.6$ can be a good model




REGULARIZATION


Overfitting

- Want to extract general trends
- Danger: "memorizing" the training set
- A model is **overfit** when model performance on test set is much worse than on training set.


Overfitting



$$\theta_0 + \theta_1 x$$



$$\theta_0 + \theta_1 x + \theta_2 x^2$$




$$\theta_0 + \theta_1 x + \theta_2 x^2 + \theta_3 x^3 + \theta_4 x^4$$


Complexity

- What makes a good model overfit?
 - Nature of training data
 - **Complexity of model**
- How do we handle these?
 - Cross validation
 - Manual model constraint
 - Regularization

Intuition



$$\theta_0 + \theta_1 x + \theta_2 x^2 + \theta_3 x^3 + \theta_4 x^4$$



$$\theta_0 + \theta_1 x + \theta_2 x^2 + \theta_3 x^3 + \theta_4 x^4$$

- Want to discourage complex models automatically – How?
- Adjust the cost function!
 - Penalize models with large high-order θ terms

$$J'(\theta) = J(\theta) + \text{Penalty}$$

Definitions

- Two most common

- L1 regularization

- lasso regression

$$J_{L1}(\theta) = \frac{1}{2n} \sum_{i=1}^n (h_{\theta}(x^i) - y^i)^2 + \lambda \sum_{j=1}^m |\theta_j|$$

- L2 regularization

- ridge regression
- weight decay

$$J_{L2}(\theta) = \frac{1}{2n} \sum_{i=1}^n (h_{\theta}(x^i) - y^i)^2 + \lambda \sum_{j=1}^m \theta_j^2$$

Regularized Regression

$$J_{L1}(\theta) = \frac{1}{2n} \sum_{i=1}^n (h_{\theta}(x^i) - y^i)^2 + \lambda \sum_{j=1}^m |\theta_j|$$

$$J_{L2}(\theta) = \frac{1}{2n} \sum_{i=1}^n (h_{\theta}(x^i) - y^i)^2 + \lambda \sum_{j=1}^m \theta_j^2$$

- Find the best fit
- Keep the θ_j terms as small as possible.
- λ is a user-set parameter which controls the trade off

Regularized Regression

$$J_{L1}(\theta) = \frac{1}{2n} \sum_{i=1}^n (h_{\theta}(x^i) - y^i)^2 + \lambda \sum_{j=1}^m |\theta_j|$$

$$J_{L2}(\theta) = \frac{1}{2n} \sum_{i=1}^n (h_{\theta}(x^i) - y^i)^2 + \lambda \sum_{j=1}^m \theta_j^2$$

- Size of λ important
 - λ too high \Rightarrow no fitting
 - λ too low \Rightarrow no regularization

QUESTIONS