

Cotard's Syndrome

Le Délire des Négations · The Walking Corpse Delusion

“

I have no brain, no nerves, no chest, no stomach, no intestines ... all that is left is the rotten skin and the disorganized bones of a decomposing body.”

—Patient described by Jules Cotard, 1880

Table of Contents

A	Diagnostic Criteria	Slides 3–6
B	History & Background	Slide 7
C	Epidemiology	Slide 8
D	Etiology	Slides 9–11
E	Treatment, Case & Stigma	Slides 12–15
F	References	Slide 17–18

Diagnostic Criteria

DSM-5-TR & ICD-11 STATUS

Cotard's is **not** a standalone diagnosis in the DSM-5-TR or ICD-11. It is coded under:

- **Delusional Disorder, somatic type**
DSM-5-TR 297.1 / ICD-11 6A24
- **Major Depressive Disorder with psychotic features**
When mood symptoms dominate
- **Schizophrenia spectrum or organic psychosis**

Three Clinical Subtypes

Psychotic Depression

Nihilistic mood, mild delusions

Cotard Type I

"Pure" delusional, less depressive

Cotard Type II

Mixed: anxiety, immortality, suicidality

CORE CLINICAL FEATURES

Nihilistic delusions

About the body (no organs, no blood) or existence ("I am dead")

Severe depressive mood

Often with psychomotor retardation, mutism, or stupor

Paradoxical immortality beliefs

Patient believes they cannot die—yet may attempt suicide

Self-neglect

Refusing food ("the dead don't eat") or medical care

(Berrios & Luque, 1995)

Diagnostic Framework

DELUSIONAL DISORDER CRITERIA

- A.** ≥1 delusion for ≥1 month
- B.** Criterion A for schizophrenia never met (no prominent hallucinations, disorganized speech, etc.)
- C.** Functioning not markedly impaired apart from delusion's impact
- D.** Any mood episodes brief relative to delusional periods
- E.** Not attributable to a substance or another medical condition

(American Psychiatric Association, 2022)

DIFFERENTIAL DIAGNOSIS


- **Capgras Delusion**
External—familiar people experienced as impostors. Same disconnection mechanism, opposite direction.
- **MDD with Psychotic Features**
Mood-congruent delusions, less systematized. Cotard's features more elaborate nihilistic content.
- **Schizophrenia**
Includes hallucinations and formal thought disorder; Cotard's delusions are more isolated and fixed.
- **Depersonalization-Derealization**
Patient knows the feelings of unreality are not literally true—no fixed delusional belief.

Same neural disconnection mechanism (fusiform–limbic), pointed inward in Cotard's, outward in Capgras.

Symptom Profile

Figure 1

Frequency of Symptoms in 100 Documented Cases of Cotard's Syndrome



Note. Symptom prevalence based on factor analysis of 100 historical cases. Adapted from "Cotard's syndrome: Analysis of 100 cases," by G. E. Berrios and R. Luque, 1995, *Acta Psychiatrica Scandinavica*, 91(3), p. 187.

KEY INSIGHTS

89%

of cases present with depressive mood—the strongest psychiatric link.

86% / 69%

believe parts of their body are absent or that they don't exist at all.

55%

hold paradoxical delusions of immortality—the central contradiction.

Creative Section

A simulated journal entry: what Cotard's might sound like from the inside

Tuesday—I think.

Mom keeps telling me to eat. I keep telling her there's no reason to. The food just sits in the bowl. I know this sounds wrong, but I don't have a stomach anymore. I haven't for a few weeks now.

I look in the mirror and the face is mine but the eyes belong to someone who isn't there. The doctor showed me the EKG—said “your heart is fine”—but the line on the paper isn't proof of anything. Paper can lie. Machines can lie. I can feel that I am not feeling anything, and that is the only thing I trust.

The strangest part is the certainty. People keep asking me “don't you think it's strange that you can walk around if you're dead?” and I want to tell them: yes, of course it's strange. But being dead doesn't mean what you think it means. It means the inside is gone. It means I will be like this forever—that's the worst part. I cannot die because there's nothing left to die.

Tomorrow Mom is taking me to a new psychiatrist. I'll go. I don't think it will help. But I will go.

Fictional composite drawing on documented features (Berrios & Luque, 1995; Dieguez, 2018).

History & Background

1880 ●

Jules Cotard describes "Mademoiselle X"

French neurologist presents a 43-year-old woman who denies the existence of God, the devil, and several parts of her body, and refuses food. He calls it le délire des négations (Berrios & Luque, 1995).



1882 ●

Coined as a syndrome

Cotard formally separates the condition from melancholia, framing it as a form of agitated, anxious depression rather than a discrete disorder (Debruyne et al., 2009).

1893 ●

Renamed Cotard's Syndrome

After Cotard's death (1889), Emil Régis names the condition in his honor; adoption spreads through European psychiatry (Berrios & Luque, 1995).

1995 ●

Berrios & Luque reframe it

A 100-case statistical analysis in Acta Psychiatrica Scandinavica identifies three subtypes and treats Cotard's as a syndrome cluster rather than a single disorder.

2026 ●

Still not a DSM diagnosis

DSM-5-TR and ICD-11 continue to subsume Cotard's under delusional disorder or psychotic depression; debate over standalone status continues (American Psychiatric Association, 2022).

Epidemiology

C · EPIDEMIOLOGY

BREAKDOWN BY UNDERLYING DIAGNOSIS



Synthesized from Berrios & Luque (1995) and Ramirez-Bermudez et al. (2010); approximate proportions.

0.57%

Overall prevalence

psychogeriatric inpatients,
Hong Kong (n = 349)

Chiu, 1995

3.2%

Severe late-life depression

rate climbs with age and
depression severity

Debruyne et al., 2009

M = 52

Mean age of onset

SD = 14.5; range 16–81 yrs
(n = 100)

Berrios & Luque, 1995

≈ 1 : 1

Male : Female ratio

no significant gender difference
(n = 100)

Berrios & Luque, 1995

TWO SUBGROUP COMPARISONS

By age: Cotard's risk increases with age. Berrios & Luque (1995) and Huarcaya-Victoria et al. (2020) both found major depression and organic brain disorders are significantly more common in older Cotard's cases ($p < 0.001$). Nearly absent in pre-pubertal children—only a handful of pediatric cases ever reported.

By underlying diagnosis: roughly 60–70% of cases occur within psychiatric illness (psychotic depression, schizophrenia spectrum); the remainder are linked to organic neurological conditions including TBI, multiple sclerosis, brain tumors, epilepsy, migraine, and adverse reactions to valacyclovir, bridging psychiatry and neurology (Ramirez-Bermudez et al., 2010).

Etiology · Biological Basis

STRUCTURAL & FUNCTIONAL

- **Frontal–parietal–temporal hypoperfusion**

SPECT scans show reduced blood flow in non-dominant hemisphere; abnormalities resolve after ECT (Petracca et al., 1995).

- **Median frontal lobe atrophy**

Higher rates of cerebral atrophy than controls (Joseph & O'Leary, 1986).

- **Fusiform gyrus involvement**


Disrupts self-recognition; sister mechanism to Capgras delusion.

NEUROCHEMISTRY

Dopamine: reduced striatal D2 receptor binding; Cotard's responds to dopamine-modulating antipsychotics like clozapine (De Risio et al., 2004).

Serotonin: low serotonin tone consistent with the strong depression overlap; SSRIs help in depressive subtypes.

IMPLICATED REGIONS



1 Frontal lobe median atrophy & hypoperfusion


2 Parietal lobe right > left lesion concentration

3 Temporal lobe secondary involvement

Non-dominant hemisphere bias across studies (Joseph & O'Leary, 1986; Petracca et al., 1995).

Biological Mechanisms

TWO-FACTOR MODEL OF DELUSION · Coltheart (2007)



ORGANIC TRIGGERS (catalogued causes)

- **Anti-NMDA receptor encephalitis**
autoantibodies against NMDA receptors → cortical/limbic dysregulation; reversible with immunotherapy (Taib et al., 2022)
- **Valacyclovir / aciclovir neurotoxicity**
iatrogenic; CMMG metabolite accumulates in renal impairment, crosses BBB (Hellden et al., 2007)
- **Traumatic brain injury / parieto-temporal lesions**
classic right-hemisphere lesion sites (Young et al., 1992)
- **Multiple sclerosis, brain tumors, epilepsy, migraine**
especially involving non-dominant frontal–parietal regions
- **Severe psychotic depression**
most common psychiatric trigger; HPA-axis dysregulation implicated

CONVERGING MECHANISMS

Predictive coding failure

Brain constantly predicts interoceptive signals (heartbeat, gut, breath). When right insula and anterior cingulate fail, predicted body signals never arrive — the brain's best inference becomes "I have no body" (Seth et al., 2012).

Fusiform–limbic disconnection

Mirror image of Capgras: the fusiform face area still recognizes the self in the mirror, but the affective tag from the amygdala/insula is missing. Patient sees themselves but feels nothing.

HPA-axis hyperactivation

Severe depression elevates cortisol chronically → hippocampal atrophy, prefrontal hypoactivity. Treating the depression often resolves the delusion downstream.

Etiology · Cognitive Models

THE “MIRROR” MODEL · Young & Leafhead (1996)

Both Cotard's and Capgras delusions follow when perceptual recognition is intact but the limbic emotional response is disconnected—the brain confabulates an explanation.

CAPGRAS DELUSION

“This person looks like my mother—but feels unfamiliar. So she must be an impostor.”

Trigger: loss of emotional response to *external* familiar faces.
(“Who is this guy?”)

Confabulation: blame the world—“she's been replaced.”

COTARD'S DELUSION

“This face in the mirror looks like mine—but I feel nothing. So I must be dead.”

Trigger: loss of emotional response to the *self*.
(“My own body feels alien.”)

Confabulation: blame the self—“I must be dead.”

Internalized attributional style (Young & Leafhead, 1996): patients with depression already attribute everything to themselves; this cognitive bias channels the disconnection signal toward “I am dead” rather than “the world is fake.”

Psychodynamic view: extreme guilt and self-loathing produce a defensive “death of the self,” a symbolic punishment for imagined moral failure (Debruyne et al., 2009).

Treatment Approaches

01 Pharmacotherapy

Combination strategy

- **Antidepressant + Antipsychotic**

SSRI (e.g. sertraline) paired with olanzapine or clozapine; targets both mood and delusion.

- **Lithium augmentation**

Effective in depressive Cotard's (Terao et al., 1994).

- **Mechanism**

Restores serotonin / dopamine balance;
lifts depression so the delusion follows.

- **Limitation**

Effective in only ~50% of severe cases—
many are treatment-resistant.

02 Electroconvulsive Therapy

Most reliably effective treatment

- **What it does**

Brief generalized seizure under anesthesia; restores frontal-parietal-temporal blood flow on follow-up SPECT.

- **First-line for treatment-resistant cases**

Particularly effective in severe psychotic depression with Cotard's features (Moschopoulos et al., 2016).

- **Course**

Typically 6–12 sessions over 2–4 weeks; remission often within days.

- **Side effects**

Short-term memory loss, headache; reversible in most cases.

Treatment Approaches

STEPPED-CARE ALGORITHM

STEP 1

Combination Pharmacotherapy

SSRI + atypical antipsychotic (e.g., sertraline + olanzapine); ± lithium augmentation for depressive subtype (Terao et al., 1994). Trial 4–6 weeks.

response

≈
50%



STEP 2

Switch / Augment

Change antipsychotic class (e.g., to clozapine); add lithium or thyroid hormone if mood-driven; reassess underlying diagnosis.

additional

+15–
20%



STEP 3

Electroconvulsive Therapy

First-line for treatment-resistant or life-threatening cases (refusal to eat, suicidality). 6–12 bilateral sessions over 2–4 weeks; SPECT-confirmed mechanism (Petracca et al., 1995).

remission

70–
90%



STEP 4

Maintenance & Supportive Care

Continue meds 12+ months; consider maintenance ECT for relapses; CBT-p, family psychoeducation, safety planning, nutritional rehab.

stability

Long
term

Case Study & Long-Term Outlook

CASE · COTARD'S SYNDROME · Petracca et al. (1995)

“He believed his stomach had vanished.”

Patient: 27-year-old man presenting with Cotard's syndrome arising in schizophreniform disorder (Cotard's is not its own DSM diagnosis—it sits within an underlying disorder).

Cotard's features: nihilistic somatic delusions (“no stomach,” “no cardiovascular function”); refused food; lost 12 kg in 1 month from self-neglect.

Imaging: MRI showed lateral ventricle enlargement; SPECT revealed hypoperfusion in left frontal, parietal, and temporal lobes.

Course: antipsychotics failed → ECT initiated → full remission of nihilistic delusions. Follow-up SPECT showed normalized perfusion. Maintained on olanzapine.

Outcome: remission at 4-month follow-up; no recurrence; returned to work and social functioning.

Significance: first case to document brain perfusion changes before AND after ECT, providing direct evidence that the treatment normalizes the underlying neural dysfunction.

FINDING TREATMENT

Often presents as a psychiatric **emergency**—refusal to eat or suicidality. Care path: primary care or ER → inpatient psychiatry → ECT referral if treatment-resistant.

LONG-TERM HEALTH RISKS

- Malnutrition, dehydration
- Suicide (paradox: “already dead” + unbearable suffering)
- Cognitive decline if untreated
- Recurrence with depression relapse

Prognosis: good when treated early—psychiatric origins respond well; organic lesions may leave residual symptoms.

Media & Stigma

IN POPULAR CULTURE

Figure 2

Hannibal: "Buffet Froid"



Note. From (Fuller, 2013, NBC, S1 E10).

✓ **Right:** Captures appetite loss, ECT as treatment, dissociation from one's own face.

✗ **Wrong:** Conflates Cotard's with violent murder; links a rare delusion to dangerousness.

Figure 3

Synecdoche, New York



Note. From (Kaufman, 2008, film).

✓ **Right:** Protagonist named Caden Cotard; captures existential despair & dissolution of self.

✗ **Wrong:** Metaphorical rather than clinical; viewers may not recognize it as a real disorder.

Figure 4

Scrubs: "My Bright Idea"



Note. From (Lawrence, 2005, ABC, S4 E14).

✓ **Right:** Brief but accurate mention of a patient who believes he died years earlier.

✗ **Wrong:** Used as a one-line gag, no exploration of treatment or lived experience.

STIGMA TODAY

Decreasing: Cotard's is too rare to carry a strong public stereotype, and popular neuroscience writing (Eagleman, 2011; Sacks, 1985) frames it as a window into self-awareness rather than horror. **Still needed:** depictions of recovery (not just symptoms), reducing ECT stigma, and teaching that delusions are clinical symptoms rather than character failings.

Five things to remember about Cotard's Syndrome

- 01 It is a syndrome, not a DSM diagnosis**
Always sits beneath delusional disorder, psychotic depression, or organic psychosis.
- 02 The signature is paradox**
“I am dead” + “I cannot die”—the contradiction is what distinguishes it from depression alone.
- 03 It bridges psychiatry and neurology**
Two-thirds psychiatric, one-third organic—rare clinical evidence of mind/brain unity.
- 04 Cotard's mirrors Capgras**
Same disconnection between recognition and emotion, pointed inward instead of outward.
- 05 ECT remains the most reliable cure**
Brain imaging confirms—perfusion deficits resolve after treatment.

Cotard's Syndrome reminds us that even "being alive" is something the brain has to construct—and that construction can break.

References

- American Psychiatric Association. (2022). *Diagnostic and statistical manual of mental disorders* (5th ed., text rev.). American Psychiatric Association Publishing. <https://doi.org/10.1176/appi.books.9780890425787>
- Berrios, G. E., & Luque, R. (1995). Cotard's syndrome: Analysis of 100 cases. *Acta Psychiatrica Scandinavica*, 91(3), 185–188. <https://doi.org/10.1111/j.1600-0447.1995.tb09764.x>
- Chiu, H. F. K. (1995). Cotard's syndrome in psychogeriatric patients in Hong Kong. *General Hospital Psychiatry*, 17(1), 54–55. [https://doi.org/10.1016/0163-8343\(94\)00066-M](https://doi.org/10.1016/0163-8343(94)00066-M)
- Debruyne, H., Portzky, M., Van den Eynde, F., & Audenaert, K. (2009). Cotard's syndrome: A review. *Current Psychiatry Reports*, 11(3), 197–202. <https://doi.org/10.1007/s11920-009-0031-z>
- De Risio, S., De Rossi, G., Sarchiapone, M., Camardese, G., Carli, V., Cuomo, C., Di Giannantonio, M., Janiri, L., & Bria, P. (2004). A case of Cotard syndrome: 123I-IBZM SPECT imaging of striatal D2 receptor binding. *Psychiatry Research: Neuroimaging*, 130(1), 109–112. <https://doi.org/10.1016/j.psychresns.2003.10.003>
- Dieguez, S. (2018). Cotard syndrome. *Frontiers of Neurology and Neuroscience*, 42, 23–34. <https://doi.org/10.1159/000475679>
- Eagleman, D. (2011). *Incognito: The secret lives of the brain*. Pantheon Books.
- Huarcaya-Victoria, J., Bojórquez-De la Torre, J., & De la Cruz-Oré, J. (2020). Factor structure of Cotard's syndrome: Systematic review of case reports. *Revista Colombiana de Psiquiatría*, 49(3), 187–193. <https://doi.org/10.1016/j.rcp.2018.10.008>

References

- Joseph, A. B., & O'Leary, D. H. (1986). Brain atrophy and interhemispheric fissure enlargement in Cotard's syndrome. *Journal of Clinical Psychiatry*, 47(10), 518–520.
- Moschopoulos, N. P., Kaprinis, S., & Nimatoudis, J. (2016). Cotard's syndrome: Case report and a brief review of the literature. *Psychiatriki*, 27(4), 296–302. <https://doi.org/10.22365/jpsych.2016.274.296>
- Petracca, G., Migliorelli, R., Vázquez, S., & Starkstein, S. E. (1995). SPECT findings before and after ECT in a patient with major depression and Cotard's syndrome. *Journal of Neuropsychiatry and Clinical Neurosciences*, 7(4), 505–507.
- Ramirez-Bermudez, J., Aguilar-Venegas, L. C., Crail-Melendez, D., Espinola-Nadurille, M., Nente, F., & Mendez, M. F. (2010). Cotard syndrome in neurological and psychiatric patients. *Journal of Neuropsychiatry and Clinical Neurosciences*, 22(4), 409–416. <https://doi.org/10.1176/jnp.2010.22.4.409>
- Sacks, O. (1985). *The man who mistook his wife for a hat and other clinical tales*. Summit Books.
- Taib, N. I. A., Wahab, S., Khoo, C. S., Tan, H. J., Kamaruzaman, L., Woon, L. S.-C., & Gan, L. L. Y. (2022). Case report: Cotard's syndrome in anti-NMDA receptor encephalitis. *Frontiers in Psychiatry*, 13, 779520. <https://doi.org/10.3389/fpsy.2022.779520>
- Terao, T., Fujino, A., Egashira, K., & Abe, K. (1994). Lithium-antidepressant combination in the treatment of depressive Cotard's syndrome. *Clinical Neuropharmacology*, 17(3), 270–272.
- Young, A. W., & Leafhead, K. M. (1996). Betwixt life and death: Case studies of the Cotard delusion. In P. W. Halligan & J. C. Marshall (Eds.), *Method in madness: Case studies in cognitive neuropsychiatry* (pp. 147–171). Psychology Press.