



Covariant Formulation of 2D Chiral Scalars



Calvin YR Chen
LeCosPA

18th Apr 2025
NTU String Theory Seminar

Covariant Formulation of 2D Chiral Scalars



based on 2501.16463 in collaboration with E. Joung, K. Mkrtchyan, and J. Yoon

Warm-Up: Maxwell Theory

Electromagnetic Duality

Consider the theory for gauge field $A \in \Omega^1(M)$ on a 4-manifold (M, g) , described by the **Maxwell action**

$$S_{\text{Maxwell}} = \int_M -\frac{1}{2} F \wedge \star F, \quad F = dA$$

The field strength obeys:

Bianchi Identity

$$d^2 = 0 \longrightarrow dF = 0$$

Equation of Motion

$$\delta S = 0 \longrightarrow d \star F = 0$$

- **Electromagnetic Duality** refers to invariance of equations under

$$F \mapsto \star F, \quad \star F \mapsto -F$$

- Here: Enhancement to invariance of equations under $(S)O(2)$ rotations

$$\begin{pmatrix} F \\ \star F \end{pmatrix} \mapsto \begin{pmatrix} \cos(\beta) & \sin(\beta) \\ -\sin(\beta) & \cos(\beta) \end{pmatrix} \begin{pmatrix} F \\ \star F \end{pmatrix}$$

Off-Shell Duality Invariance

Switch to vector notation

$$E_i = F_{0i}, \quad B^i = \frac{1}{2} \epsilon^{ijk} F_{jk}$$

- Action takes the form

$$S_{\text{Maxwell}} = \frac{1}{2} \int d^4x (|\mathbf{E}|^2 - |\mathbf{B}|^2)$$

- **Duality rotation** is now

$$\begin{pmatrix} \mathbf{E} \\ \mathbf{B} \end{pmatrix} \mapsto \begin{pmatrix} \cos(\beta) & \sin(\beta) \\ -\sin(\beta) & \cos(\beta) \end{pmatrix} \begin{pmatrix} \mathbf{E} \\ \mathbf{B} \end{pmatrix}$$

→ Action only invariant under **hyperbolic** rotations?

Is EM duality only a “symmetry” of the **equations of motion**?

Key insight: *“Formal transformation [...] only meaningful [...] if it can be implemented at the level of the basic field variable [...].”*

Off-Shell Duality Invariance

- In **first-order** form:

$$S_{\text{Maxwell}} = \frac{1}{2} \int d^4x \left(|\dot{\mathbf{A}}^T|^2 - |\nabla \times \mathbf{A}^T|^2 + |\dot{\mathbf{A}}^L - \nabla A_0|^2 \right)$$

→ Third term is constraint, and e.o.m. is $\ddot{\mathbf{A}}^T = \nabla^2 \mathbf{A}^T$.

Now consider following transformation

$$\delta \mathbf{A}^T = -\beta \nabla^{-2} \nabla \times \dot{\mathbf{A}}^T$$

- This generates

$$\delta \mathbf{E} = \delta \dot{\mathbf{A}}^T = -\beta \nabla^{-2} \nabla \times \ddot{\mathbf{A}}^T, \quad \delta \mathbf{B} = \nabla \times \delta \mathbf{A}^T = \beta \mathbf{E}$$

→ **Duality rotation on-shell.**

- Changes action by **total time derivative**, and generated by (time-local) **conserved charge**.

→ Duality invariance is off-shell **symmetry**, BUT have to break **manifest covariance** to see this.

Twisted Self-Duality

Can make **duality** manifest using **doubled field content**.

- Introduce extra index $a \in \{1, 2\}$ on gauge fields A^a . Then

$$F^a = dA^a \quad \longrightarrow \quad dF^a = 0$$

→ By construction, **invariant** under $SO(2)$ rotations in “extra directions”.

- Would like to identify

$$F^1 = \star F, \quad F^2 = F$$

such that **EM duality** is equivalent to condition:

$$\star F^2 = F^1$$

→ This is **twisted self-duality relation**: Implies original equations of motion.

Manifest Duality

Duality invariance made manifest by following **local action**

$$S_{\text{Schwarz-Sen}} = -\frac{1}{2} \int d^4x \left(\delta_{ab} \mathbf{B}^a \cdot \mathbf{B}^b - \varepsilon_{ab} \mathbf{E}^a \cdot \mathbf{B}^b \right)$$

[Schwarz & Sen '93]

[Henneaux & Teitelboim '11]

- Equation of motion for \mathbf{A}^2 sets $\mathbf{B}^2 = \mathbf{E}^1$ (up to gauge transformations). Plugging back into action, recover

$$S_{\text{Schwarz-Sen}} = -\frac{1}{2} \int d^4x \left(\mathbf{B}^1 \cdot \mathbf{B}^1 - \mathbf{E}^1 \cdot \mathbf{E}^1 \right)$$

→ **Maxwell action.**

- Manifestly **gauge invariant**.
- Not manifestly **covariant**, but it is: Manifest invariance under spatial rotations but not **boosts** – latter changes action, but by boundary term.

Central **tension** between manifest **duality** and **covariance**

(Twisted) Self-Duality

Are there other examples of (twisted) self-duality relations?

Linearised Gravity

For example, consider **linearised gravity** in D dimensions.

- In vacuum, linearised Riemann tensor satisfies

Bianchi Identity

$$R_{\mu[\nu\rho\sigma]} = R_{\mu\nu[\rho\sigma,\lambda]} = 0$$

Equation of Motion

$$R_{\mu\nu} = 0$$

- Same relations satisfied by linearised **dual Riemann** tensor

$$\tilde{R}_{\mu\nu\rho\sigma} = \frac{1}{2}\epsilon_{\mu\nu\alpha\beta}R^{\alpha\beta}{}_{\rho\sigma}$$

- **Gravitational Duality**: $R \mapsto \tilde{R}$ (enhanced to full rotations in $D = 4$) **off-shell symmetry** of theory – can phrase as twisted self-duality.

[Henneaux & Teitelboim '05; Bünster, Henneaux, and Hörtner '13]

→ Make manifest at expense of **covariance**.

- Similar holds for **(partially) massless higher-spin** fields.
- **Non-linearities**: Exchanges Schwarzschild mass with **Taub-NUT** parameter.

Form Fields

Most straightforward **generalisation** to Maxwell theory.

[Henneaux '85]

Theory of p -form gauge field $A \in \Omega^p(M)$ on manifold (M, g) with $\dim(M) = d$, described by action

$$S = \int_M -\frac{1}{2} F \wedge \star F, \quad F = dA$$

- Numerous examples in **string theory/supergravity**: $p = 2$ in Kalb-Ramond sector, and $p = 1, 3$ and $p = 0, 2, 4$ in type-IIA and -IIB respectively.
- **Electromagnetic Duality**: $F \mapsto \star F$ (enhanced to full rotations in $d = 2p + 2$) **off-shell symmetry** – can phrase as twisted self-duality relation.

→ Make manifest at expense of **covariance**.

[Bünster & Henneaux '11]

(Anti-) Self-Duality

Can also have **(anti-) self-duality** relations!

Again, consider p -form $A \in \Omega^p(M)$ satisfying

$$dA = \sigma \star dA, \quad \sigma = \pm 1$$

- For consistency (in Lorentzian signature), need $d = 2(p + 1)$, $p \in 2\mathbb{N}_{>0}$.
- Use null coordinates two-dimensional (Lorentzian) submanifold

$$dA = \partial_- A dx^- + \partial_+ A dx^+$$

Then $\star dx^\pm = \pm dx^\pm$, so **(anti-) self-duality** amounts to

$$\partial_\pm A = 0$$

→ **Chirality** condition: Halves degrees of freedom!

- Notorious example: Self-dual 4-form in **type-IIB** supergravity.

(Twisted) Self-Duality

Twisted self-duality most general one-derivative eq. for bosonic fields.

- For particular (D, p) find (anti-) self-duality/EM duality: **Dimensional reduction** of chiral fields in $D = 4k + 2$ to $D = 4k$ gives EM duality.
- **Democratic actions**: Twisted self-duality as e.o.m. (from latter).
- Expect two-derivative eq. \rightarrow difficult to find **covariant actions!**
 - For **gauge fields**: Actions with manifest EM duality.
 - For **chiral fields**: Describing right degrees of freedom. *E.g.* naïve action

$$S = \int_M F \wedge \star F + \lambda \wedge (F - \sigma \star F)$$

wrong...

Democratic Actions

Americans for Democratic Action



Democratic Formulations

Fields obeying **(twisted) self-duality relations** clearly ubiquitous. For quantisation, convenient to have **actions**.

- Correct degrees of freedom, nothing manifest.

[Siegel '84; Tseytlin '90; Floreanini & Jackiw '87]

- **Duality invariance**, no covariance.

[Schwarz & Sen '94; Henneaux & Teitelboim '11]

- Manifest **duality** and **covariance**.

[Pasti, Sorokin, and Tonin '97; Sen '15; Mkrtchyan '19]

Unifying picture: Descent from **topological field theory** in one higher dimension.

[Arvanitakis, Cole, Hulik, Sevrin, and Thompson '22]

- Description of EM, p-forms and other gauge theories from CS/BF theory.

→ Straightforward generalisation to **interactions**.

Example: Maxwell Theory

Consider $H_1, H_2 \in \Omega^2(M)$ with $\dim(M) = 5$ described by **BF-type action**

$$S = \int_M -H_2 \wedge dH_1 + dH_2 \wedge H_1 - \frac{1}{2} \int_{\partial M} H_a \wedge \star H_a$$

- Bulk equations of motion:

$$dH_a = 0$$

→ **Pure gauge** (non-dynamical) in bulk.

- Boundary equation of motion:

$$H_1 = \star H_2$$

→ **Twisted self-duality!**

Idea: *Covariant reduction to the boundary should lead to covariant action of boundary degrees of freedom!*

Topological Reduction

Trick: Decompose

$$H_a = H_a^\perp + v \wedge H_a^\parallel$$

for closed, constant, and nowhere-null $v \in \Omega^1(M)$ — defines **foliation**.

- H_a^\parallel acts as Lagrange multiplier for **constraint**

$$v \wedge dH_a^\perp = 0$$

→ **General** solution

$$H_a^\perp = dA_a + v \wedge S_a \longrightarrow H_a = dA_a + v \wedge R_a$$

- Plug back into action: Obtain **democratic action** in **Mkrtchyan form**

$$S = \int_{\partial M} \left[-\frac{1}{2} \delta^{ab} H_a \wedge \star H_b - \varepsilon^{ab} v \wedge R_a \wedge dA_b \right]$$

[Mkrtchyan '19]

**2D chiral fields can be described as
boundary degrees of freedom of 3D gravity**

Goal: Do this!

3D (Higher-Spin) Gravity

Focus on the gravitational sector for now

Frame-Like Formulation

For gravity in **first-order form**, fundamental fields: vielbein $\{e^a\}$ and connection $\{\omega^a{}_b\}$.

→ In $D = 3$, useful to dualise

$$\omega^a = \frac{1}{2}\epsilon^a{}_{bc}\omega^{bc}$$

- **Einstein-Hilbert action** with $\Lambda = -1/\ell^2 < 0$

$$S_{\text{EH}} = \frac{1}{8\pi G} \int_M e^a \wedge \left[d\omega^a + \frac{1}{2}\epsilon^a{}_{bc} \left(\omega^b \wedge \omega^c + \frac{1}{3\ell^2} e^b \wedge e^c \right) \right]$$

- Equations of motion are vanishing of **torsion** and **curvature** respectively:

$$\mathcal{T}^a := de^a + \omega^a{}_b \wedge e^b = 0,$$

$$\mathcal{R}^a{}_b := d\omega^a{}_b + \epsilon^a{}_{bc} (\omega^b \wedge \omega^c + e^b \wedge e^c) = 0$$

→ Reproduce **Einstein's equation** with

$$g_{\mu\nu} = e^a{}_\mu e^b{}_\nu \eta_{ab}$$

Chern-Simons Formulation

Can write in **Chern-Simons** formulation.

- Define $\mathfrak{sl}(2, \mathbb{R})$ -valued **gauge fields**

$$A^a = \omega^a + \frac{e^a}{\ell}, \quad \tilde{A}^a = \omega^a - \frac{e^a}{\ell}$$

→ Useful **basis** of generators is

$$[L_m, L_n] = (m - n)L_{m+n} \longrightarrow A = \sum_{n=-1}^{+1} A^{(n)} L_n$$

Identifying $k = \ell/4G$,

$$S_{\text{EH}} = S_{\text{CS}}[A] - S_{\text{CS}}[\tilde{A}], \quad S_{\text{CS}}[A] = \frac{k}{4\pi} \int_M \text{Tr} \left(A \wedge dA + \frac{2}{3} A \wedge A \wedge A \right)$$

- Einstein's equation equivalent to **flatness condition** on both field strengths.

→ (Classical) **equivalence** of AdS_3 gravity with $\mathfrak{so}(2, 2) \sim \mathfrak{sl}(2, \mathbb{R}) \oplus \mathfrak{sl}(2, \mathbb{R})$ CS theory.

[Achucarro & Townsend '86; Witten '88]

Asymptotically AdS Fluctuations

Put on **curved background**: Flat background gauge field \underline{A} , *i.e.* for which

$$d\underline{A} + \underline{A} \wedge \underline{A} = 0$$

- Action for **fluctuations**: “Minimal coupling”

$$S[a] = \frac{k}{4\pi} \int_M \text{Tr} \left(a \wedge D a + \frac{2}{3} a \wedge a \wedge a \right), \quad D \cdot = d \cdot + [\underline{A}, \cdot]$$

Most general **asymptotically** AdS₃ solution

$$ds^2 = \frac{\ell^2}{z^2} \left[\left(dx^+ + z^2 \tilde{\mathcal{L}}(x^-) dx^- \right) \left(dx^- + z^2 \mathcal{L}(x^+) dx^+ \right) + dz^2 \right]$$

[Bañados '99]

- Explicitly for chiral part:

$$\underline{A}^z = \frac{dz}{z}, \quad \underline{A}^+ = 2 \frac{dx^+}{z}, \quad \underline{A}^- = 2z \mathcal{L}(x^+) dx^+$$

Incorporate higher-spins now!

Higher-Spin Fields

Why **higher-spins**?

- Fields characterised by **mass** + **spin**: Higher-spin fields natural (in the free case).
 - Natural context: **String theory** and **AdS/CFT** (e.g. \mathcal{W}_N -minimal model holography)
 - Presence of $s > 2$ typically implies existence of spin-2: Higher-spin **gravity**.
 - For interactions: Various **no-go “theorems”**!
 - E.g. in $D \geq 4$ need infinitely many higher spins, e.g. **Vasiliev HS gravity** (dual to vector model)
- Caveat: In $D = 3$, **little group** does not admit arbitrary helicity massless rep. (no dynamical HS bulk degrees of freedom), but good **toy model**.

Fronsdal Field

Massless **spin-s** on curved space described by symmetric rank-s tensor field obeying

$$(\square - m_s^2) \phi_{\mu[s]} + \dots = 0$$

- Invariance under **gauge transformation**

$$\delta \phi_{\mu[s]} = s \nabla_{(\mu_s} \epsilon_{\mu[s-1])}, \quad \epsilon_{\mu[s-3]}{}^\lambda{}_\lambda = 0$$

$$\alpha[n] = \alpha_1 \dots \alpha_n$$

Higher-spin analogues of vielbein and spin connection $\{e^{a_1 \dots a_{s-1}}, \omega^{a_1 \dots a_{s-1}}\}$.

- Fronsdal **action** around spin-2 background, schematically

$$S_{\text{Fronsdal}} \sim \int_M \{e \wedge [d\omega + \underline{\omega} \wedge \omega] + \underline{e} \wedge [\omega \wedge \omega + e \wedge e]\}$$

- E.o.m.: Vanishing of higher-spin **torsion** and **curvature** \rightarrow Fronsdal equation.
- Identification with Fronsdal field via

$$\varphi_{\mu[s]} = \underline{e}^{a_1}{}_{(\mu_1} \dots \underline{e}^{a_{s-1}}{}_{(\mu_{s-1}} e_{\mu_s) a[s-1]}$$

Chern-Simons Formulation

For **CS formulations**, define $\mathfrak{sl}(N, \mathbb{R})$ -valued gauge fields

$$a^{a_1 \dots a_{s-1}} = \omega^{a_1 \dots a_{s-1}} + \frac{1}{\ell} e^{a_1 \dots a_{s-1}}, \quad \tilde{a}^{a_1 \dots a_{s-1}} = \omega^{a_1 \dots a_{s-1}} - \frac{1}{\ell} e^{a_1 \dots a_{s-1}}$$

- Useful **basis** of generators

$$[L_m, W_n^s] \sim W_{m+n}^s, \quad \text{Tr} \left(W_m^s W_n^{s'} \right) \sim \delta_{m, -n} \delta_{s, s'} \longrightarrow a = \sum_{n=-(s-1)}^{s-1} a^{(s, n)} W_n^{(s)}$$

- Fronsdal action** now

$$S_{\text{Fronsdal}}[e, \omega] = S_{\text{CS}}^{(2)}[a] - S_{\text{CS}}^{(2)}[\tilde{a}]$$

$$S_{\text{CS}}^{(2)}[a] = \frac{k}{4\pi} \int_M \text{Tr} (a \wedge \text{D}a), \quad \text{D}a = \text{d}a + \underline{A} \wedge a + a \wedge \underline{A}$$

Description of massless spin-s around on-shell **gravitational background**.

→ *E.g.* includes linearised gravity for spin-2!

Covariant Reduction to Boundary

Boundary Theory

Focus on the **linearised** theory.

For full reduction, careful about **boundary terms**.

- For any $\mathfrak{sl}(N, \mathbb{R})$

$$S[a, \tilde{a}] = S_L[a] - S_R[\tilde{a}]$$

with opposite sign

$$S_L[a] = \frac{k}{4\pi} \left[\int_M \text{Tr} (a \wedge D a) - \frac{1}{2} \int_{\partial M} \text{Tr} (a \wedge \star a) \right]$$
$$S_R[\tilde{a}] = \frac{k}{4\pi} \left[\int_M \text{Tr} (\tilde{a} \wedge D \tilde{a}) + \frac{1}{2} \int_{\partial M} \text{Tr} (\tilde{a} \wedge \star \tilde{a}) \right]$$

- Correct **boundary equation**/condition: **Chirality condition**

$$(a - \star a) \big|_{\partial M} = 0$$

Topological Reduction of Linearised Theory

Follow topological **boundary reduction** procedure from before.

- Trick: Decompose

$$a = b + v \psi$$

- Field ψ acts as Lagrange multiplier enforcing **constraint**

$$v \wedge D b = 0 \quad \longrightarrow \quad a = D\varphi + v\rho$$

- Plug back into action:

$$S_L = -\frac{k}{4\pi} \int_{\partial M} \text{Tr} \left[-\frac{1}{2} (D\varphi + v\rho) \wedge \star (D\varphi + v\rho) - D\phi \wedge v\rho \right]$$

→ **Covariant action!**

Is this the correct action?

Dynamical Content

Connect it to **non-covariant formulations**.

- Integrate auxiliary field ρ out:

$$S_L = -\frac{k}{4\pi} \int d^2x \frac{1}{v_-} \text{Tr} (\varepsilon^{\alpha\beta} v_\alpha D_\beta \varphi D_- \varphi)$$

→ **PST-type action** (covariant + duality symmetric, but non-polynomial)!

- Gauge fixing $v = dx^0$

$$S_L = -\frac{k}{2\pi} \int d^2x \text{Tr} (D_1 \varphi D_- \varphi)$$

→ **FJ-type action**: Up to gauge freedom, equation of motion reduces to:

$$D_- \varphi = 0$$

as expected.

→ **BUT** for right degrees of freedom, need to impose asymptotic boundary conditions.

Boundary Conditions

Fluctuations should be subdominant to background.

- Gauge-fixing $v = dx^0$, **asymptotic AdS conditions:**

$$a_1^{(s,n)} = \mathcal{O}(z^{1-n}), \quad \text{for } n > -(s-1)$$

- Spin-1 reduction: **Floresanini-Jackiw**
- Spin-2 reduction: **(non-chiral) WZW/Liouville** or **Alekseev-Shatashvili**.

[Coussaert, Henneaux, van Driel '95]

[Cotler & Jensen '18]

- **Asymptotic symmetry** algebra: *Drinfeld-Sokolov* reduction of affine- $\mathfrak{sl}(2, \mathbb{R})$ to Virasoro (matching central charge!).

[Brown & Henneaux '86]

- Higher-spin reduction: Toda or **higher-spin generalisation** of AS.
 - **Asymptotic symmetry** \mathcal{W}_N -algebra!

Higher-Order Chiral Scalar

Higher-spin AAdS conditions trivialised by

$$a^{(s,n)} = z^{-n} D \phi^{(s,n)}, \quad D_1 \phi^{(s,n)}(0, x) = 0, \quad \text{for } n > -(s-1)$$

→ Only need half of these.

- Gives **recurrence relation**

$$\phi^{(s,n)} = - \frac{\partial_1 \phi^{(s,n+1)} + (s+n+1) \mathcal{L} \phi^{(s,n+2)}}{s-n-1}$$

→ Solve everything in terms of $\phi^{(s,s-1)}$!

- Plug back into FJ-type action: **HS generalisation of FJ/AS** describing **higher-order scalar**

$$S_L \sim \int_{\partial M} d^2x D_1 \phi^{(s,1-s)} \partial_- \phi^{(s,s-1)} \sim \int_{\partial M} d^2x \mathcal{D}_{\mathcal{L}}^{(2s-1)} \phi^{(s,s-1)} \partial_- \phi^{(s,s-1)}$$

Higher-Order Chiral Scalar

What is this operator $\mathcal{D}_{\mathcal{L}}^{(2s-1)}$?

- **Differential operator** of maximal order $2s - 1$, defined by recurrence relation: *E.g.* for spin-2 and -3

$$\mathcal{D}_{\mathcal{L}}^{(3)} = \partial_1^3 - 2(\partial_1 \mathcal{L} + \mathcal{L} \partial_1),$$

$$\mathcal{D}_{\mathcal{L}}^{(5)} = \partial_1^5 - 2(2 \partial_1^3 \mathcal{L} + 3 \partial_1^2 \mathcal{L} \partial_1 + 3 \partial_1 \mathcal{L} \partial_1^2 + 2 \mathcal{L} \partial_1^3) + 8(3 \partial_1 \mathcal{L}^2 + 2 \mathcal{L} \partial_1 \mathcal{L} + 3 \mathcal{L}^2 \partial_1)$$

- For constant \mathcal{L} : Operator **factorises**

$$\mathcal{D}_{\mathcal{L}}^{(2s-1)} = \partial_1 \prod_{n=1}^{s-1} (\partial_1^2 - 4\mathcal{L}n^2)$$

→ Coincides with **Bol operators**: Invariant under projective action of $\text{SL}(2, \mathbb{R})$.

[Bol '49; Gieres & Theisen '94]

- Physical degrees of freedom are chiral scalar: Tachyonic (massive) scalars in (A)dS₂ with definite-sign momentum (energy).

[Farnsworth, Hinterbichler, and Saha '24]

For non-linear reduction for spin-2 and -3 cf. our paper 😊

Conclusion

Concluding Remarks

Various fields obey **twisted self-duality relations** in theoretical physics.

- For gauge theories, **EM duality** symmetry.
- For chiral fields, (anti-) self-duality \rightarrow **chirality**.

Long history of attempts at **covariant actions** with twisted self-duality relations as equations of motion.

\rightarrow Recent development: **Topological boundary reduction**.

- Applied this to higher-order chiral scalars: Boundary degrees of freedom in AdS_3 HS gravity.

Goal: Repeat for linearised (HS) gravity in $D = 4$.

Thanks for your attention!

Bonus Slides

Non-Linear HS Gravity

Non-linear **HS gravity** is $\mathfrak{sl}(N, \mathbb{R}) \oplus \mathfrak{sl}(N, \mathbb{R})$ CS theory: Massless fields up to spin- N with HS self-interactions.

- Including **boundary terms**:

$$S_{\text{HSG}} = S_L[\mathcal{A}] - S_R[\tilde{\mathcal{A}}], \quad \mathcal{A} = \sum_{n=-1}^1 A^{(n)} L_n + \sum_{s=3}^N \sum_{m=-(s-1)}^{s-1} a^{(s,n)} W_n^{(s)}$$

where

$$S_{L/R}[\mathcal{A}] = \int_M \text{Tr} \left[\mathcal{A} \wedge d\mathcal{A} + \frac{2}{3} \mathcal{A} \wedge \mathcal{A} \wedge \mathcal{A} \right] \pm \int_{\partial M} \text{Tr} [\mathcal{A} \wedge \star \mathcal{A}]$$

- **Bulk** equations of motion: **Flatness** condition, solved by

$$\mathcal{A} = G^{-1} dG = g_\varphi^{-1} \underline{A} g_\varphi + g_\varphi^{-1} dg_\varphi$$

- **Boundary** equation of motion: **Chirality** condition

$$(\mathcal{A} - \star \mathcal{A})|_{\partial M} = G^{-1} \partial_- G = 0$$

Boundary Reduction

Once again interested in **covariant boundary reduction**.

- Action reduces to

$$S_L = \int_{\partial M} \text{Tr} \left[-\frac{1}{2} (g^{-1} dg + v\rho) \wedge \star (g^{-1} dg + v\rho) + v \wedge (\rho + \lambda) g^{-1} dg \right] + S_{\text{WZW}}[g]$$

- Connect to **non-covariant formalisms** by integrating and fixing gauge:

$$S_L = -2 \int_{\partial M} d^2x \text{Tr} [g^{-1} \partial_1 g g^{-1} \partial_- g] - \frac{1}{3} \int_M \text{Tr} [(g^{-1} dg)^3]$$

- Need to supply with **asymptotic conditions**

$$v \wedge \mathcal{A}^{(s,n)} \Big|_{\partial M} = v \wedge \underline{\mathcal{A}}^{(s,n)}, \quad \text{for } n > 1 - s$$

→ In terms of group elements,

$$g^{-1} \partial_1 g = L_{+1}, \quad \text{for } n > -(s-1)$$

Spin-2 Reduction

Consider *e.g.* reduction procedure for spin-2.

- **Gauss decomposition** of group element

$$g_\phi = e^{\varepsilon(L_1 - \mathcal{L}L_{-1})} e^{\sigma L_0} e^{fL_{-1}}$$

- In fund. rep., **asymptotic conditions** are

$$e^{-\sigma} = \partial_1 \chi, \quad f = -\frac{1}{2} \partial_1 \sigma$$

upon which action becomes

$$S_L = \int_{\partial M} d^2x \left(-\frac{\partial_1^2 \chi \partial_- \partial_1 \chi}{(\partial_1 \chi)^2} - 4\mathcal{L} \partial_1 \chi \partial_- \chi \right)$$

→ **Alekseev-Shatashvili action.**

For spin-3, possible but long expressions...