

An Integral Projection Modeling Approach to Demographic Consequences of Multiple Partner Mutualisms

Alexandra Campbell^{1,*}

Tom E.X. Miller^{1,†}

1. Rice University, Houston, Texas 77004;

* Corresponding author; e-mail: amc49@rice.edu.

† Corresponding author; e-mail: tom.miller@rice.edu.

Manuscript elements: Figure 1, figure 2, table 1, appendix A (for print; including figure A1, figure A2, and table A1), supplemental PDF. Figure 2 is to print in color.

Keywords: Examples, model, template, guidelines.

Manuscript type: Article.

Prepared using the suggested L^AT_EX template for *Am. Nat.*

Abstract

Mutualisms are among the most widespread species interactions with diverse and dynamic con-
sequences. They are considered more context dependent than other species interactions, meaning
there are many different factors which influence the impacts of a mutualistic interaction, includ-
ing partner diversity. Partner diversity has become a central focus in the field of mutualisms
in recent years shifting the focus to multispecies mutualisms where the focus was previously
on pairwise studies. It has been shown that pairwise studies are poor predictors of the effects
of multispecies mutualistic interactions. The diversity of partners in a multi-species mutualism
causes varied demographic effects on the population of the focal mutualist which can be ex-
plained by several mechanisms, portfolio effect, complementarity, and sampling effect. In this
paper, I will focus on defensive ant-plant mutualisms. These involve plants which provide extra-
floral nectar and/or “housing” to ants which in turn defend them from herbivores. While these
interactions have been well studied in the literature, few have considered how diversity within
partner guilds affect the overall benefits of mutualism for the plant partner. I use the plant,
Cylindropuntia imbricata (tree cholla), and ant, *Crematogaster opuntiae* (Crem.), *Liometopum*
apiculatum (Liom.), and more, multispecies mutualism in which the cacti provide extrafloral nec-
tar in exchange for defense from various herbivores and seed predators. I used 18 years of data
collected from demographic censuses, which includes data such as size, survival, reproductive
status, flowers produced, ant partner, and herbivory for 8 30x30 m plots, in New Mexico. With
this data I parameterize a series of Bayesian generalized linear vital rate models to determine
the impacts of different partners on the focal mutualists. I found that different ant partners did
have different impacts on the vital rates of the tree cholla. Specifically, Crem. tended plants had
advantages in both growth and survival when small, and Liom. tended plants had advantages
in colonizing partners as well as floral viability. With these models I constructed an Integral
Projection Model in which I could vary the presence of each partner, creating different “diver-
sity scenarios”, to determine under which diversity scenario the focal mutualist experienced the

27 highest fitness, and which of the above mechanisms may explain the effects of partner diversity
in this system. I found that the real-life scenario (all possible ants are present) lead to the highest
fitness for the tree cholla, indicating that partner diversity is beneficial in this system. It also
30 shows that complementarity is at play in this system, meaning different partners offer different
benefits leading to synergistic benefits for the tree cholla.

Introduction

33 Mutualisms are species interactions where all participants benefit, leading to higher individual fitness and increased population growth rates. They are among the most widespread species interactions[Chamberlain2014, BoucherDouglasH.1985] with diverse and dynamic consequences
36 [Bronstein1994, Chamberlain2014, Frederickson2013], often deteriorating into other types of interactions, including parasitism [Rodriguez-Rodriguez2017, Song2020, Mandyam2014, Thrall2007, Bahia2022]. One reason they are so ubiquitous is they take many forms in nature, including defense for food or housing[Willmer1997], pollination and dispersal for food[Sakai2002, Burns2004], resource uptake for housing[Holland2010], etc. Mutualisms are typically considered more context dependent than other species interactions[Chamberlain2014, Frederickson2013], meaning
39 strength of the interaction is often determined by the surrounding environment and species. Historically, pairwise studies, where two species engage in mutualism with only each other, were central in the field of mutualisms, it has since been shown that pairwise studies are poor
42 predictors of the effects of multi-species mutualistic interactions[Afkhami2014, Palmer2010], where more than two species engage in mutualisms. The differences in mutualistic outcomes are due to differences in partner quality[Bascompte2019, Stanton2013, Frederickson2013, Jones2015, Ness2006], differences in the types of benefits offered[Kiers2003, Afkhami2014], and direct interactions between partners[Sun2019, Heath2009, Heath2014, Grutter2003]. For these reasons, pairwise studies cannot be accurately used to predict the outcomes of multi-species mutualisms[Palmer2010,
45 Stanton2013, Chamberlain2014, Song2020]. Studies on the effects of diverse multi-species mutualism systems are necessary to help us understand the demographic effects of partner diversity in mutualisms[Bascompte2019]. Multi-species mutualism partner diversity causes varied demographic effects on the population of the focal mutualist which can be explained by
51 a number of mechanisms. In some cases, the quality of the benefits offered varies leading to some partners which benefit the individual, some which have neither a positive or negative

tive impact on the individual, and some which negatively impact the host¹ [Bronstein1994, Bronstein2001a, Afkhami2014, Song2020, West2007, Frederickson2013, Jones2015]. In cases where the partner offers benefits, pinkthe function, or type of benefits, offered by the partner can vary² [Stanton2003].

[width=0.58]Complementarity copy.png [width=0.40]SamplingEffectcopy.png

Figure 1: asdasdf

[width=0.60]PortfolioEffectcopy.png

Figure 2: asdasdf

This variation in partners can lead to pinknet benefits, no net benefits, or net costs for the focal mutualist³.In this study I focus on the positive effects of partner diversity which can be explained by several mechanisms described below. When there is a consistent partner hierarchy based on benefits offered, a more diverse sample of the partner community may be more likely to include the highest quality partner (the one which offers the focal mutualist the most benefit)[Frederickson2013].When this leads to the fitness of a focal mutualist interacting with multiple partners being equal to the fitness of a focal mutualist interacting with only the highest quality partner, as shown in Figure ??b, sampling effect can explain the positive effects of partner diversity[Batstone2018]. Partners can also vary functionally, meaning each partner offers a different type of reward to the focal mutualist[Stachowicz2005, Bronstein2006, Stanton2003].When this leads to a higher fitness for the focal mutualist than any single interaction,as shown in Figure ??a, complementarity explains the benefits of partner diversity[Batstone2018]. Partners can also have asynchronous responses to the environment, either spatially[Ollerton2006] or temporally[Alarcon2008].Multiple partners can act as a 'portfolio' leading to more even fitness

¹pinkIs this the best way to introduce this or should I talk about quality of benefits compared to cost of rewards? AKA some partners offer a net benefit when the benefits outweigh the cost of rewards to the partner ... etc.

²pinkIs this enough detail or do I need to go into examples?

³pinkIs this the best way to start this paragraph? If so should I include a figure panel?

benefits across environmental or temporal heterogeneity, as shown in Figure ??, through the portfolio effect [Batstone2018, Lazaro2022]. Partner diversity can have different effects depending on whether partners are present all at once or sequentially (partner turnover) [Djieto-Lordon2005, Ness2006, Bruna2014]. Turnover can happen at different timescales [Oliveira1999, Horvitz1986] from minutes to years. The degree and order of partner turnover can have important impacts on the fitness of the focal mutualist. The degree, or rapidity, of partner turnover can impact the level of benefits received by the focal mutualist particularly if the benefits continue to accumulate or if they saturate over time [Sachs2004].⁴ The direction of turnover can also have a significant impact, particularly if the most beneficial partner changes across the ontogeny of the focal mutualists life [Fonseca2003]. For example, susceptibility to enemies can change across life stages [Boege2005, Barton2010] so the focal mutualist benefits the most when defensive partners align with the more vulnerable life stages of focal mutualists [Djieto-Lordon2005]. Ant visitation to extra-floral nectar producing plants are a classic and well-studied example of interspecific mutualisms which are often multispecies and have dynamic turnover patterns⁵, making these interactions great systems to study the effects of partner diversity. A subset of these interactions, defensive ant-plant mutualisms are common interactions in which plants provide food and/or “housing” to ants which in turn defend them from herbivores [Bronstein1998, Bronstein2006]. While these interactions have been well studied [Ness2006, Beattie1985, Schultheiss2022], few have considered how diversity within ant defender guilds [Stanton2013] or temporal fluctuations in partner interactions [Trjelsgaard2015] affect the overall benefits of mutualism for the plant partner. This study focuses on the tree cholla cactus, a long lived *Cylindropuntia imbricata*, EFN-bearing plant that associates with multiple species of ant partners. Insect herbivory negatively affects the growth of cholla, in turn reducing population growth rates [Miller2009]. Ant partners defend the cholla from these insect herbivores, reducing the negative effects of herbivory [Miller2007]. The multiple ant partners do not co-occur on individual plants, but a

⁴*pink* Not quite sure what to say here.

⁵*pink*... Is this a better transition? I tried to tie it in a bit more conceptually.

single cholla may interact with multiple types of partners over its lifetime. While the quality of ant partners in this system has been studied, with some seen as superior defenders and some being viewed as net-negative because they can deter pollinators [Ohm2014], no one has integrated demographic effects over the life cycle of these tree cholla.

[width=.8]herb_{ant}only_{flow}.png

Figure 3: asdasdf

In this study we will answer three questions about the demographic effects of partner diversity across host lifetimes in multi-species mutualisms:

1. What are the contrasting demographic effects of multiple partners?
2. How do host size and partner identity impact the directionality of turnover between multiple partners?
3. What mechanisms explain the effects of partner diversity in a multispecies mutualism?

To answer these questions, we used a 20-year long-term dataset of individual-level demographic information and ant partner identity and quantity data to track the structure of the population across time as well as individual level impacts of ant partners on the cacti. We used vital rate functions to evaluate the first two questions. We then used the composite integral projection model (IPM) to evaluate how vital rate differences affect intrinsic population growth rates and identify mechanisms at play.

Methods

Study System

This study was conducted in the Los Piños mountains, a small mountain chain located on the Sevilleta National Wildlife Refuge, a Long-term Ecological Research (LTER) site in central New Mexico. This is an area characterized by steep, rocky slopes, and perennial vegetation like cacti

and junipers. Tree cholla cacti are common in high Chihuahuan desert habitats, with their native range spanning the southwestern USA[Benson1982]. These arborescent cholla produce cylindrical segments with large spines. In the growing season, May to August in New Mexico, the plants initiate new vegetative segments and flower buds at the ends of existing segments. While most plants produce new segments every season, only those which are reproductively mature produce flower buds. Tree cholla generally reach at least 9 years of age before beginning to reproduce [Ohm2014]. Like other EFN-bearing cacti, tree cholla secrete nectar from specialized glands on young vegetative and reproductive structures[Ness2006, Oliveira1999].

This EFN is harvested by various ant species in return for defense. At the Sevilleta, the cholla are visited primarily by two species of ground-nesting ants from the formicoid clade, *Crematogaster opuntiae* and *Liometopum apiculatum*, as well as other rarer species, including *Forelius pruinosus*, a red *Phenogaster* spp.⁶, and *Camponotus* spp.. *L. apiculatum* are the most frequent visitors with 25% – 60% of tree cholla tended by these ants, followed by *C. opuntiae* visiting between 0% – 20% of cacti[Donald2022] depending on the year. Up to 80% of cacti remain vacant in any given year.

These ants rarely co-occur on a plant, probably due to interspecific competition[Miller2007]. Each cholla is visited by a single ant species for the duration of a season, and the species of the visitors can change from one season to the next. At the beginning of the growing season, when EFN production begins, the ground-nesting ants will begin visiting tree cholla. They will visit the cholla every day during the season around the clock, with the most activity around sunrise or sunset[Ohm2014]. Smaller cholla are less likely to be visited because they produce very little EFN, so larger cholla are generally more highly tended[Miller2014]. In late August, the tree cholla stop producing EFN and the ants vacate until the next growing season.

There are a variety of insect herbivores and seed predators which attack the cholla, focusing either on the vegetative segments and the reproductive segments[Mann1969]. An unidentified weevil of the genus *Gerstaekeria* feeds on vegetative and reproductive structures and im-

⁶redIncorrect spelling.

plants their larvae within the plant tissue for the winter. A cactus bug, *Narnia pallidicornis*, (Hemiptera: Coreidae) feeds on all cholla parts with a preference for the reproductive structures [Miller2006]. A seed predator, *Cahela ponderosella*, (Lepidoptera: Pyralidae) attacks developing fruits pre-dispersal and oviposits in open flowers mid-growing season where larvae burrow into the ripening ovary. These predators can have significant negative impacts on the fitness of individual cholla and depress population growth [Miller2009]. There is experimental evidence that tree cholla tended by *L. apiculatum* and *C. opuntiae* experience less herbivory from all of the mentioned insect predators [Miller2007].

Data

The data collected are from a long-term dataset spanning 2004 to 2023 taken from 30×30 meter plots at the Sevilleta LTER. The data initially included 134 naturally occurring plants across 4 spatial blocks censused annually from 2004 to 2008. Six of the plots were established in 2009 by tagging all existing plants within a 30×30 meter area. The final two plots were added to this census from 2011 onwards. Annually, in May we surveyed these plots, taking many types of demographic and partner data. For each cholla, we recorded plant survival from the last survey to the current survey. We recorded the height (cm), maximum crown width, and crown width perpendicular to the maximum, which are used to calculate plant volume (cm^3) based on the volume of a cone with the mean of maximum crown width and perpendicular crown width as the diameter. We recorded the total number of flower buds, including how many were aborted and how many were not. We recorded all ant species present and the number of ants we could count in 30 seconds. We also recorded the species of herbivore and the number present on the plant.

Statistical Modeling – Construction

With the data described above we fit a series of generalized linear mixed models (GLMMs) in
171 a hierarchical Bayesian framework with both fixed and random effects. Many of the vital rates
are estimated as a function of plant size, ant partner, or both. Ant partner type is included
as a predictor only where there are biological pathways through which ants could impact the
174 outcome of that process. The biological sources of variance (including individual, spatial, and
annual variance) are accounted for by including year-to-year and plot-to-plot random effects
in the models. Unless otherwise mentioned, all models use vague priors. The growth model
177 ($G_j(y, x)$) estimates the size of cholla, with fixed effects of the previous size and ant partner and
random effects of plot and year, using a Skew Normal distribution, with ω and α varying with
the previous size. Ants are included as a predictor here because ant partners defend plants
180 from herbivory, therefore decreasing the likelihood of segment loss. The survival model ($S_j(x)$)
estimates the probability of survival, with fixed effects of the previous size of the cholla and ant
partner and random effects of plot and year, using a Bernoulli distribution. Ants are included
183 as predictors here because ant partners defend cholla from herbivores and predators, decreasing
the likelihood of mortality due to either of these. The reproduction model ($P(x)$) estimates the
probability of reproducing each year, with fixed effects for the size and random effects of plot and
186 year, using a Bernoulli distribution. The total flowers model ($F(x)$) estimates the total flowers
produced by a plant, with fixed effects of size and random effects of plot and year, using a
Negative Binomial distribution. The viability model ($V_i(x)$) estimates the proportion of flowers
189 produced by a plant which are viable (not aborted), with fixed effects of the ant partner of the
cactus and random effects of plot and year, using a Binomial distribution. Ants are included
as predictors here because they defend the cacti from seed predation which can lead to floral
192 abortion. The ant transition rates model ($\tau_{i,j}(x)$) estimates the probability of a cactus being
visited by an ant partner, with fixed effects of the previous size of the cholla and the previous ant
partner and random effects of plot and year, using a Multinomial distribution. Ant partners are

195 included as predictors here because partners may choose to return to the same cholla repeatedly
 or choose new ones, therefore the previous partner may be a good indicator of the next partner.
 The recruit size model ($n_j(x)$) estimates the size distribution of all recruits from a given year,
 198 with no fixed or random effects, using a Normal distribution. With germination data from Miller
 et al., 2009, we fit two Bayesian generalized linear models for the probability of germinating
 from a seed in the first year (γ_1) or the second year ($1 - \gamma_1$), with no fixed or random effects,
 201 using a Binomial distribution. With data collected in a 2005-2006 recruit census, we fit a Bayesian
 generalized linear model for the probability of a seedling surviving to May (δ) (accounting for
 missed mortality events), with fixed effects of the previous size and random effects of the transect,
 204 using a Bernoulli distribution. With data from Miller 2007, we fit a Bayesian generalized linear
 model for the number of seeds produced by every flower on a cholla (κ) based on the ant partner,
 using a Negative Binomial distribution. Ant partners are included as predictors here because
 207 they reduce floral abortion rates and therefore may lead to higher numbers of seeds.

To obtain posterior estimates of the demographic parameters, we fit models using Markov
 chain Monte Carlo (MCMC) simulations via STAN run through version 4.0.2 of R. For each model,
 210 we obtained 3 chains of 10,000 iterations, each with randomly chosen initial conditions. The
 first 1,500 iterations were discarded as burn-in to eliminate transience associated with initial
 conditions. We did not thin the chains, thus all samples were retained. To assess the convergence
 213 of our models we assessed between and within chain convergence, the resulting figures are
 included in supplemental documents. To assess the overall model fit we carried out posterior
 predictive checks to examine how well the fitted model can generate simulated data similar to
 216 the real data. Large differences in the two indicate a poor model fit and can be assessed visually
 (figures included in supplemental documents). All estimated parameters are described in table
 1. Data and code for all vital rate models is included in the supplemental information.

Integral Projection Model Construction

Integral Projection Models describe population dynamics in discrete time, with continuous functions that relate vital rates to continuous and discrete state variables. We constructed a composite IPM which allows us to analyze the long term population growth rate of cholla with ant transition dynamics explicitly included. Many of the vital rate models are dependent not only on continuous size variables, but also on discrete variables representing the ant partner present. There is also a transition rate which determines what proportion of cholla are tended by each ant in a given year based on the previous ant partner and the size of the plants. This unique composite structure allows us to determine the individual effects of each ant species as well as the composite effects of several partners across the cholla population.

Following previous studies, we modeled the life cycle of cholla using continuously size-structured plants, $n_i(x)$, where x is the size of the cholla and i is the ant partner, and two discrete seed banks ($B_{1,t}$ and $B_{2,t}$) corresponding to 1 and 2-year old seeds. The dynamics of these 1 and 2-year old seedbanks are given by the following equations:

$$B_{1,t+1} = \kappa \delta \sum_i^4 \int_L^U P(x) V_i(x) F(x) n_i(x) dx$$

$$B_{2,t+1} = (1 - \gamma_1) B_{1,t}$$

The functions $P(x)$ and $F(x)$ give the probability of flowering, the number of flowerbuds produced based on the plant size x and the year t . The proportion of flowerbuds which will produce seeds ($V_i(x)$) is dependent on the plant size x and the ant species present on the plant i in year t . The integral is multiplied by the number of seeds per fruit (κ) and the probability of seed dispersal/survival (δ) to give the number of seeds that enter the one-year old seed bank. Parameters U and L are the upper and lower bounds, respectively, of the plant size distribution. Plants can recruit out of the one-year seed bank with the probability of γ_1 or transition to the two-year seed bank with a probability of $1 - \gamma_1$. Seeds in the two-year seed bank are assumed to either germinate with a probability of γ_2 or die.

The size dynamics of the cholla are given by:

$$n(y, i)_{t+1} = (\gamma_1 B_{1,t} + \gamma_2 B_{2,t}) \eta(y) \omega \beta_i + \sum_j^4 \int_L^U S_j(x) G_j(y, x) \tau_{ij}(x) n_j(x) dx$$

The first term gives the recruitment from one and two-year seed banks to a plant of size y , where $\eta(y)$ $N(* * *)$ gives the seedling size distribution and $\omega * **$ gives the proportion of seedlings which survive from germination (late summer) to the census (May). The second term reflects the changes in the population of the cholla which are not recruits, where $S_j(x)$ gives the probability of a plant of size x in year t surviving to year $t + 1$ with partner j . $G_j(y, x)$ gives the probability of growing from size x in year t to size y in year $t + 1$, respectively with partner j . Finally, τ_{ij} is the probability of a cholla which is size x with ant partner i in year t being tended by ant partner j in year $t + 1$.

Deterministic IPM Analysis

Analyzing an IPM requires discretizing the composite IPM into a matrix to calculate the dominant eigenvalue. Each component of the IPM (each set of ant partner specific vital rates) can be discretized into its own 200×200 matrix shown in figure ***. The possibility of eviction, when individuals are predicted to grow outside of the possible size classes, was avoided by adding probabilities of growing smaller or larger than the existing boundaries, as done by many others***. We used the composite IPM to quantify the effects of partner diversity on the intrinsic growth rate of cholla, $r \ln(\lambda)$. We calculated the r for each combination of ant partners: complete vacancy; *L. apiculatum* and vacancy; *C. opuntiae* and vacancy; other and vacancy; *L. apiculatum*, *C. opuntiae*, and vacancy, *L. apiculatum*, other, and vacancy; *C. opuntiae*, other, and vacancy; and all ant partners and vacancy. The relative distributions allowed us to determine if either complementarity or sampling effect were at play in the tree cholla-ant defense system. Sampling Effect requires that the r when all possible partners are present is equal to the r of the cholla

population when only the best partner is present. Complementarity requires that the r when all possible partners are present is greater than the r of any other combination of partners or any single partner.

Stochastic IPM Analysis

The third mechanism considered in this study requires annual variation to be explicitly considered in the IPM so we constructed a stochastic version of the IPM described above. The year of record was a random effect in all vital rate models meaning we were able to include an intercept for specific years. In order to include stochasticity in the analysis, we randomly sampled from the year-random effect intercepts 1,000 times. For each of these 1,000 iterations we calculated the r for every possible combination of ant partners. With these values we calculated δr between the scenario with all possible partners and the scenario with no partners. This δr measures the net benefits offered by partner diversity with annual variation. We also calculated the δr of the deterministic IPM to measure the net benefits offered by partner diversity without annual variation. These two relative differences allow us to determine if portfolio effect is at play. Portfolio effect is demonstrated when the net benefit with annual variation is larger than the net benefit without annual variation.

Acknowledgments

OEC would like to thank Madlen Wilmes, Gyuri Barabás, Flo Débarre, Vlastimil Křivan, and Greg Dwyer for their comments and suggestions on this template.

Statement of Authorship

OEC conceived the experiments, collected the data, and wrote the original draft. GHC provided specimens and analyzed the model. AQE oversaw data analysis and developed the code. All authors reviewed and edited the writing at all stages of revision.

Parameter	Median (95% CI)	Prior
growth xi intercept vacant ω_0^g	-5.210899 (-5.686865, -5.491787)	
growth xi intercept other ω_2^g	-5.8288 (-5.956217, 1.766021)	
growth xi intercept <i>C. opuntiae</i> ω_3^g	-4.529523 (-6.0770390, 0.1222112)	
growth xi intercept <i>L. apiculatum</i> ω_4^g	-5.106802 (-5.4499944, 0.5453901)	
growth xi size dependent vacant ω_1^g		
growth xi size dependent other ω_2^g		
growth xi size dependent <i>C. opuntiae</i> ω_3^g		
growth xi size dependent <i>L. apiculatum</i> ω_4^g		
growth omega intercept ω_0^g		
growth omega size dependent ω_1^g		
growth alpha intercept α_0^g		
growth alpha size dependent α_1^g		
1-year germination intercept α^{γ_1}		
2-year germination intercept α^{γ_2}		

Data and Code Availability

On initial submission, you may use this section to provide a URL for editors and reviewers that is ‘private for peer review’. After acceptance, this section must be updated with correct, working DOIs for data deposits (such as on the Dryad Digital Repository, ?) and code deposits (such as in Zenodo).

Appendix A: Additional Methods and Parameters

Fox–dog encounters through the ages

The quick red fox jumps over the lazy brown dog. The quick red fox has always jumped over the lazy brown dog. The quick red fox began jumping over the lazy brown dog in the 19th century and has never ceased from so jumping, as we shall see in figure ??. But there can be surprises (figure ??).

If the order and location of figures is not otherwise clear, feel free to include explanatory dummy text like this:

[Figure A1 goes here.]

[Figure A2 goes here.]

Further insights

Tables in the appendices can appear in the appendix text (see table ?? for an example), unlike appendix figure legends which should be grouped at the end of the document together with the other figure legends.

Lorem ipsum dolor sit amet, as we have seen in figures ?? and ??.

Table A1: Various rivers, cities, and animals

River	City	Animal
Chicago	Chicago	Raccoon
Des Plaines	Joliet	Coyote
Illinois	Peoria	Cardinal
Kankakee	Bourbonnais	White-tailed deer
Mississippi	Galena	Bald eagle

Note: See table ?? below for further table formatting hints.

References

- 312 Cook, O. E., G. H. Collaborator, and A. Q. Expert. 2015. Data from: Template and guidelines
for using \LaTeX in *The American Naturalist*. *American Naturalist*, Dryad Digital Repository,
<https://dx.doi.org/10.5061/dryad.XYZAB123>.
- 315 Darwin, C. 1859. On the origin of species by means of natural selection, or the preservation of
favoured races in the struggle for life. J. Murray, London.
- Davis, E. B., K. A. Brakora, and A. H. Lee. 2011. Evolution of ruminant headgear: a review.
318 *Proceedings of the Royal Society B* 278:2857–2865.
- Inglis, R. F., P. G. Roberts, A. Gardner, and A. Buckling. 2011. Spite and the scale of competition
in *Pseudomonas aeruginosa*. *American Naturalist* 178:276–285.
- 321 Fastovsky, D. E. 2009. Ideas in dinosaur paleontology: resonating to social and political context.
Pages 239–253 in D. Sepkoski and M. Ruse, eds. *The Paleobiological Revolution*. University
of Chicago Press, Chicago IL.
- 324 Xiao, X., D. J. McGlinn, and E. P. White. 2015. A strong test of the maximum entropy theory of
ecology. *American Naturalist* 185:E705–E80.

References Cited Only in the Online Enhancements

- ³²⁷ Tytler, W. 1759. *The Inquiry, Historical and Critical, into the Evidence against Mary Queen of Scots, and an Examination of the Histories of Dr. Robertson and David Hume with respect to that Evidence.* W. Creech, Edinburgh.

Tables

Table 1: Founders of *The American Naturalist*

Early editor	Years with the journal
Alpheus S. Packard Jr.	1867–1886
Frederick W. Putnam	1867–1874
Edward S. Morse	1867–1871
Alpheus Hyatt	1867–1871
Edward Drinker Cope ^a	1878–1897
J. S. Kingsley	1887–1896

Note: Table titles should be short. Further details should go in a ‘notes’ area after the tabular environment, like this.

^a Published the first description of *Dimetrodon*.

Figure legends

Figure 1: Figure legends can be longer than the titles of tables. However, they should not be excessively long—in most cases, they should be no more than 100 words each.

Figure 2: In this way, figure legends can be listed at the end of the document, with references that work, even though the graphic itself should be included for final files after acceptance. Instead, upload the relevant figure files separately to Editorial Manager; Editorial Manager should insert them at the end of the PDF automatically.

Figure A1: *A*, the quick red fox proceeding to jump 20 m straight into the air over not one, but several lazy dogs. *B*, the quick red fox landing gracefully despite the skepticism of naysayers.

Figure A2: The quicker the red fox jumps, the likelier it is to land near an okapi. For further details, see ?.

Video S1: Video legends can follow the same principles as figure legends. Counters should be set and reset so that videos and figures are enumerated separately.