

Thank you, next: partner turnover elevates benefits of mutualism for an ant-tended plant

Alexandra Campbell^{1,*}

Tom E.X. Miller¹

1. Program in Ecology and Evolutionary Biology, Department of BioSciences, Rice University, Houston, Texas 77005;

* Corresponding author; e-mail: amc49@rice.edu.

Manuscript elements:

Keywords:

Manuscript type: Article.

Prepared using the suggested L^AT_EX template for *Am. Nat.*

Abstract

Introduction

Mutualisms are species interactions where all participants benefit, leading to higher individual fitness and increased population growth rates. They are among the most widespread species interactions (10; 16; 22), but can deteriorate into commensalism or parasitism (4; 25; 34; 37; 40). Mutualisms are considered more context dependent than other species interactions (16; 22), meaning the magnitude and sign of interaction strength are often determined by environmental conditions and species' identities.

Mutualism is defined at the level of a species pair (+/+) but these interactions are embedded within multi-species communities, and growing evidence suggests that pairwise interactions are poor predictors of the net effects of multi-species mutualism (1; 33). A focal mutualist may interact with multiple guilds of partner types (e.g., plants that interact with pollinators, seed dispersers, soil microbes, and ant defenders) or with multiple partner species within the same guild (e.g., plants visited by multiple pollinator species). Within a mutualist guild, partner species often differ in the amount or type of goods or services they provide, making partner identity an important source of contingency in mutualism (39). Whether and how partner diversity modifies the demographic effects of mutualistic interactions remain open questions within relevance in applied settings (?).

There are multiple mechanisms by which partner diversity can influence the net benefits accrued by a focal mutualist – mirroring the mechanisms by which, at a larger scale of organization, biodiversity can influence ecosystem function (cite BEF chapter). First, when there is a hierarchy of fitness effects – a consistent ranking of best to worst mutualists – a more diverse sample of the partner community may be more likely to include the best partner (22). This can lead to an apparent benefit of diversity driven by a sampling effect (7). However, if partner associations are mutually exclusive then partner diversity may impose opportunity costs, leading to negative effects of a diverse mutualist assemblage relative to exclusive association with the single best partner (26). Second, even within a single mutualist guild, the benefits conferred by alternative

partner species can vary in type, and not just degree (12; 38; 39). This can lead to a positive effect of partner diversity through complementarity of alternative functions (7). Interference or synergies between partners can make their combined effect different than the expected from the sum of complementary functions (cite). Third, partner species can have species-specific responses to the environment, either spatially (32) or temporally (3). Multiple partners can therefore act as a 'portfolio' that stabilizes fitness benefits across spatial or temporal heterogeneity, leading to positive effects of partner diversity through the portfolio effect (7; 24).

Partner diversity can have different effects depending on whether partners are present all at once or sequentially (partner turnover) (13; 17; 28). Sequential associations are likely when alternative partners engage in interference competition for access to a shared mutualist (cite examples, including non-ant-plant examples). Turnover can happen at different timescales, from minutes to years (23; 31). The frequency of partner turnover can impact the level of benefits received by the focal mutualist, particularly if the benefits continue to accumulate (e.g., when sequential partners provide complementary functions) or if they saturate over time (36). Directionality of turnover can also influence diversity effects, particularly if partner identity changes consistently across ontogeny of a focal mutualist (19). For example, plant susceptibility to enemies can change across life stages (6; 9), so the benefits of defensive mutualism with ants are greatest when more defensive partner species align with more vulnerable life stages (17).

Defensive ant-plant mutualisms – where plants provide food and/or housing to ants that in turn defend them from enemies – are widespread interactions that offer valuable model systems for the ecology and evolution of mutualism (11; 12). Extrafloral nectar (EFN) bearing plants can serve as dietary resources that promote ant abundance and colony size (14; 28; 29). Presence of defensive ant partners is often linked to herbivory reduction (41?) and demographic advantages for the plant partner (Báez et al.). Defensive ant-plant mutualisms are commonly multi-species, where a guild of ant partner species share, and often compete for, a plant mutualist (2; 8; 11; 41). Ant partners can vary in their ability to deter herbivores (?), and visitation by low quality ant partners can prevent visitation by higher quality partners, consequently causing a reduction

in fitness through missed opportunity costs (20; 21). Another source of temporal variation is susceptibility to herbivory can also vary significantly throughout the life stages of the plant (9), suggesting that the order and timing of successive partners is important to the effectiveness of ant partners. Temporal dynamics of partner visitation therefore have important impacts on the fitness of the plant partners in these interactions (6; 9; 19). Recently many studies have investigated how partner diversity in these guilds has been shown to impact plant fitness (1; 5; 33; 42), however, none have considered the combined effects of partner identity, directional partner turnover, and temporal stochasticity on the fitness of plants with a longterm dataset.

This study examined the consequences of partner diversity in a food-for-protection mutualism between the tree cholla cactus (*Cylindriopuntia imbricata*), a long-lived EFN-bearing plant, and multiple species of ant partners. Previous studies have shown that herbivory by specialized insect herbivores negatively affects plant fitness (27), and ant defense reduces herbivore damage (26). Tree cholla are tended by two common ant species (*Liometopum apiculatum* and *Crematogaster opuntiae*) and several infrequent species, all of which are ground-nesting. These ant species locally co-occur at the scale of meters, but individual plants are typically tended by only one species that patrols the plant around-the-clock and maintains control of the plant's nectar resources for an entire growing season (30). Switches between partner species, or between vacancy and ant occupancy, commonly occur from one growing season to the next (26). Prior experiments suggested a hierarchy of mutualist quality, with *Liometopum apiculatum* providing strong fitness benefits and *Crematogaster opuntiae* having net negative fitness effects because herbivore deterrence is outweighed by deterrence of pollinators (26; 30). However, those studies did not integrate the demographic effects of ant defense across the plant life cycle, nor did they account for inter-annual fluctuations in the benefits provided by alternative partners, and therefore may have missed important mechanisms through which different partner species, and their combination, may be beneficial.

We used a unique long-term data set that allows us to explore mutualistic associations with multiple partner species and how the demographic effects of alternative partner species varied

across plant size structure and nearly 20 years of inter-annual fluctuations. We used this observational data set, contextualized by previous experiments, to ask whether and through which mechanism(s) partner diversity affects the fitness benefits of ant visitation for the focal plant partner. Specifically, we asked:

1. What are the demographic effects of association with alternative partners and how do these effects fluctuate across years?
2. What are the frequency and direction of partner turnover across the plant life cycle?
3. What is the net effect of partner diversity on plant fitness, and what mechanism(s) explain(s) this effect?

We used a hierarchical Bayesian statistical approach to estimate demographic vital rates for hosts in different states of ant occupancy, and to quantify state-dependent partner turnover. We then used a stochastic, multi-state integral projection model (IPM) that combines diverse effects on vital rates and pathways of partner turnover to quantify effects of partner diversity on plant fitness.

Methods

Study System

This study was conducted in the Los Piños mountains, a small mountain chain located on the Sevilleta National Wildlife Refuge, a Long-term Ecological Research site (SEV-LTER) in central New Mexico, USA. This is an area characterized by steep, rocky slopes, and perennial vegetation including grasses (*Bouteloua eriopoda* and *B. gracilis*), yuccas, cacti, and junipers. Tree cholla cacti are common in high Chihuahuan desert habitats, with their native range spanning the southwestern USA (?). These arborescent plants produce cylindrical segments with large spines. In the growing season (May to August in New Mexico), the plants initiate new vegetative segments and flower buds at the ends of existing segments. While most plants produce new segments

every season, only those which are reproductively mature produce flower buds. Like other EFN-bearing cacti, tree cholla secrete nectar from specialized glands on young vegetative segments and flower buds (28; 31). Flower buds produce more and higher-quality EFN than vegetative segments, making reproductive cholla valuable mutualist partners.

Tree cholla EFN is harvested by various ant species. At SEV-LTER, cholla are visited primarily by two species of formicoid, ground-nesting ants, *Crematogaster opuntiae* and *Liometopum apiculatum*, as well as other rarer species, including *Forelius pruinosus* and unidentified species of *Aphaenogaster*, and a *Camponotus*. *L. apiculatum* are the most frequent visitors with 25% – 60% of tree cholla tended by these ants, followed by *C. opuntiae* visiting between 5% – 20% of cacti (?) depending on the year. Between 30%–80% of cacti remain vacant in any given year. These ants rarely co-occur on a plant, likely due to interspecific competition (26): staged introductions of *C. opuntiae* to *L. apiculatum*-tended plants, and vice versa, provoke aggressive responses by resident ants (A. Campbell, *personal observation*). Each cholla is visited by a single ant species for the duration of a season, and the species of the visitors can change from one season to the next. At the beginning of the growing season, when EFN production begins, the ground-nesting ants will begin visiting tree cholla. They will visit the cholla every day during the season around the clock, with the most activity around sunrise or sunset (30). Smaller cholla are less likely to be visited because they produce very little EFN, so larger cholla, especially flowering individuals, are generally more highly tended (?). In late August, the tree cholla stop producing EFN and the ants vacate until the next growing season.

There are a variety of insect herbivores and seed predators which attack the cholla (?). An unidentified weevil of the genus *Gerstaekeria* feeds on vegetative and reproductive structures and implants their larvae within the plant tissue for the winter. A cactus bug, *Narnia pallidicornis*, (Hemiptera: Coreidae) feeds on all cholla parts with a preference for the reproductive structures (?). A seed predator, *Cahela ponderosella*, (Lepidoptera: Pyralidae) attacks developing fruits pre-dispersal and oviposits in open flowers mid-growing season where larvae burrow into the ripening ovary. These predators can have significant negative impacts on the fitness of individual

cholla and depress population growth (27). There is experimental evidence that tree cholla tended by *L. apiculatum* and *C. opuntiae* experience less herbivory than plants from which ants were excluded (26).

Data Collection

We used a longterm dataset spanning 2004 to 2023 taken from natural plants (2004 – 2008) and 30 × 30 meter plots (established in 2009) at SEV-LTER. The data initially included 134 naturally occurring plants across 4 spatial blocks censused annually from 2004 to 2008. Six of the plots were established in 2009 by tagging all existing plants within a 30 × 30 meter area. The final two plots were added to this census from 2011 onwards. Annually, in May we surveyed all individuals in these plots, taking demographic and partner data. For each cholla, we recorded plant survival from the last survey to the current survey. For surviving plants, we recorded the height (cm), maximum crown width (cm), and crown width perpendicular to the maximum (cm), which are used to calculate plant volume (cm^3) based on the elliptical volume of a cone with the mean of maximum crown width and perpendicular crown width as the diameter. We recorded the total number of flower buds, including how many were aborted and how many were not. We recorded all ant species present and the number of ants we could count in 30 seconds. In the rare case that more than one species of ant was present, we recorded the ant state as the more prevalent of the species, and recorded the identity and number of the secondary species present in the comments. We also recorded the numbers and species of herbivores on each plant. We conducted recruit and new plant surveys by splitting up each plot and walking the entire area to search for untagged plants. Plants less than 1 cm tall were marked as recruits and larger plants as new plants, these were all tagged, GPS located, and recorded. red¹

¹redAlso need to describe recruit data, seed banks, seed counts per fruit.

Statistical Modeling

156 With the data described above we fit a series of generalized linear mixed models (GLMMs) in
a hierarchical Bayesian framework to serve as vital rate sub-models of the IPM. Many of the
vital rates are estimated as a function of plant size, ant partner, or both. Ant partner type is
159 included as a predictor in the growth model, survival model, and floral viability model, where
there are biological pathways through which ants could impact the outcome of that process. The
ant partners defend the cacti from herbivores, meaning their presence can lead to higher survival
162 rates and less vegetative structure loss, and seed predators, meaning their presence can lead to
fewer attacked and aborted flowers. The biological sources of variance (including spatial, and
annual variance) are accounted for by including year-to-year and plot-to-plot random effects in
165 the models. Unless otherwise mentioned, all models use vague priors.

The growth model $\text{pink}(G(x', x, a))$ estimates the size of cholla y^{G^2} in year $t + 1$, with fixed effects of the previous size x and ant partner i in year t and random effects of plot w and year u , using a Skew Normal distribution, with ω and α varying with the previous size. A skew normal distribution was chosen here because the data had evidence of skew and previous efforts using a gaussian distribution poorly predicted the sizes of these plants, overestimating the next sizes for large plants. ω denotes the scale of the data in a Skew Normal distribution while α denotes the slant, when α is 0, the ω is equivalent to the sd of a Normal distribution. The ω and α estimates vary across size because the scale and skew of the data varied across size.

$$y^G \text{ SkewN}(\hat{G}, \hat{\omega}, \hat{\alpha})$$

$$\hat{G} = \beta_0^G + \beta_1^G \times x + \beta_2^G \times a + \beta_4^G \times x \times a + u + w$$

$$\hat{\omega} = \beta_0^\omega + \beta_1^\omega \times x$$

$$\hat{\alpha} = \beta_0^\omega + \beta_1^\omega \times x + \beta_2^\omega \times x^2$$

²*pink* This is where I get a little confused between the x and x' sizes and the ys . Should this all be noted like this?

where $u \sim N(0, \sigma_{yr}^2)$ is the year random effect with year specific variance of σ_{yr} in year t , and $w \sim N(0, \sigma_{plot}^2)$ is the plot random effect with plot specific variance of σ_{plot} . Ants are included
 168 as a predictor here because ant partners defend plants from herbivory, therefore decreasing the likelihood of segment loss.

The survival model ($S(a, x)$) estimates the probability of survival y^S from year t to year $t + 1$, with fixed effects of the previous size of the cholla x and ant partner i in year t and random effects of plot w and year u , using a Bernoulli distribution:

$$y^S \sim \text{Bern}(\hat{S})$$

$$\text{logit}(\hat{S} = \beta_0^S) + \beta_1^S \times x + \beta_2^S \times a + \beta_3^S \times x \times a + u + w$$

where $u \sim N(0, \sigma_{yr}^2)$ is the year random effect with year specific variance of σ_{yr} in year t , and
 171 $w \sim N(0, \sigma_{plot}^2)$ is the plot random effect with plot specific variance of σ_{plot} . A Bernoulli distribution was chosen here because the only two possible outcomes are survival or death. Ants are included as predictors here because ant partners defend cholla from herbivores and predators, decreasing
 174 the likelihood of mortality due to either of these.

The reproduction model ($P(x')$) estimates the probability of reproducing y^P in year $t + 1$, with fixed effects for the size x' in year $t + 1$ and random effects of plot w and year u , using a Bernoulli distribution:

$$y^P \sim \text{Bern}(\hat{P})$$

$$\text{logit}(\hat{P}) = \beta_0^P + \beta_1^P \times x' + u + w$$

where $u \sim N(0, \sigma_{yr}^2)$ is the year random effect with year specific variance of σ_{yr} in year t , and $w \sim N(0, \sigma_{plot}^2)$ is the plot random effect with plot specific variance of σ_{plot} .

The total flowers model ($F(x)$) estimates the total flowers produced by a plant y^F in year $t + 1$, with fixed effects of size x in year t and random effects of plot w and year u , using a Negative Binomial distribution:

$$y^F \sim \text{NegativeBinom}(\hat{F}, \hat{\phi})$$

$$\log(\hat{F}) = \beta_0^F + \beta_1^F \times x + u + w$$

$$\log(\hat{\phi}) = \beta_0^\phi$$

177 where $u \sim N(0, \sigma_{yr}^2)$ is the year random effect with year specific variance of σ_{yr} in year t , and
 $w \sim N(0, \sigma_{plot}^2)$ is the plot random effect with plot specific variance of σ_{plot} .

The viability model ($V_i(x)$) estimates the proportion of flowers produced by a plant which are viable (not aborted) y^V in year $t + 1$, with fixed effects of the ant partner of the cactus i in year t and random effects of plot w and year u , using a Binomial distribution:

$$y^V \sim \text{Binom}(\hat{V})$$

$$\text{logit}(\hat{V}) = \beta_0^V \times a + u + w$$

180 where $u \sim N(0, \sigma_{yr}^2)$ is the year random effect with year specific variance of σ_{yr} in year t , and
 $w \sim N(0, \sigma_{plot}^2)$ is the plot random effect with plot specific variance of σ_{plot} . Ants are included as predictors here because they defend the cacti from seed predation which can lead to floral abortion.

The ant transition rates model ($\tau(x, a, a')$) estimates the probability of a cactus being visited by an ant partner a' y^τ , with fixed effects of the previous size of the cholla x and the previous ant partner a in year t and random effects of plot w and year u , using a Multinomial distribution:

$$y^\tau \sim \text{Multinomial}(\hat{\tau})$$

$$\text{logit}(\tau) = \beta_0^\tau + \beta_1^\tau \times x + \beta_2^\tau \times a + \beta_3^\tau \times x \times a + u + w$$

183 where $u \sim N(0, \sigma_{yr}^2)$ is the year random effect with year specific variance of σ_{yr} in year t , and
 $w \sim N(0, \sigma_{plot}^2)$ is the plot random effect with plot specific variance of σ_{plot} . Ant partners are included as predictors here because partners may choose to return to the same cholla repeatedly
186 or choose new ones, therefore the previous partner may be a good indicator of the next partner.

The recruit size model ($n(x, a')$) estimates the size distribution of all recruits y^n from a given year $t + 1$, with no fixed or random effects, using a Normal distribution:

$$y^n \sim N(\hat{\eta}, \hat{\sigma})$$

$$e\hat{t}a = \beta_0^\eta$$

where $\hat{\sigma}$ is estimated with a non-informative prior.

With germination data from Miller et al., 2009****, we fit two Bayesian generalized linear models for the probability of germinating from a seed in the first year (γ_1) or the second year ($1 - \gamma_1$) in year $t + 1$, with no fixed or random effects, using a Binomial distribution:

$$y^{\gamma_1} \text{ Binomial}(\hat{\gamma}_1)$$

$$y^{\gamma_2} \text{ Binomial}(\hat{\gamma}_2)$$

$$\text{logit}(\hat{\gamma}_1) = \beta_0^{\gamma_1}$$

$$\text{logit}(\hat{\gamma}_2) = \beta_0^{\gamma_2}$$

With data collected in a 2005-2006 recruit census, we fit a Bayesian generalized linear model for the probability of a seedling surviving to May (δ) of year $t + 1$ (accounting for missed mortality events), with fixed effects of the previous size x and random effects of the transect m , using a Bernoulli distribution:

$$y^\delta \text{ Bern}(\hat{\delta})$$

$$\text{logit}(\hat{\delta}) = \beta_0^\delta + m$$

where $m \sim N(0, \sigma_{\text{transect}}^2)$ is the random effect of transect where the recruited individual was analyzed for survival.

With data from Miller 2007, we fit a Bayesian generalized linear model for the number of seeds produced by every flower on a cholla (κ) in year $t + 1$ based on the ant partner i in year $t + 1$, using a Negative Binomial distribution:

$$y^\kappa \text{ NegativeBinomial}(\hat{L}, \hat{\phi})$$

$$\hat{L} = \beta_0^L \times j$$

$$\hat{\phi} = \beta_0^\phi$$

Ant partners are included as predictors here because they reduce floral abortion rates and therefore may lead to higher numbers of seeds.

To obtain posterior estimates of the demographic parameters, we fit models using Markov chain Monte Carlo (MCMC) simulations via STAN run through version 4.0.2 of R ³ For each model, we obtained 3 chains of 10,000 iterations, each with randomly chosen initial conditions. The first 1,500 iterations were discarded as burn-in to eliminate transience associated with initial conditions. We did not thin the chains, thus all samples were retained. To assess the convergence of our models we assessed between and within chain convergence, the resulting figures are included in supplemental documents. To assess the overall model fit we carried out posterior predictive checks to examine how well the fitted model can generate simulated data similar to the real data. Large differences in the two indicate a poor model fit and can be assessed visually (figures included in supplemental documents). All estimated parameters are available in the data. Data and code for all vital rate models is included in the supplemental information.

Deterministic Integral Projection Model Construction and Analysis

Integral Projection Models describe population dynamics in discrete time, with functions that relate vital rates to continuous state variables. While IPMs are a natural choice for populations with continuous size structure, they can also be modified to accommodate a combination of continuous and discrete state variables, as we do here. We constructed a deterministic composite IPM which allows us to analyze the long term population growth rate of cholla with ant transition dynamics explicitly included. This novel structure allows us to determine the individual effects of each ant species and the composite effects of several partners across the cholla population.

Following previous studies of this population, we modeled the life cycle of cholla using continuously size-structured plants, $n(x', a', t + 1)$, where x' is the size of the cholla in year $t + 1$ and a' is the ant partner in year $t + 1$, and two discrete seed banks ($B_1(t + 1)$ and $B_2(t + 1)$) corresponding to 1 and 2-year old seeds. The dynamics of these 1 and 2-year old seedbanks are

³*red*You need to cite R and RStan

given by the following equations:

$$\begin{aligned}
B_1(t+1) &= \delta \sum_a^A \int_L^U \kappa(a) P(x') V(a) F(x) n(x, a, t) dx \\
B_2(t+1) &= (1 - \gamma_1) B_1(t)
\end{aligned}$$

The functions $P(x')$ and $F(x)$ give the probability of flowering, the number of flowerbuds produced based on the plant size x' in year $t+1$ and x in year t respectively. The proportion of flowerbuds which will produce seeds ($V(a)$) is dependent on the ant species present on the plant i in year t . The function for the number of seeds per fruit ($\kappa(a)$) is dependent on the ant partner present a in year t . These functions are integrated across size, from the smallest observed size L to the largest observed size U .⁴ The integral is multiplied by the probability of seed dispersal/survival (δ) to give the number of seeds that enter the one-year old seed bank. Parameters U and L are the upper and lower bounds, respectively, of the plant size distribution. Plants can recruit out of the one-year seed bank with the probability of γ_1 or transition to the two-year seed bank with a probability of $1 - \gamma_1$. Seeds in the two-year seed bank are assumed to either germinate with a probability of γ_2 or die.

The size and ant dynamics of the cholla are given by:

$$\begin{aligned}
n(x', a', t+1) &= (\gamma_1 B_1(t) + \gamma_2 B_2(t)) \eta(x') \omega \rho_{a'} + \\
&\sum_{a'}^A \int_L^U S(x, a) G(x', x, a) \tau(x, a, a') n(x, a, t) dx
\end{aligned}$$

$n(x', a', t+1)$ explains the dynamics of all plants of size x' with ant partner x' at time $t+1$. The first term in the equation shows how many of these plants come from the seedbank, either year one or year two seed banks, while the second term describes how many of these plants come from previously existing plants. The larger size x' is the less likely these plants were recently in the seedbanks. The number of recruits generated from year-one seedbanks is given by the number of seeds in the seedbank $B_1(t)$ multiplied by the year-one seed bank germination rate

⁴pinkI know that technically L is smaller than the smallest size and U is larger, but I am not sure if here is the place to put it.

γ_1 . The number of recruits generated from year-two seedbanks is calculated in a similar way with year-two specific estimates. The first term estimates the number of individuals recruited from a one or two-year seed bank to a plant of size x' as the number of individuals germinated from a first year seed bank $\gamma_1 B_1(t)$ and a second year seed bank $\gamma_2 B_2(t)$ multiplied by the recruit size distribution term $\eta(x')$, the proportion of seedlings which survive from germination (late summer) to the census (May) ω . Finally, this term is multiplied by $\rho_{a'}$, which is an ant specific term which is 1 when a' indicates vacancy and 0 when a' indicates any ant presence, to ensure all new recruits are vacant as observed in nature. Plants not coming from the seedbank to size x' and ant state a' were previously existing plants $n(x, a, t)$ of size x and ant state a . The number of these plants can be described by the proportion of plants of size x and ant state a in year t ($n(x, a, t)$) which survive to year $t + 1$ ($S(x, a)$), from size x to size x' ($G(x', x, a)$), and transition from ant partner a to a' ($\tau(x, a, a')$).

Analyzing an IPM requires discretizing the composite IPM into a matrix to calculate the dominant eigenvalue. In a traditional IPM x is discretized into b bins, replacing the continuous kernel into a $b \times b$ matrix, in this case there is additional complexity in the form of transitions between ant partners. Each combination of previous ant state a and next ant state a' represents a unique set of plants in the population and therefore must be discretized individually, leading to a matrix size of $mb \times mb$ where m is the number of unique combinations of a and a' (how many possible ant transitions there are). In this model we have two additional discrete states (year one and year two seed banks) leading to a final matrix size of $m(b + 2) \times m(b + 2)$. We used $b = 200$ bins. We extend the integration limits L and U to avoid unintentional eviction (?). We estimate the asymptotic population growth rate λ as the dominant eigenvalue of the discrete kernel for each combination of ant partners: complete vacancy; *C. opuntiae* and vacancy; other and vacancy; *L. apiculatum*, *C. opuntiae*, and vacancy, *L. apiculatum*, other, and vacancy; *C. opuntiae*, other, and vacancy; and all ant partners and vacancy. Rather than estimating a single λ value for each possible population, we used 1,000 random parameter estimations from iterations of our bayesian statistical models to estimate 1,000 λ values as a distribution. We compare the distributions of

λ across each combination of ant partners to whether sampling effect or complementarity is at play in the system. To compare the distributions of two hypothetical populations, A and B, we subtract the vector of 1,000 λ_A estimations of one population from the 1,000 λ_B estimations of the other ($\lambda_A - \lambda_B$). If the average of these differences is positive, population A has a higher λ than B, if the average is 0, they are equal, and if the average is negative, population B has a higher λ than A.

To determine if sampling effect is at play in a system, we must first determine if there is a "best" partner, by determining which single ant association is correlated with the highest λ estimation. We can do this by comparing the distributions of each λ and finding the one which is larger than all others. Then we must show that the λ of a population with all possible partners is equal to that of λ of a population with only the "best" partner, by comparing the distributions as described earlier. If it is significantly larger then complementarity is potentially at play (more on this in the next paragraph). If it is significantly smaller, this indicates that rather than positive effects of partner diversity, there are actually important costs of partner diversity that dampen the population growth rate.

To determine if complementarity is at play in a system, we must determine if the partner scenario which leads to the highest λ is the most partner diverse scenario. We can do this by comparing the distribution of λ of a population with all possible partners to the distribution of λ of all other populations. If the λ of a population with all possible partners is the largest, it indicates that complementarity is at play. If neither the conditions for sampling effect or complementarity, it indicates that rather than positive effects of partner diversity, there are actually costs of diversity.

Stochastic Integral Projection Model Construction and Analysis

The third mechanism considered in this study requires annual variation to be explicitly considered in the IPM so we constructed a stochastic version of the IPM described above. This stochastic IPM draws from the same statistical models as the deterministic version, and is described below:

$$\begin{aligned}
291 \quad B_1(t+1) &= \delta \sum_a^A \int_L^U \kappa(a) P(x', u, w) V(a, u, w) F(x, u, w) n(x, a, t) dx \\
B_2(t+1) &= (1 - \gamma_1) B_1(t) \\
n(x', a', t+1) &= (\gamma_1 B_1(t) + \gamma_2 B_2(t)) \eta(x') \omega \rho_{a'} + \\
294 \quad \sum_a^A \int_L^U S(x, a, u, w) G(x', x, a, u, w) \tau(x, a, a', u, w) n(x, a, t) dx
\end{aligned}$$

This system of equations still describes the contribution of seedbanks and existing plants to the group of plants of size x' and ant partner a' in year $t+1$, but now includes the random effects of plot u and year w . These effects are included in the form of linear scales on each statistical model where they were biologically relevant. Year effects w are both year and ant specific, meaning for each year there are different ant and year-specific linear scales for each year of data. When calculating stochastic λ , every one of the 1,000 parameter-iteration associated λ s actually come from the mean of 1,000 year-specific estimations. For each parameter estimate taken from the bayesian models we calculate 1,000 estimations of λ , each associated with a year random effect which was randomly selected from the *****18 years of data. The mean of these estimations become the parameter-iteration specific λ estimation. This is repeated 1,000 times for each iteration of parameters to get the stochastic posterior distribution of λ . We compare the stochastic distributions to the deterministic ones to determine if portfolio effect is at play in this system.

To determine if portfolio effect is at play we must show that partner diversity buffers the population from environmental variation (in our case annual variation assessed by the inclusion of year random effects). We can estimate the effects of partner diversity on the focal mutualist by calculating the difference in λ distributions between populations with no partners and populations with all partners present. Then we compare the difference in the effects of partner diversity on the focal mutualist with and without stochasticity. If the stochastic difference in the effects of partner diversity is greater than the deterministic difference, this indicates that partner diversity is more beneficial under varying environments and therefore portfolio effect is at play.

Results

Statistical Modeling

318 Plants tended by other ants experienced positive growth rates across all sizes, with the highest
chances of shrinking occurring at the largest sizes (?)a. They experienced significantly higher
growth than vacant plants ($p = 2.2 \times 10^{-16}$), and similar mean growth to *L. apiculatum* tended
321 plants ($p = 0.9135$) and *C. opuntiae* tended plants ($p = 0.4864$). Plants tended by *C. opuntiae* ants
experienced positive growth rates across all sizes, with the highest chances of shrinking occurring
at the largest sizes (?)b. They experienced significantly higher mean growth than vacant plants
324 ($p = 4.88 \times 10^{-14}$). Plants tended by *L. apiculatum* experienced positive growth rates across all
sizes, with the highest chances of shrinking occurring at the largest sizes (?)c. Vacant plants
experienced the lowest observed growth, except at the largest sizes where they are less likely to
327 shrink than tended plants (?)d.

Plants tended by *C. opuntiae* ants experienced significantly higher mean survival rates than
L. apiculatum tended plants ($p = 2.908 \times 10^{-14}$), plants tended by other ants ($p = 4.488 \times 10^{-14}$),
330 and vacant plants ($p = 2.2 \times 10^{-16}$) (?). These *C. opuntiae* plants have the highest mean survival
at small sizes before reaching a mean survival rate of about 100% when medium to large. Plants
tended by *L. apiculatum* experienced similar mean survival rates to plants tended by other ants
333 and vacant plants. *L. apiculatum* tended plants and other tended plants reached a mean survival
rate of nearly 100% by medium sizes, with other tended plants reaching this rate slower. Vacant
plants never reach mean survival rates of 100%.

336 There is a clear size effect on the number of flowers produced. The mean number of flowers
produced by a plant remains at 0 until the plant reaches medium sizes after which the mean
number of flower produced increases exponentially to about 40 at large sizes.

339 Plants tended by *L. apiculatum* ants experienced the highest viability rates, significantly higher
than *C. opuntiae* tended plants ($p = 2.2 \times 10^{-16}$), other tended plants ($p = 2.2 \times 10^{-16}$), and

vacant plants ($p = 2.2 \times 10^{-16}$). The mean viability rate of *L. apiculatum* tended plants was
342 84.888%. Plants tended by *C. opuntiae* ants experienced significantly higher viability rates than
vacant plants ($p = 2.2 \times 10^{-16}$), but not than other tended plants ($p = 0.1168$). The mean viability
rate of *C. opuntiae* plants was 75.934%. Plants tended by other ants also experienced significantly
345 higher viability rates than vacant plants ($p = 2.2 \times 10^{-16}$). The mean viability rate of other tended
plants was 76.367%. The mean viability rate of vacant plants was 70.442%.

The probability of a plant reproducing in a given year is highly size dependent. The mean
348 probability of reproducing remains at about 0% until the plant reaches a medium size, after
which the mean probability of reproducing increases steadily before reaching about 100% at
large sizes.

351 The mean number of seeds produced by *C. opuntiae* tended plants is significantly lower than
the mean number of seeds produced by *L. apiculatum* plants ($p = 2.2 \times 10^{-16}$) and vacant plants
($p = 2.2 \times 10^{-16}$). *C. opuntiae* tended plants produce a mean of 2.561 seeds per flower. The
354 mean number of seeds produced by *L. apiculatum* tended plants is slightly higher than the mean
number produced by vacant plants ($p = 0.02521$). *L. apiculatum* tended plants produce a mean of
3.563 seeds per flower. Vacant plants produce a mean of 3.435 seeds per flower.

357 The mean pre-census seed survival rate is 16.99% with 90% of model iterations landing be-
tween 1.13% and 40.72%.

Seeds have a significantly higher probability of germinating in year one than in year two
360 ($p = 2.2 \times 10^{-16}$). Seeds in year one have a mean germination rate of 0.578% and seeds in year
two have a mean germination rate of 0.221%.

The mean size of new recruits is $-1.88 \log(\text{cm}^3)$ with 90% of new recruits falling between the
363 sizes of $-2.12 \log(\text{cm}^3)$ and $-1.64 \log(\text{cm}^3)$.

Plants that were previously vacant will most likely remain vacant until large. These plants
have significantly higher probabilities of being vacant rather than becoming tended by *C. opuntiae*,
366 *L. apiculatum*, or other ants (respectively, $p = 2.2 \times 10^{-16}$, $p = 2.2 \times 10^{-16}$, and $p = 2.2 \times 10^{-16}$).
The probability of previously vacant plants becoming tended by *L. apiculatum* ants is significantly

higher than the probabilities of becoming tended by *C. opuntiae* or other ants (respectively, $p = 3.83 \times 10^{-14}$, $p = 2.2 \times 10^{-16}$). The probability of previously vacant plants becoming tended by other ants is significantly lower than the probability of becoming tended by *C. opuntiae* ($p = 0.00227$).

Plants that were previously tended by *C. opuntiae* will most likely remain vacant until large. These plants have significantly higher probabilities of becoming vacant rather than remaining tended by *C. opuntiae* or becoming tended by *L. apiculatum* or other ants (respectively, $p = 2.2 \times 10^{-16}$, $p = 2.2 \times 10^{-16}$, $p = 2.2 \times 10^{-16}$). The probability of these plants remaining tended by *C. opuntiae* ants is significantly higher than the probability of becoming tended by *L. apiculatum* or other ants (respectively, $p = 2.2 \times 10^{-16}$, $p = 3.797 \times 10^{-9}$). The probability of these plants becoming tended by *L. apiculatum* is significantly higher than the probability of becoming tended by other ants ($p = 3.797 \times 10^{-9}$).

Plants that were previously tended by *L. apiculatum* ants will most likely remain vacant until large. These plants have significantly higher probabilities of becoming vacant than remaining tended by *L. apiculatum* or becoming tended by *C. opuntiae* or other ants (respectively, $p = 2.2 \times 10^{-16}$, $p = 2.2 \times 10^{-16}$, and $p = 2.2 \times 10^{-16}$). These plants have significantly higher probabilities of remaining tended by *L. apiculatum* ants than becoming tended by *C. opuntiae* or other ants (respectively, $p = 2.2 \times 10^{-16}$, $p = 2.2 \times 10^{-16}$). The probability of these plants becoming tended by *C. opuntiae* ants is significantly higher than the probability of them becoming tended by other ants ($p = 0.007925$).

Plants that were previously tended by other ants will most likely remain vacant until they are large. These plants have significantly higher probabilities of becoming vacant rather than remaining tended by other ants or becoming tended by *C. opuntiae* or *L. apiculatum* ants (respectively, $p = 2.2 \times 10^{-16}$, $p = 2.2 \times 10^{-16}$, $p = 2.2 \times 10^{-16}$). The probability of these plants becoming tended by *L. apiculatum* ants is significantly higher than the probability of remaining tended by other ants or becoming tended by *C. opuntiae* (respectively, $p = 6.309 \times 10^{-12}$, $p = 6.256 \times 10^{-13}$). The probability of these plants remaining tended by other ants is not higher than the probability

of becoming tended by *C. opuntiae* ants ($p = 0.954$).

In all plants, the most likely scenario is vacancy. In most cases, the second most likely next ant partner is *L. apiculatum* with the exception of plants which were previously tended by *C. opuntiae* who are second most likely to retain *C. opuntiae* as their partners in successive years. The only case where two next partners do not have statistically significantly different probabilities is when previously other tended plants either remain tended by other ants or become tended by *C. opuntiae*.

Deterministic IPM

Stochastic IPM

Discussion

Acknowledgments

Data and Code Availability

Appendix A: Additional Methods and Parameters

Literature Cited

[1] Afkhami (2014). Multiple mutualist effects: conflict and synergy in multispecies mutualisms. *Ecology*, 95(4):833–844.

[2] Agrawal, A. A. and Rutter, M. T. (1998). Dynamic anti-herbivore defense in ant-plants: The role of induced responses.

[3] Alarcón, R., Waser, N. M., and Ollerton, J. (2008). Year-to-year variation in the topology of a plant-pollinator interaction network. *Oikos*, 117(12):1796–1807.

[4] Bahia, R., Lambertucci, S. A., Plaza, P. I., and Speziale, K. L. (2022). Antagonistic-mutualistic interaction between parrots and plants in the context of global change: Biological introductions and novel ecosystems. *Biological Conservation*, 265(November 2021):109399.

[5] Barrett, L. G., Bever, J. D., Bissett, A., and Thrall, P. H. (2015). Partner diversity and identity impacts on plant productivity in acacia-rhizobial interactions. *Journal of Ecology*, 103:130–142.

[6] Barton, K. E. and Koricheva, J. (2010). The Ontogeny of Plant Defense and Herbivory: Characterizing General Patterns Using Meta-Analysis. *The American Naturalist*, 175(4):481–493.

[7] Batstone, R. T. (2018). Using niche breadth theory to explain generalization in mutualisms. *Ecology*, 99(5):1039–1050.

[8] Beattie, A. (1985). *The Evolutionary Ecology of Ant-Plant Mutualisms*.

[9] Boege, K. and Marquis, R. J. (2005). Facing Herbivory as You Grow up: The Ontogeny of Resistance in Plants. *Trends in Ecology and Evolution*, 20:441–448.

[10] Bronstein, J. L. (1994). Conditional Outcomes in Mutualistic Interactions. *TREE*, 9(6):214–217.

[11] Bronstein, J. L. (1998). The Contribution of Ant-Plant Protection Studies to Our Understanding of Mutualism. *bioTropica*, 30(2):150–161.

[12] Bronstein, J. L., Bronstein, J. L., Alarcón, R., and Geber, M. (2006). The evolution of plant – insect mutualisms. pages 412–428.

[13] Bruna, E. M., Izzo, T., Inouye, B., and Vasconcelos, H. (2014). Effect of mutualist partner identity on plant demography. *Ecology*, 95(12):3237–3243.

[14] Byk, J. and Del-Claro, K. (2011). Ant-plant interaction in the neotropical savanna: Direct beneficial effects of extrafloral nectar on ant colony fitness. *Population Ecology*, 53:327–332.

[Báez et al.] Báez, S., Donoso, D. A., Queenborough, S. A., Jaramillo, L., Valencia, R., and Dangles, O. Ant mutualism increases long-term growth and survival of a common amazonian tree.

[16] Chamberlain, S. A., Bronstein, J. L., and Rudgers, J. A. (2014). How context dependent are species interactions? *Ecology Letters*, 17(7):881–890.

[17] Djiéto-Lordon, C., Dejean, A., Ring, R. A., Nkongmeneck, B. A., Lauga, J., and McKey, D. (2005). Ecology of an improbable association: The pseudomyrmecine plant-ant *Tetraponera tessmanni* and the myrmecophytic liana *Vitex thyrsoflora* (Lamiaceae) in Cameroon. *Biotropica*, 37(3):421–430.

[18] Donald, M. L. and Miller, T. E. (2022). Does ant–plant mutualism have spillover effects on the non-partner ant community? *Ecology and Evolution*, 12(1):e8524.

[19] Fonseca, C. R., Benson, W. W., and Zoologia, D. D. (2003). Nordic Society Oikos Ontogenetic Succession in Amazonian Ant Trees Author (s): Carlos Roberto Fonseca and Woodruff Whitman Benson Published by : Wiley on behalf of Nordic Society Oikos Stable URL : <http://www.jstor.org/stable/3548044> JSTOR is a not-for-. *Oikos*, 102(2):407–412.

[20] Fraser, A. M., Axen, A. H., and Pierce, N. E. (2001). Assessing the quality of different ant species as partners of a myrmecophilous butterfly. *Oecologia*, 129:452–460.

[21] Frederickson, M. E. (2005). Ant species confer different partner benefits on two neotropical myrmecophytes. *Oecologia*, 143:387–395.

[22] Frederickson, M. E. (2013). Rethinking Mutualism Stability: Cheaters and the Evolution of Sanctions. *Quarterly Review of Biology*, 88(4):269–295.

[23] Horvitz, C. and Schemske, D. (1986). Seed Dispersal of a Neotropical Myrmecochore : Variation in Removal Rates and Dispersal Distance Author (s): Carol C . Horvitz and Douglas

W . Schemske Published by : Association for Tropical Biology and Conservation Stable URL :
http://www.jstor.org/st. *bioTropica*, 18(4):319–323.

- 462 [24] Lázaro, A., Gómez-Martínez, C., González-Estévez, M. A., and Hidalgo, M. (2022). Portfolio
effect and asynchrony as drivers of stability in plant–pollinator communities along a gradient
of landscape heterogeneity. *Ecography*, 2022(3):1–14.
- 465 [25] Mandyam, K. G. and Jumpponen, A. (2014). Mutualism-parasitism paradigm synthesized
from results of root-endophyte models. *Frontiers in Microbiology*, 5(DEC):1–13.
- [26] Miller, T. E. (2007). Does Having Multiple Partners Weaken the Benefits of Faculative Mutu-
468 alism? A Test with Cacti and Cactus-Tending Ants. *Oikos*, 116(3):500–512.
- [27] Miller, T. E., Louda, S. M., Rose, K. A., and Eckberg, J. O. (2009). Impacts of insect herbivory
on cactus population dynamics: Experimental demography across an environmental gradient.
471 *Ecological Monographs*, 79(1):155–172.
- [28] Ness, J. H., Morris, W., and Bronstein, J. L. (2006). INTEGRATING QUALITY AND QUAN-
TITY OF MUTUALISTIC SERVICE TO CONTRAST ANT SPECIES PROTECTING FEROCAC-
474 TUS WISLIZENI. *Ecology*, 87(4):912–921.
- [29] Ness, J. H., Morris, W. F., and Bronstein, J. L. (2009). For ant-protected plants, the best
defense is a hungry offense. *Ecology*, 90:2823–2831.
- 477 [30] Ohm, J. R. and Miller, T. E. (2014). Balancing Anti-Herbivore Benefits and Anti-Pollinator
Costs of Defensive Mutualists. *Ecology*, 95(10):2924–2935.
- [31] Oliveira, P. S., Rico-Gray, V., Díaz-Castelazo, C., and Castillo-Guevara, C. (1999). Interaction
480 between ants, extrafloral nectaries and insect herbivores in Neotropical coastal sand dunes:
Herbivore deterrence by visiting ants increases fruit set in *Opuntia stricta* (Cactaceae). *Func-
tional Ecology*, 13(5):623–631.

- 483 [32] Ollerton, J., Johnson, S. D., and Hingston, B. A. (2006). Geographical Variation In Diversity and Specificity of Pollination Systems — Jeff Ollerton - Academia.edu. In *Plant–pollinator interactions: from specialization to generalization.*, number January 2006, pages 282–308.
- 486 [33] Palmer, T. M., Doak, D. F., Stanton, M. L., Bronstein, J. L., Kiers, T. E., Young, T. P., Goheen, J. R., and Pringle, R. M. (2010). Synergy of multiple partners, including freeloaders, increases host fitness in a multispecies mutualism. *PNAS*, 107(40):17234–17239.
- 489 [34] Rodriguez-Rodriguez, M. C., Pedro, J., and Valido, A. (2017). Functional consequences of plant-animal interactions along the mutualism-antagonism gradient. *Ecology*, 98(5):1266–1276.
- [35] Rogers, S. R., Tarpy, D. R., and Burrack, H. J. (2014). Bee species diversity enhances productivity and stability in a perennial crop. *PloS one*, 9(5):e97307.
- 492 [36] Sachs, J. L., Mueller, U. G., Wilcox, T. P., and Bull, J. J. (2004). The evolution of cooperation. *Quarterly Review of Biology*, 79(2):135–160.
- 495 [37] Song, C., Ahn, S. V., Rohr, R. P., and Saavedra, S. (2020). Towards Probabilistic Understanding About the Context-Dependency of Species Interactions. *Trends in Ecology and Evolution*, 35(5):384–396.
- 498 [38] Stachowicz, J. J. and Whitlatch, R. B. (2005). Multiple mutualists provide complementary benefits to their seaweed host. *Ecology*, 86(9):2418–2427.
- [39] Stanton, M. L. (2013). Interacting Guilds: Moving beyond the Pairwise Perspective on Mutualisms. *The American Naturalist*, 162:510–523.
- 501 [40] Thrall, P. H., Hochberg, M. E., Burdon, J. J., and Bever, J. D. (2007). Coevolution of symbiotic mutualists and parasites in a community context. *Trends in Ecology and Evolution*, 22(3):120–126.
- 504 [41] Trager, M. D., Bhotika, S., Hostetler, J. A., Andrade, G. V., Rodriguez-Cabal, M. A., Mckeeon, C. S., Osenberg, C. W., and Bolker, B. M. (2010). Benefits for plants in ant-plant protective mutualisms: A meta-analysis.

507 [42] Ushio, M. (2020). Interaction capacity underpins community diversity.

Tables

Parameter	Median (95% CI)	Prior Distribution
growth xi intercept vacant β_{01}^g	-5.210899(-5.686865, -5.491787)	sDE
growth xi intercept other β_{02}^g	-5.8288(-5.956217, 1.766021)	asdf
growth xi intercept <i>C. opuntiae</i> β_{03}^g	-4.529523(-6.0770390, 0.1222112)	asdf
growth xi intercept <i>L. apiculatum</i> β_{04}^g	-5.106802(-5.4499944, 0.5453901)	asdf
growth xi size dependent vacant β_{11}^g	asdf	asdf
growth xi size dependent other β_{12}^g	asdf	asdf
growth xi size dependent <i>C. opuntiae</i> β_{13}^g	asdf	asdf
growth xi size dependent <i>L. apiculatum</i> β_{14}^g	sadf	asdf
growth omega intercept ω_0^g		
growth omega size dependent ω_1^g		
growth alpha intercept α_0^g		
growth alpha size dependent α_1^g		
1-year germination intercept α^{γ_1}		
2-year germination intercept α^{γ_2}		
survival intercept vacant β_{01}^s		
survival intercept other β_{02}^s		
survival intercept <i>C.opuntiae</i> β_{03}^s		
survival intercept <i>L. apiculatum</i> β_{04}^s		
survival size dependent vacant β_{11}^s		
survival size dependent other β_{12}^s		
survival size dependent <i>C. opuntiae</i> β_{13}^s		
survival size dependent <i>L. apiculatum</i> β_{14}^s		
flowering intercept β_0^f		
flowering size dependent β_1^f		
viability intercept vacant $\beta_0 1^v$		
viability intercept other $\beta_0 2^v$		
viability intercept <i>C. opuntiae</i> $\beta_0 3^v$		
viability intercept <i>L. apiculatum</i> $\beta_0 4^v$	28	

Figure legends

⁵¹⁰ Videos If you have videos, journal style for them is generally similar to that for figures.