

Demographic consequences of partner diversity and turnover in a multi-species ant-plant mutualism

Alexandra Campbell^{1,†} (ORCID: 0009-0008-4893-8029)

Tom E.X. Miller¹ (ORCID: 0000-0003-3208-6067)

¹ Program in Ecology and Evolutionary Biology, Department of BioSciences, Rice University, Houston, Texas 77005

[†] Corresponding author: ali.campbell@berkeley.edu

Keywords: Integral Projection Model, *Cylindropuntia imbricata*, multi-species mutualism, complementarity, sampling effect, portfolio effect

Abstract

Mutualisms commonly involve multiple partner species but the ecological consequences of partner diversity remain poorly understood. Partner diversity may benefit a focal mutualist through mechanisms that mirror positive effects of species diversity on ecosystem function, such as sampling effect, complementarity or portfolio effect. In sequential mutualisms, partner identity can change over time due to environmental variability or competitive displacement. Here we asked how partner diversity and turnover influence host demography and long-term fitness, and which mechanisms explain these effects. We studied the sequential multi-species food-for-protection mutualism involving an EFN-bearing plant (tree cholla cactus *Cylindropuntia imbricata*) and multiple species of nectar-feeding ants (*Crematogaster opuntiae*, *Liometopum apiculatum*, and other less frequent ants) that provide defense from herbivores. We used a 20-year dataset on plant vital rates and part-

ner identity to quantify the demographic consequences of association with different ant partners and the frequency and direction of turnover between mutually exclusive partner associations across the plant life cycle. We then constructed a stochastic, multi-state Integral Projection Model (IPM) to investigate how different patterns of partner richness, composition, and sequential turnover affect population fitness under fluctuating environmental conditions. We found that, while ant partners had different impacts on plant vital rates (*C. opuntiae*-tended plants had advantages in growth and survival when small, and *L. apiculatum*-tended plants had floral viability advantages), plant fitness was similarly elevated by ant visitation for any level of partner identity and richness, because effects on the highest-sensitivity vital rates were consistent across partner species. Furthermore, there was no evidence that benefits of partner diversity manifest in the context of fluctuating environments (i.e., the portfolio effect). In contrast to much of the previous literature, this study highlights that demographic benefits of mutualism can be surprisingly robust to diversity and turnover of partner species.

Introduction

Mutualisms, species interactions where all participants receive net benefits, are key components of ecological communities and can enhance the fitness and population growth rates of the partners involved. While traditionally framed as pairwise (+/+) interactions, mutualisms are often embedded within multispecies communities in which a focal mutualist interacts with multiple partner species. Growing evidence suggests that pairwise interactions may be poor predictors of the net effects of multi-species mutualism (Afkhami et al., 2014; Bascompte, 2009; Dattilo et al., 2014; Palmer et al., 2010).

A focal mutualist may interact with multiple guilds of partner types (e.g., plants that interact with pollinators, seed dispersers, soil microbes, and ant defenders) or with multiple partner species within the same guild (e.g., plants visited by multiple pollinator species). Within a mutualist guild, partner species often differ in the amount or type of goods or services they provide, making partner identity an important source of contingency in mutualism (Stanton, 2003). Partner diversity can arise either from simultaneous interactions or sequential ones, where individual hosts experience turnover in partner identity over time (Djiéto-Lordon et al., 2005; Ness et al., 2006). Understanding how partner diversity and turnover affect the overall fitness benefits of mutualism is an open question, especially in the context of environmental variability, where different partners may confer benefits unevenly across space or time (Frederickson, 2005; Palmer et al., 2010; Rogers et al., 2014; Thibaut et al., 2012).

There are multiple mechanisms by which partner diversity can influence the net benefits accrued by a focal mutualist, mirroring the mechanisms by which, at a larger scale of organization, biodiversity can influence ecosystem function (Barrett et al., 2015; Ushio, 2020; Winfree, 2020). First, when there is a consistent ranking of best to worst mutualists, a more diverse sample of the partner community may be more likely to include the best partner (Frederickson, 2013). This can lead to an apparent benefit of diversity driven by a sampling effect (Batstone et al., 2018). However, if partner associations are mutually exclusive (a focal mutualist may engage with only one partner at

a time), then partner diversity may impose opportunity costs, leading to negative effects of a diverse mutualist assemblage relative to exclusive association with the single best partner (Miller, 2007). Second, even within a mutualist guild, the benefits conferred by alternative partner species can vary by method, efficiency, or timing (Bronstein et al., 2006; Stachowicz and Whitlatch, 2005; Stanton, 2003). This can lead to a positive effect of partner diversity through complementarity, whereby diversity increases the net benefits received by focal mutualist beyond what a single partner species could offer (Batstone et al., 2018). Positive diversity effects through complementarity need not be additive: interference or synergies between partners can make their combined effect different than expected from the sum of complementary functions (Afkhani et al., 2014). Third, partner species can have species-specific responses to environmental variation, either spatially (Ollerton et al., 2006) or temporally (Alarcón et al., 2008). Multiple partners can therefore act as a ‘portfolio’ that stabilizes fitness benefits across spatial or temporal heterogeneity, leading to positive effects of partner diversity through the portfolio effect (Batstone et al., 2018; Horvitz and Schemske, 1990; Lázaro et al., 2022). In the context of temporally fluctuating environments, the portfolio effect implies that inter-annual variation in partner identity could stabilize overall plant fitness across years. For such a portfolio effect to operate, two conditions must be met: (i) partner species must differ in the timing or magnitude of their responses to environmental variation (i.e., their benefits fluctuate asynchronously), and (ii) the combined effect of these fluctuations must result in increased net benefit to the host across years relative to the net benefit expected if all partners responded in parallel (Hautier et al., 2014; Loreau, 1999; Tilman et al., 1994). While portfolio effects have been well studied in plant and animal communities, their role in mutualisms—particularly where partner diversity is expressed sequentially rather than simultaneously—remains less explored (Ushio, 2020).

For exclusive, sequential associations, the timing, frequency, and directionality of partner turnover can influence the net fitness benefits to a focal mutualist, particularly if turnover allows accumulation of complementary benefits or buffers environmental variability via portfolio effects (Fiala et al., 1994; Sachs et al., 2004; Ushio, 2020). Turnover can happen at different timescales, from minutes to years (Horvitz and Schemske, 1986; Nakazawa, 2020; Oliveira et al., 1999; Ventre-Lespiauq et al.,

2021). If partner identity changes systematically across ontogeny, such as through succession of partners aligned with host life stages, this may further affect fitness outcomes associated with partner diversity (Dejean et al., 2008; Fonseca and Benson, 2003; Noe and Hammerstein, 1994). For example, plant susceptibility to enemies can change across life stages (Barton and Koricheva, 2010; Boege and Marquis, 2005), so the benefits of a diverse guild of defensive mutualists are greatest when more defensive partner species align with more vulnerable life stages (Dejean et al., 2008; Djiéto-Lordon et al., 2005). Thus, the duration and order of interactions may be just as important as the identities of the interacting partners.

Defensive ant-plant mutualisms – where plants provide food and/or housing to ants that in turn defend them from enemies – are widespread interactions that offer valuable model systems for the ecology and evolution of mutualism (Bronstein, 1998; Bronstein et al., 2006). Extrafloral nectar (EFN) -bearing plants can serve as dietary resources that promote ant abundance and colony size (Byk and Del-Claro, 2011; Donald and Miller, 2022; Ness et al., 2006, 2009). In turn, presence of defensive ant partners is often linked to reductions in herbivory (Rudgers, 2004; Trager et al., 2010) and demographic advantages for the plant partner (Báez et al., 2016). Defensive ant-plant mutualisms are commonly multi-species, where a guild of ant partner species share, and often compete for, a plant mutualist (Agrawal and Rutter, 1998; Beattie, 1985; Bronstein, 1998; Trager et al., 2010). Ant partners can vary in their ability to deter herbivores (Bruna et al., 2014) and visitation by low quality ant partners can prevent visitation by higher quality partners, causing a reduction in relative fitness through opportunity costs (Fraser et al., 2001; Frederickson, 2005). Susceptibility to herbivory can also vary significantly throughout the life stages of the plant (Boege and Marquis, 2005), suggesting that the order and timing of successive partners is important to the fitness impacts of the combined partner guild (Barton and Koricheva, 2010; Boege and Marquis, 2005; Fonseca and Benson, 2003). Finally, herbivore identity and pressure can vary inter-annually (Wetzel et al., 2023), much like mutualist identity and presence, meaning the threat plants face can vary just as much as the protection they receive due to temporal stochasticity. Previous studies have investigated how ant partner diversity affects plant fitness (Dattilo et al., 2014; Fiala et al.,

1994; Gaume et al., 1998; Ludka et al., 2015; Palmer et al., 2010), however little is known about the combined effects of partner identity, directional partner turnover, and temporal stochasticity, particularly because long-term demographic studies on ant-plant interactions are exceedingly rare (Donald and Miller (2022); Moraes and Vasconcelos (2009)).

Long-term demographic data are essential to understanding the fitness consequences of multi-species mutualisms for several reasons. First, fitness effects of sequential partnering play out over long time scales due to the accumulation of partners across time rather than the effects of multiple partners at once. While turnover of mutualist partners is known to occur (Djiéto-Lordon et al., 2005; Fonseca and Benson, 2003; Ness et al., 2006), long-term data documenting patterns and consequences of partner switching are uncommon, especially in field settings (Palmer et al., 2010). Second, certain effects of partner diversity are fluctuation-dependent (i.e., portfolio effect) and would only be detectable in the context of temporally stochastic environments in which alternative partners may fluctuate asynchronously. Quantifying contributions from the portfolio effect therefore requires long-term data on partner (a)synchrony across inter-annual fluctuations. Finally, alternative partners may have different demographic effects (e.g., one that enhances survival and another that enhances reproduction), and these effects may contribute differently to overall fitness of a focal mutualist. Most studies evaluate short-term fitness proxies, such as seed set or herbivory, often focusing on a single life stage (Palmer et al., 2010; Trager et al., 2010), while fewer trace demographic effects across multiple stages (Bruna et al., 2014). Integrating long-term data from across the life cycle within a demographic modeling framework therefore provides a powerful opportunity to diagnose the fitness consequences of partner diversity and turnover in the context of environmental stochasticity.

This study examined the consequences of partner diversity in a food-for-protection mutualism between the tree cholla cactus (*Cylindropuntia imbricata*), a long-lived EFN-bearing plant, and multiple species of ant partners. Tree cholla are tended by two common ant partners and several additional rarer species, all of which collect EFN in exchange for defense against herbivores. These ant species locally co-occur, but individual plants are typically tended by only one species that

patrols the plant around-the-clock and maintains control of the plant’s nectar resources, usually for an entire growing season (Donald and Miller, 2022; Ohm and Miller, 2014). Switches between partner species, or between vacancy and ant occupancy, commonly occur from one growing season to the next (Miller, 2007). Prior experiments suggested a hierarchy of mutualist quality, with *Liometopum apiculatum* providing strong anti-herbivore defense and *Crematogaster opuntiae* having net negative effects because herbivore deterrence is outweighed by deterrence of pollinators (Miller, 2007; Ohm and Miller, 2014). However, previous studies in this system focused on single life stages (adult plants) or vital rates (seed production), did not integrate the demographic effects of ant defense across the life cycle, and did not incorporate temporal stochasticity – gaps we aimed to fill in the present study.

Here we used a unique long-term data set that allows us to explore mutualistic associations with multiple partner species, longitudinal dynamics of partner turnover, and demographic effects of alternative partners that vary across plant size structure and across nearly 20 years of inter-annual fluctuations. The long-term nature of this observational dataset gives us the power to investigate whether and through which mechanism(s) partner diversity affects the fitness benefits of ant visitation. Specifically, we asked: What are the vital rate effects of association with alternative partners and how do these effects fluctuate across years? What are the frequency and direction of partner turnover across the plant life cycle? What is the net effect of partner diversity on plant fitness, and what mechanism(s) explain(s) this effect?

To answer these questions we used a hierarchical Bayesian statistical approach to estimate demographic vital rates for hosts in different states of ant occupancy and to quantify state-dependent partner turnover from the long-term data. We then built a stochastic, multi-state integral projection model (IPM) which combines diverse effects on vital rates and pathways of partner turnover. Finally, we used the model to quantify plant fitness under different scenarios of partner richness, composition, and synchrony.

Methods

Study System

This study was conducted at the Sevilleta Long-term Ecological Research site (SEV-LTER), located within the Sevilleta National Wildlife Refuge in central New Mexico, USA. Our study area in the Los Piños mountains ($34^{\circ}20'5.3''\text{N}$, $106^{\circ}37'53.2''\text{W}$) is characterized by steep, rocky slopes and perennial vegetation including grasses (*Bouteloua eriopoda* and *B. gracilis*), yuccas, cacti, and junipers. Tree cholla cacti are common in such high Chihuahuan desert habitats (Benson, 1982). These arborescent plants produce cylindrical segments with large spines. Early each growing season (May to August in New Mexico), they initiate new vegetative segments and flowerbuds at the ends of existing segments. While most plants produce new segments every season, only those which are reproductively mature produce flowerbuds. Like other EFN-bearing cacti, tree cholla secrete nectar from specialized glands on young vegetative segments and flowerbuds (Ness et al., 2006; Oliveira et al., 1999). Flowerbuds produce more and higher-quality EFN than vegetative segments, making reproductive cholla valuable mutualist partners (Miller, 2014). Small, non-flowering cholla produce little to no EFN and are commonly vacant (no ant visitation) (Miller, 2014).

Tree cholla EFN is collected by several ground-nesting ant species. At SEV-LTER, cholla are visited primarily by two species, *Crematogaster opuntiae* and *Liometopum apiculatum*, as well as other rarer species, including *Forelius pruinosus* and unidentified species in the genera *Aphaenogaster* and *Camponotus*. *L. apiculatum* are the most frequent visitors with 25% – 60% of tree cholla tended by these ants in a given year, followed by *C. opuntiae* (5% – 20%) (Donald and Miller, 2022). Between 30% – 80% of cacti are vacant in any given year. Workers of different species rarely co-occur on individual plants, likely due to interspecific competition. For example, staged introductions of *C. opuntiae* to *L. apiculatum*-tended plants, and vice versa, provoke aggressive responses by residents (*personal observation*).

Several insect herbivores and seed predators specialize on tree cholla (Mann, 1969), and defense

against these enemies is the main pathway by which ant visitation affects plant demography. The Cerambycid beetle *Moneilema appressum* and an unidentified weevil (Coleoptera: Curculionidae) of the genus *Gerstaekeria* feed on vegetative and reproductive structures as adults and their larvae feed internally. Two species of cactus bugs, *Narnia pallidicornis* and *Chelinidea vittiger* (Hemiptera: Coreidae), feed on all cholla parts with a preference for flower buds; their damage can induce floral abortion (Miller et al., 2006). A seed predator, *Cahela ponderosella* (Lepidoptera: Pyralidae), oviposits in open flowers and larvae eat seeds in developing fruits. These consumers can have significant negative impacts on plant fitness and depress population growth (Miller et al., 2009). Prior experiments showed that ant-tended tree cholla experience less herbivory and seed predation than plants from which ants were excluded (Miller, 2007; Ohm and Miller, 2014).

Data Collection

This study is based on long-term data spanning 2004 to 2023 that documented plant demographic rates in relation to their ant partner status. From 2004 to 2008, we censused 134 plants distributed across three spatial blocks. This initial census group was discontinued in 2009, when we established six 30×30 -meter plots and tagged all tree cholla within those plots. Two additional 30×30 -meter plots were added in 2011, and this group of eight plots (mean $\pm$ SD tree cholla per plot: 87.30 ± 18.28) was censused annually through 2023 (with the exception of 2020 due to the pandemic shutdown). For all plants, in May or early June of each year we recorded plant survival since the last survey (if the plant had any green flesh it was considered alive) and, for survivors, we recorded height (cm), maximum crown width (cm), and crown width perpendicular to the maximum (cm). Size measurements were used to calculate plant volume (cm^3) based on the volume of an elliptical cone. We measured reproduction by counting flowerbuds, and in most years we distinguished between flowerbuds that were viable versus aborted (aborted flowerbuds are distinguishable by dry and withered true leaves and petals). We recorded the ant species presence and identity (or vacancy if no ants present). Occurrences of more than one ant species on one plant were rare (less than 5% of observations), and for the purpose of this analysis we classified the plant as being

occupied by the more abundant species. We also quantified the abundances and identities of any cactus herbivores observed, and recorded the occurrence of herbivore damage (described further in Appendix ??). Each year, the field team searched exhaustively for new recruits by walking slowly in a standard formation across the plot area, and these were added to the census (cactus seedlings are small ($< 2\text{cm}$ in height) and cryptic so it is likely that some are missed even with a standardized search protocol). These data allowed us to link each plant’s demographic fate (survival, growth, and reproduction) to its state of ant visitation. In total, the data set includes a total of 9,787 observations of 1,141 individuals across 15 complete transition years (spanning May/June of year t to May/June of year $t + 1$) (Miller, 2020). In addition to missing the year 2020, there are gaps in the time series where we switched plots or plants (and thus broke up transition years for growth/survival and partner turnover) or where we did not distinguish between viable and aborted flower buds (and therefore are missing some vital rate information for those years).

To complete the tree cholla life cycle, we used additional, smaller data sets from previously published studies to estimate seed and seed bank parameters. First, Ohm et al. 2014 provide data on the number of seeds per fruit for plants tended by *L. apiculatum*, *C. opuntiae*, or no ants (experimental exclusion), accounting for their effects on pollinator visitation. Second, Elder and Miller 2016 provide data on seed entry to the seed bank and seedling germination and survival rates based on surveys of fruit predation and seed addition experiments. Because these seed and seed bank data were not collected as part of our long-term study, these demographic parameters are based on data from a single year and are assumed to be temporally constant.

Multi-state Integral Projection Model

Model overview

The demographic data were used to parameterize a multi-state Integral Projection Model (IPM). IPMs describe population dynamics in discrete time with functions that relate vital rates to continuous state variables, typically size (Ellner et al., 2016). While IPMs are a natural choice for

populations with continuous size structure, they can also be modified to accommodate a combination of continuous and discrete state variables, as we do here. We constructed a stochastic, multi-state IPM that stitches together population structure associated with plant size and ant state, allowing us to determine the individual fitness effects of each ant species in isolation and the composite effects of multiple partners, with ant transition dynamics and inter-annual variability modeled explicitly.

Given the low frequency of ant occupancy states other than *L. apiculatum* and *C. opuntiae* (<8% of observations) we combined all other ants into an “Other” category, such that our multi-state IPM included four possible ant states: vacant, *L. apiculatum*, *C. opuntiae*, and Other. The “Other” category was made up of *Forelius pruinosus* (3.5% of observations), unidentified species belonging to the genera *Camponotus* (0.9%), *Aphaenogaster* (0.4%), *Myrmecocystus* (0.08%), *Tetramorium* (0.02%), *Brachymyrmex* (0.02%), and additional ants not identified to genus or species (2.8%). Given our objective of quantifying demographic effects across the plant size distribution, there was not sufficient size representation of these low-frequency partner species in the long-term data to treat these as separate partner states.

Ant state is included as a predictor variable in IPM sub-models where there are biologically realistic pathways through which ants could have an impact. For example, prior experimental work indicates that ant tending can reduce vegetative tissue loss and floral abortion, while also potentially reducing pollinator visitation (Miller et al., 2009; Ohm and Miller, 2014). Therefore, ant state was included in sub-models for survival, growth, flowerbud viability, and seed number per flowerbud. In contrast, we have no reason to expect that ant tending can directly influence the probability of flowering or flowerbud production independently of its influence on plant size, so these sub-models do not include ant state as a predictor variable.

The biological “story” of the model is as follows: The tree cholla population is comprised of plants characterized by their size, which can take any continuous value between lower and upper limits, and ant visitation status, which can take one of the four discrete levels described above. An individual’s size and ant status jointly predict its demographic fate such as survival, growth, and

reproductive contributions to the seed bank. Plants may be lost to mortality and new individuals may be gained through recruitment from the seed bank. Size and ant status are updated annually. The environment fluctuates from year to year such that demographic rates exhibit stochastic deviations from their long-term average. Plants in different ant states can fluctuate independently; for example, a year that is better than average for *L. apiculatum*-tended plants may be worse than average for *C. opuntiae*, and vice versa, which could occur if there are year-to-year fluctuations in the natural enemies against which each defender species is most effective. Our long-term demographic data define the “rules” of tree cholla demography and ant partner transition. In the analyses that follow, we tinker with those rules to create novel scenarios of partner diversity and inter-annual synchrony.

Mathematical formulation

Mathematically, we modeled the tree cholla life cycle using continuously size-structured plants where number of plants of size x and ant state a in year t ($n(x, a)_t$) predicts the number of plants of size x' and ant state a' in year $t + 1$ ($n(x', a')_{t+1}$) based on a size- and ant-specific vital rates. The model also includes two discrete seed banks (B_t^1 and B_t^2) corresponding to 1 and 2-year old seeds. Seed bank dynamics are given by:

$$B_{t+1}^1 = \delta \sum_{a=1}^A \int_L^U \kappa(a) P(x; \boldsymbol{\tau}^P) F(x; \boldsymbol{\tau}^F) V(a; \boldsymbol{\tau}_a^V) n(x, a)_t dx \quad (1)$$

$$B_{t+1}^2 = (1 - \gamma_1) B_t^1 \quad (2)$$

Functions $P(x; \boldsymbol{\tau}^P)$ and $F(x; \boldsymbol{\tau}^F)$ give the probability of flowering in year t and the number of flowerbuds produced in year t , respectively, by plants of size x in year t . The proportion of flowerbuds that remain viable through fruit set ($V(a; \boldsymbol{\tau}_a^V)$) and the number of seeds per fruit ($\kappa(a)$) are dependent on ant state a . The vectors $\boldsymbol{\tau}$ give year-specific deviates (mean zero) and appear in functions for which we can estimate temporal stochasticity from the long-term data; superscripts indicate the corresponding vital rate and, when present, the a subscript indicates that deviates are

specific to plants in ant state a . For example, temporal deviates τ_a^V describe better- and worse-than-average years for flowerbud viability, and plants in different ant states can fluctuate independently. Seed production is integrated over the size distribution, from the lower L to upper U size limits, and summed over all possible ant states ($A = 4$) giving total seed production. Seeds are multiplied by the probability of escaping post-dispersal seed predation (δ) to give the number of seeds that enter the one-year old seed bank. Plants can recruit out of the year-one seed bank with probability γ_1 or transition to the two-year seed bank with a probability of $1 - \gamma_1$. Seeds in the two-year seed bank are assumed to either germinate with probability γ_2 or die.

For the above-ground part of the life cycle, the number of plants of size x' and ant state a' in year $t + 1$ is given by survival/growth transitions from size x and ant state a in year t , plus germination out of the seed banks:

$$n(x', a')_{t+1} = (\gamma_1 B_t^1 + \gamma_2 B_t^2) \eta(x') \omega \rho_0(a') + \sum_{a=1}^A \int_L^U S(x, a; \tau_a^S) G(x', x, a; \tau_a^G) \rho(x, a, a') n(x, a)_t dx \quad (3)$$

The first term in Eq. (3) estimates the number of individuals recruiting from a one- or two-year seed bank to a plant of size x' and ant state a' based on the recruit size distribution $\eta(x')$ and the probability of over-winter seedling survival (ω) from germination (late summer) to the census (May). This term is multiplied by $\rho_0(a')$, which gives the probability that a new recruit has ant state a' upon its first appearance in the census ($\sum \rho_0(a') = 1$). The second term represents all possible transitions from size x and ant a to size x' and ant a' , conditioned on survival. Survival ($S(x, a; \tau_a^S)$) and growth from size x to x' ($G(x', x, a; \tau_a^G)$) are both dependent on initial size and ant state. As above, these functions include inter-annual variability through year-specific deviates that can vary by ant state (τ_a). Finally, ant transition function $\rho(a', a, x)$ gives the probability that an individual transitions from ant state a to a' in the next census, conditional on initial size x .

Statistical modeling and parameter estimation

We parameterized the IPM using a series of generalized linear mixed models in a hierarchical Bayesian framework. Vital rate models included spatial and temporal random effects associated with plot and year variation, respectively (only year variation is used in the IPM), and included plant size (the natural logarithm of volume, $\log(\text{cm}^3)$; x, x'), ant partner state (a, a'), or both as fixed-effect predictor variables. As in the IPM, our statistical modeling assumed that demographic effects of ant occupancy are limited to survival, growth, and flowerbud viability.

Growth. Following best practices for modeling size transitions (Miller and Ellner, 2025), we fit the growth sub-model ($G(x', x, a; \tau_a^G)$) to data on size in year $t + 1$ (y^G) using the skewed normal distribution to account for left-skewed size transitions (at some initial sizes, transitions below the expected future size were more common than transitions above it). The skew-normal has three parameters corresponding to location ($\hat{G}$), shape (σ), and scale (α):

$$y_i^G \sim \text{SkewedNormal}(\hat{G}_i, \sigma_i, \alpha_i) \quad (4)$$

$$\hat{G}_i = \beta_{a[i]}^0 + \beta_{a[i]}^1 x_i + \beta_{a[i]}^2 x_i^2 + u_{\text{year}[i], a[i]} + w_{\text{plot}[i]} \quad (5)$$

$$\log(\sigma_i) = \beta^3 + \beta^4 x_i \quad (6)$$

$$\alpha_i = \beta^5 + \beta^6 x_i \quad (7)$$

Here, the location parameter for the i th observation $\hat{G}_i$ is defined as a second-order polynomial with ant-size interactions because preliminary analysis found this was an improvement over a linear relationship. The location parameter of the skew-normal is not the mean, but the mean can be derived as $\hat{G} + (\sigma\alpha/\sqrt{1+\alpha^2})\sqrt{2/\pi}$. The year- and ant-specific random effect u (which parameterizes the τ_a^G vectors) and plot-specific random effect w are normally distributed with variances σ_{year}^2 and σ_{plot}^2 , respectively (note that all ant states are assumed to share the same inter-annual variance σ_{year}^2 , even as they are allowed to fluctuate independently). Parameters σ_i and α_i control residual variance and skewness, respectively, and were defined as linear functions of initial size x_i (σ_i is

strictly positive and was modeled with a log link function). Due to data constraints, we modeled growth variance and skewness as size-dependent but not dependent on ant occupancy state.

Survival. The survival sub-model ($S(a, x; \tau_a^S)$) estimates the probability of survival from year t to year $t + 1$, with fixed effects of size x and ant partner a in year t . We fit this model to the survival data (alive or dead) using a Bernoulli distribution with a similar linear predictor for the probability of survival as in the growth model but with a logit link function and without the second-order influence of size.

Reproduction. The flowering sub-model ($P(x; \tau^P)$) estimates the probability of reproducing in year t , with fixed effects of size x in year t and random effects of plot and year. We fit this model to the reproductive status data (vegetative or flowering) using a Bernoulli distribution and a logit link function, similar to the survival model above but with no ant effects. The flowerbud function $F(x; \tau^F)$ estimates the total flowers produced by a reproducing plant in year t , with fixed effects of size x in year t . We fit this model to flowerbud count data (sum of viable and aborted buds) using a zero-truncated negative binomial distribution with a log link and normally distributed year and plot random effects.

The flowerbud viability sub-model ($V(a; \tau_a^V)$) estimates the probability that flowerbuds produced in year t remain viable, with fixed effects of ant partner a in year t . We fit this model to floral viability data using a binomial distribution where trials and successes are given by the total number of flowerbuds and the number that are viable, respectively. This model used a logit link function and included random effects for plot and year following the same structure as the survival model, with ant-specific year random effects.

Estimates for the number of seeds per fruit were obtained from a field experiment which excluded ants from flowering plants (Ohm and Miller, 2014). This experiment only included *L. apiculatum* and *C. opuntiae* -tended plants. We therefore assumed that plants tended by Other ants had the same number of seeds per fruit as vacant plants (our results showed very little sensitivity to this assumption). Additional reproductive parameters for probability of entry to the seed bank,

germination rates, and recruit size were estimated following methods described in Appendix ??.

All of these parameters are assumed to be temporally constant.

Ant Transitions. The ant transition model ($\rho(x, a, a'; \boldsymbol{\tau}^e)$) estimates the probability of a cactus being occupied by ant partner a' in year $t + 1$ given that it was occupied by initial ant partner a in year t , with fixed effect of initial size x . We fit this model to ant partner data using a multinomial distribution with a logit link function.

Model-fitting. We fit all models using Stan run through version 4.0.2 of R (R Core Team, 2023; Stan Development Team, 2023). We used vague priors for all parameters. For each model, we obtained three chains of 10,000 iterations, discarding the first 1,500 iterations. We visually assessed parameter convergence between and within chains and assessed overall model fit with posterior predictive checks to examine how well the fitted model can generate simulated data similar to the real data.

IPM Analysis

Analyzing the IPM required that we discretize the continuous IPM kernel into an approximating matrix. Size variable x was discretized into b bins, resulting in a $b \times b$ matrix. In our model there is additional complexity in the form of transitions between A ant states and two additional discrete states (year one and year two seed banks), leading to a matrix size of $A(b + 2) \times A(b + 2)$. We used $b = 500$ bins, which we found to be sufficient for numerically stable outputs, and extended the integration limits beyond the minimum (L) and maximum (U) observed sizes to avoid unintentional eviction using the “floor-and-ceiling” method (Williams et al., 2012).

For stochastic analyses, we estimated the approximating matrix corresponding to each t to $t + 1$ transition year. To estimate population mean fitness in a stochastic environment (λ_S) we simulated population dynamics for 500 years by randomly sampling among the 15 annual transition matrices, discarding the first 100 years of the simulation to minimize the influence of initial conditions.

Sampling observed transition matrices (rather than independently sampling regression coefficients) produces demographic time series that realistically capture inter-annual variation by preserving correlations between vital rates (Metcalf et al., 2015). We tallied the total population size at each time step as $N_t = B_t^1 + B_t^2 + \sum_{a=1}^A \int n(x, a)_t dx$ and calculated the stochastic growth rate as

$$\log(\lambda_S) = E[\log(N_{t+1}/N_t)]$$

(Rees and Ellner, 2009). We propagated uncertainty from the vital rate models using 100 draws from the joint posterior distribution of model parameters, resulting in a posterior distribution of λ_S and the derived quantities described next.

Partner diversity scenarios

Using the fully parameterized multi-state IPM, we conducted simulations to quantify how diversity and identity of ant partners influenced plant fitness. From the full version of the model (described above) corresponding to the observed assemblage of partners and observed frequencies of partner transition, we created alternative model versions corresponding to all eight “counter-factual” scenarios of diversity and composition: no ant partners (complete vacancy); one ant partner (*C. opuntiae* only, *L. apiculatum* only, Other only); two partners (all pairwise combinations of *C. opuntiae*, *L. apiculatum*, and Other); and three partners (observed scenario of all ant states). These simulations were made possible by extrapolating ant-specific demographic performance across the size distribution, even for combinations of size and ant occupancy that were rarely observed. For example, the no-partner scenario modeled a hypothetically ant-free cactus population, even though no such population exists to our knowledge, by applying the statistical knowledge gleaned from vacant plants across the size distribution. We refer to stochastic fitness associated with partner number or identity using superscripts, e.g. λ_S^0 for vacant plants (zero partners), λ_S^{1+} for any state of ant tending, λ_S^C for tending by only *C. opuntiae*, λ_S^{CO} for tending by *C. opuntiae* and Other ants, etc.

In all scenarios that included any ant partners, we preserved the observed pattern of size-

dependent vacancy/occupancy (estimated through the ant transition sub-model) and manipulated partner identity conditional on occupancy by any ant. This means, for example, that the *C. opuntiae*-only scenario included two possible states, vacancy and occupied by *C. opuntiae*. Our statistical models allow us to extrapolate the demographic performance of ant-tended plants to small sizes that are typically vacant, but the natural history of this system tells us that this is not biologically sensible. Small, non-reproductive plants are typically vacant because they do not produce extrafloral nectar, and once plants begin producing nectar they are nearly always ant-tended (Miller, 2014). Our simulations preserved this basic biology, avoiding tiny ant-occupied plants that do not and could not occur in nature. While technically such plants must exist in the model (because all ant states share the same L and U size limits), they are so infrequent, as guided by our data, as to have virtually no bearing on the results.

The partner diversity scenarios required additional assumptions about the mechanisms that give rise to observed occupancy patterns. Based on evidence that EFN-bearing cacti are nearly always ant-occupied (Miller, 2014), we assume that ant partners competitively exclude one another from EFN-bearing cacti and that competition for plant partners is zero-sum. This means that, in scenarios that remove species from the partner community, remaining species gain access to plants that the removed species would have tended. In Appendix ??, we present results under an alternative assumption, that ant visitation is limited by factors other than availability of cactus EFN (e.g., nesting sites or off-plant dietary resources), such that when a species is removed from the partner community, the plants it would have occupied remain vacant.

Partner synchrony scenarios

Under the portfolio effect hypothesis, partner diversity may confer a fitness advantage when the benefits of alternative partners are not perfectly synchronized across temporal environmental variation, yielding an advantage of a diverse “portfolio” of partners when the environment fluctuates. We constructed two versions of the stochastic, multi-state IPMs that allowed us to test this hypothesis. The baseline, “non-synchronous” model described above included ants effects that could

vary uniquely across time, according to the parameter estimates for the random effects (τ_a). We quantified from the fitted random effects how tightly inter-annual variation was correlated between ant states for each vital rate. The “synchronous” version included temporal fluctuations that were forced to be the same across ant states. To synchronize ant states, we averaged the ant-specific year random effects, thus ensuring that plants in all ant states fluctuated synchronously in response to temporal environmental variation. This second, synchronous version of the model effectively turns off any portfolio effect, holding all else equal. Both scenarios of temporal stochasticity, non-synchronized and synchronized, were run for all eight ant partner scenarios described above. In Appendix ??, we explore additional scenarios that vary the correlation of ant-specific fluctuations, to determine if contributions from the portfolio effect are constrained by high correlations seen in the long-term data (Results).

Statistical inference on fitness consequences of partner identity and diversity

The range of models we created could generate many outputs; we focus our inference on the following specific contrasts. First, to determine whether ant occupancy and partner diversity are beneficial, we calculated a posterior distribution of λ_S for each of four partner richness levels (λ_S^0 , λ_S^1 , λ_S^2 , λ_S^3), averaging over composition scenarios within each level. If cactus fitness increases with partner richness, this would be interpreted as evidence for benefits of partner diversity. Second, to determine whether each partner, in isolation, confers a fitness advantage and to rank alternative partners, we contrasted the fitness of each single partner scenario (λ_S^C , λ_S^L , λ_S^O) against vacancy (λ_S^0). Third, to determine whether any benefits of diversity are due to the sampling effect or complementarity, we contrasted the fitness of multi-partner scenarios against the single best partner scenario. If the best multi-partner scenario exceeds the fitness associated with the best single partner, this would be interpreted as evidence of complementarity. Alternatively, the sampling effect hypothesis predicts that no multi-partner scenario yields higher plant fitness than the best single partner. It is also possible that multi-partner scenarios yield lower fitness than the single best partner, which would be consistent with an opportunity cost of diversity. Fourth, to quantify any contribution of

the portfolio effect, we contrasted λ_S of the full (four-state) scenario to vacancy for synchronized and non-synchronized responses to temporal stochasticity. If the portfolio effect confers a benefit of diversity, the fitness advantage of having all vs. no partners should be greater when temporal fluctuations are not synchronized across ant states.

We base our statistical inferences on the posterior probability distributions of the contrasts described above. For example, the contrast of *C. opuntiae* (λ_S^C) with vacancy (λ_S^0) yields a posterior distribution of the difference $\Delta\lambda_S^{C-0}$. We can quantify from this distribution our certainty in the mutualistic effect of *C. opuntiae*, given the data, as $Pr(\Delta\lambda_S^{C-0} > 0)$. We apply similar logic to other contrasts described above. We interpret contrast probabilities ≥ 0.95 as statistically significant differences.

Results

What are the effects of association with alternative partners on vital rates and how do these effects fluctuate across years?

Over the 20-year data set, we found that ant partners influenced cactus demographic performance, and different ant partners had contrasting effects across host vital rates. Ant-tended plants had a growth advantage over vacant plants, especially at smaller sizes (Figure 1). For the smallest sizes that were likely to be ant-tended (minimum observed size of ant-tended plants was $0.8 \log(\text{cm}^3)$; solid lines in Figure 1) there were modest differences between partner species, with the greatest growth advantage associated with *C. opuntiae* followed by *L. apiculatum* and then Other ants. At the largest sizes, growth trajectories of ant-tended and vacant plants were nearly indistinguishable. For all ant states, growth was left-skewed, with transitions to sizes below the mean being more frequent than sizes above the mean.

Similarly, for plants that were large enough to have ant visitors, visitation enhanced cactus survival (Figure 2). Mean survival rates ranged from 7.7% to 99.9%, with the smallest plants

being the most vulnerable to mortality. *C. opuntiae*-occupied plants had a survival advantage over other ant-tended plants, particularly at smaller sizes, consistent with positive effects on growth. At larger sizes, plants in any state of ant occupancy had a survival advantage over vacant plants. Plants smaller than $-2 \log(cm^3)$ were predicted to experience negative effects of ant visitation, but this was based entirely on extrapolated survival estimates of ant-tended plants. Plants in this size range were never observed to be ant-tended and, because the IPM preserves the size-dependence of vacancy, benefits of vacancy at small sizes are never realized in the IPM.

Figure 1:

Figure 2:

Ant visitation was associated with increased floral viability and ant identity determined the strength of viability benefits. Mean viability rates spanned 55–81% (Figure 3). *L. apiculatum*-tended plants had the highest mean viability rate (86.1% [95% credible interval: 77.6–92.4%]), while there were similar viability rates for vacant (60.0% [44.3–75.0%]), Other-tended (60.6% [43.7–75.5%]), and *C. opuntiae*-tended plants (57.1% [40.6–72%]). Furthermore, *C. opuntiae*-tended plants had lower mean seeds per fruit (115.0[79.5–165.5] seeds) than vacant (147.2[114.1–189.9] seeds) or *L. apiculatum*-tended plants (142.4[100.7–200.2] seeds), likely due to pollinator interference (Ohm and Miller, 2014).

Figure 3:

Inter-annual fluctuations in demographic rates (estimated as statistical random effects) were generally positively correlated between ant states, limiting the potential for benefits of diversity through the portfolio effect (Figure 4). However, the degree of correlation depended on the vital rate and pair of ant partners. Across vital rates, random effects for cactus growth were the most strongly correlated between ant states (mean pairwise correlation: 0.63) and random effects for survival were the least correlated (mean pairwise correlation: 0.36). Certain partner pair / vital

rate combinations were effectively independent in their yearly fluctuations (such as survival of *C. opuntiae* and Other-tended plants; mean pairwise correlation: -0.0016) while others were almost completely synchronous (such as growth of Other and *L. apiculatum*-tended plants; mean pairwise correlation: 0.93).

Figure 4:

What are the frequency and direction of partner turnover across the plant life cycle?

A majority (55%) of plants surveyed in the long-term data experienced at least one ant state transition, with distinct size-dependent and directional patterns (Figure 5). Vacancy was the most likely ant state of small plants ($\leq 10 \log(\text{cm}^3)$). Even when small plants were ant-tended at the start of the transition year, they were most likely to transition back to vacancy (Figure 5 b-d). The probability of becoming ant-tended increased with size, though it was not equally likely to be tended by any partner. For large plants that were initially vacant or tended by *L. apiculatum* or Other ants, *L. apiculatum* was the most likely next partner, suggesting that this species is able to colonize plants that were previously vacant or occupied by Other ants, and effectively retain plants that it previously occupied. *C. opuntiae* were also able to retain plants they previously occupied, but not as well as *L. apiculatum*: for plants that began the transition year with *C. opuntiae*, the probability that those plants remain occupied by *C. opuntiae* at the end of the transition year was only slightly greater than the probability of take-over by *L. apiculatum*, while take-over in the other direction was extremely rare. It is also notable that transitions away from the initial state of *L. apiculatum* were almost always transitions to vacancy (Figure 5 d), while transitions away from the initial states of *C. opuntiae* and Other were often transitions to other ants. This suggests a competitive hierarchy whereby *L. apiculatum* may abandon low-value plants with little nectar production but is almost never displaced from high-value plants.

Figure 6:

Figure 5:

What is the net effect of partner diversity on plant fitness, and what mechanism(s) explain(s) this effect?

By integrating vital rates, temporal fluctuations, and ant transition dynamics into the stochastic, multi-state IPM we can evaluate the fitness implications of different scenarios of partner identity and diversity (Figure 6). First, there was strong evidence that ant visitation had mutualistic fitness effects on plant partners. The lowest mean stochastic fitness was λ_S^0 , the fitness of the cholla with no partners (Figure 6 a). Across all 1+ partner scenarios, we found that the λ_S^{1+} posterior distributions were greater than λ_S^0 with nearly 100% certainty. This indicates that ant visitation elevates fitness no matter the number of partners. Furthermore, we found no benefit of partner diversity, with the fitness associated with one, two, or three partners roughly equivalent (Figure 6 a). The one- and two-partner scenarios were statistically consistent but there was a modest reduction in fitness in the three-partner scenario compared to two partners ($Pr(\Delta\lambda_S^{2-3} > 0) = 0.95$), consistent with a weak cost of diversity at the highest level. Patterns of λ_S in relation to partner richness were generally consistent between scenarios of non-synchronous and synchronous inter-annual fluctuations.

Partner identity and composition were not strongly consequential for plant fitness (Figure 6 b-c). Among the one-partner scenarios, there was no strong evidence for any single best partner species (Figure 6 b). While a hypothetical *L. apiculatum*-only population had the highest mean fitness, it was not significantly higher than *C. opuntiae*- ($Pr(\Delta\lambda_S^{L-C} > 0) = 0.7$) or Other- ($Pr(\Delta\lambda_S^{L-O} > 0) = 0.69$) only populations. Furthermore, the fitness of *L. apiculatum*-only was consistent with all 2-partner scenarios ($Pr(\Delta\lambda_S > 0)$ ranged from 0.37 to 0.83). However, as above, there was evidence for an opportunity cost of diversity wherein fitness of the three-partner scenario was lower than any of the 2-partner scenarios ($Pr(\Delta\lambda_S > 0) > 0.95$). The lack of diversity benefits are not

driven by the high overall frequency of *L. apiculatum*. Using simulations in which all ant species were artificially forced to be equally frequent (further explained in Appendix ??), we found the same fitness patterns as described above.

We found no evidence that the portfolio effect generated positive effects of partner diversity, as effects of partner richness and composition were highly consistent between the baseline, non-synchronous model and the model version that synchronized all ant states (Figure 6). The effect of all ant partners can be measured as $\lambda_S^3 - \lambda_S^0$. We have high confidence that this contrast is positive and equally so for the synchronous and non-synchronous scenarios (Figure S??), indicating that the full suite of partners is similarly beneficial with and without the portfolio effect operating. In Appendix ??, we show that contributions from the portfolio effect were constrained by positively correlated fluctuations between ant states (Figure 4). In scenarios in which partner fluctuations were uncorrelated or, especially, negatively correlated, the non-synchronous model exhibited a stronger benefit of diversity than the synchronous model (Figure ??), as expected under the portfolio effect hypothesis.

This lack of evidence for a sampling effect likely stems from the strong competitive dominance and high frequency of *L. apiculatum*, which restricts the population-level impacts of alternative partners. To explore the conditions under which a sampling effect might emerge, we conducted an alternative simulation in which ant species did not competitively exclude one another (Appendix ??). In this counterfactual scenario, we found that a sampling effect can indeed emerge when partner transitions are governed by random colonization rather than competitive hierarchies. This suggests that the benefits of partner diversity are highly sensitive to the rules of partner turnover; in systems like the cholla-ant mutualism, where a single high-quality partner is numerically and competitively dominant, the potential for diversity to increase fitness via sampling is constrained.

Discussion

Mutualisms commonly involve multiple partners but the ecological consequences of partner diversity remain poorly understood. Here we show that while alternative partners may be ecologically different, their fitness effects on a shared mutualist can be effectively interchangeable and redundant. The results of our hierarchical models revealed that different ant partners exhibit different effects on vital rates, with *C. opuntiae*-tended plants experiencing advantages in growth and survival when small, and *L. apiculatum*-tended plants experiencing floral viability advantages. These results, alone, would suggest some potential benefits of partner diversity through complementarity. Yet, our stochastic, multi-state IPM revealed that all scenarios which included any partners resulted in similarly high cactus fitness, with a statistically significant but quantitatively weak cost of diversity at the highest level. Furthermore, evidence from nearly two decades of demographic data failed to support the “portfolio effect” hypothesis, whereby benefits of diversity manifest in the context of environmental fluctuations, because fluctuations in the demographic effects of alternative partners are too synchronized. Overall, our results indicate that while ant visitation is strongly consequential for plant demography, partner composition is much less so, suggesting that the fitness benefits of mutualism from the plant’s perspective may be surprisingly resilient to changes in partner diversity.

We attribute weak effects of partner identity and diversity to the vital rate sensitivity structure of this population. Like other long-lived, iteroparous species (Franco and Silvertown, 2004), tree cholla fitness is most sensitive to the growth and survival of established individuals (Elderd and Miller, 2016; Miller et al., 2009), which are virtually guaranteed many years of reproductive opportunities once they reach a size that is protected from mortality. Differences between alternative partners were most pronounced either in reproductive rates (floral viability and seed number), which contribute relatively weakly to fitness because a long reproductive lifespan overrides individual reproductive bouts, or in growth and survival at small sizes, where mortality risk is relatively high. At intermediate sizes, ant-tended plants had consistent growth and survival advantages over vacant plants regardless of partner identity, and that result dominates our integrative measures of

fitness; differences between partner species in other vital rates and at other sizes do not register nearly as strongly. The weak reduction in fitness at the highest partner diversity level is harder to explain, but suggests some form of opportunity cost in a more crowded partner environment.

Our results broaden the literature on multi-species mutualisms, a majority of which has demonstrated positive effects of partner diversity through complementarity (Afkhami et al., 2021; Afkhami and Stinchcombe, 2016; Fehling, 2022; Gustafson and Casper, 2005; Hernandez et al., 2019; Larimer et al., 2014; McKeon et al., 2012; Palmer et al., 2010; Stachowicz and Whitlatch, 2005; Yang et al., 2024) or portfolio effect (Lázaro et al., 2022; Rogers et al., 2014; Stevens et al., 2024; Thibaut et al., 2012). For example, working in another sequentially-partnered ant protective mutualism of long-lived plants (*Acacia drepanolobium*), Palmer et al. 2010 showed that synergies between alternative partners elevate plant fitness above what could be expected from the single best partner. However, the *Acacia* partner guild includes a “castrating” parasite that is uniquely able to boost plant growth by suppressing reproduction. We speculate that partners that differ in degree but not type, such as those in our system, may be less likely to synergize. Furthermore, our simulations exploring the absence of competitive dominance (Appendix ??) suggest that the turnover dynamics of the partner guild dictate which diversity mechanisms can manifest. Specifically, a sampling effect emerged only when we removed the competitive hierarchy that biases transitions toward *L. apiculatum*, allowing this species to dominate the cactus population. This suggests that when a high-quality partner is also high-frequency due to competitive dominance, there may be little benefit of adding more species, as the host is already likely to be occupied by the best available partner. Fewer studies have shown costs of mutualist diversity. Bruna et al. 2014 showed that ant diversity can depress fitness of the sequentially-partnered ant-plant *Maieta guianensis* due to reduced interactions with the better defender. Costs of diversity have also been documented in systems in which multiple partners interact simultaneously with a shared mutualist, likely due to interference via the host resource budget (Keller et al., 2018) or negative higher-order interactions (Barrett et al., 2015). One previous study (in a different ant-cactus system) showed neutral effects of partner diversity, but this study also showed that ant visitation was itself neutral, with no detectable effects on plant

fitness (Ford et al., 2015). To our knowledge, ours is the first study documenting a strong, positive fitness effect of mutualism, regardless of mutualist identity or guild composition. As this field grows, particularly with long-term studies that integrate multiple vital rates across the life cycle, we will be better positioned to understand whether positive, neutral, or negative effects of partner diversity predominate, and the conditions under which each might be more likely.

When partners exhibit different reactions to varying environments, interacting with a diverse portfolio of partners can lead to more consistent benefits across time (Batstone et al., 2018). Our work explicitly incorporated temporal environmental stochasticity, yet we find no evidence for the portfolio effect as a mechanism of diversity benefits. In ant-plant defensive mutualisms, any portfolio effect would emerge from fluctuations in the community of herbivores, and ant-specific responses to each herbivore species. Instead, we found that inter-annual fluctuations in vital rates were generally positively correlated across plants in different ant states, suggesting other drivers of inter-annual variation that are shared across the population (notably, weather) override any fluctuations in herbivore composition and ant-herbivore interactions. To further explore the potential for portfolio effects, we conducted simulations in which inter-annual fluctuations in ant effects were manipulated to be independent or negatively correlated; we found evidence of the portfolio effect in both scenarios, but especially when ant effects were negatively correlated (Appendix ??). These results indicate that while the portfolio effect can emerge under hypothetical conditions of anti-synchronous partner responses, shared drivers of inter-annual variability in our study system dampen this potential benefit of partner diversity. Dallas et al. 2022 found that while the portfolio effect was easy to show in theoretical models, it is infrequently detected in empirical data across many systems. This indicates that the portfolio effect may be difficult to detect, disguised by other mechanisms, or uncommon in nature.

This study highlights the value of long-term data in investigating species interactions. Previously studies have analyzed how partner identity and partner turnover impact focal mutualist fitness (Barrett et al., 2015; Bruna et al., 2014; Dejean et al., 2008; Fonseca and Benson, 2003; Noe and Hammerstein, 1994; Trojelsgaard et al., 2015). Separate studies have analyzed how inter-annual

variability impacts focal mutualists (Alarcón et al., 2008; Alonso, 1998; Horvitz and Schemske, 1990; Lázaro et al., 2022; Ollerton et al., 2006). The long term data set we used gave us the ability to consider, for the first time, the combined effects of partner identity, partner turnover, and temporal stochasticity. By piecing together complete life cycle information from long-term data, we gain a more nuanced understanding of the fitness consequences of ant visitation. For example, our previous study that focused only on reproductive vital rates suggested that *C. opuntiae* has overall parasitic fitness effects because activity of this species within tree cholla flowers can deter pollinators and reduce seed set (Figure ??) (Ohm and Miller, 2014). Yet, the more complete analysis presented here, which accounts for reduced seed set alongside other demographic advantages in higher-sensitivity vital rates, indicates that this species is clearly a mutualist.

As with any study, there are limitations to consider when interpreting our results. First, we used observational data to infer ant effects on plant demography. However, we have previously conducted experimental manipulations which revealed that ant presence has causal effects on plant demographic rates through anti-herbivore defense (Miller, 2007; Ohm and Miller, 2014). The combination of long-term observational data backed up by experimental results gives us greater confidence in our causal interpretations. Second, we have not explicitly incorporated ant-herbivore interactions, even though these are the primary pathway through which ants influence plant demography. Surveys of herbivore damage from our long-term data (Figure ??) support the protective benefits of ant visitation. Specifically, herbivore pressure was highest on plants in which no ant was present (vacancy), intermediate under Other and *C. opuntiae* tending, and lowest under *L. apiculatum* tending. This gradient aligns with the expected defensive hierarchy of ant species, providing indirect evidence that differences among ant partners influence herbivory rates and, by extension, plant demographic outcomes (Miller, 2007; Ohm and Miller, 2014). Future work explicitly incorporating ant-herbivore-plant interactions in the demographic models could more directly quantify how variation in ant species composition mediates herbivore pressure and plant fitness. Third, the rarity of several ant partner species imposed a practical constraint on assessing effects of partner diversity, since we needed to combine these into an Other category that might mask important differences.

As more long-term data accumulate, we may gain additional information to further unpack this group of species.

Conclusion. This study highlights that while partners within a mutualistic guild can be ecologically different, they may still be interchangeable in their net effects on fitness of a shared mutualist based on where their effects occur in the life cycle, highlighting the importance of a demographic perspective in understanding the fitness consequences of mutualisms. Contrary to much of the literature on multi-species mutualisms, we find that while ant partners are beneficial, partner diversity has no measurable benefit to their shared mutualist and may even be somewhat costly. As studies accumulate across different types of multi-partner mutualisms, particularly those spanning simultaneous/sequential partnerships and leveraging demographic data from the complete life cycle, our field will be better poised to understand whether, how, and under what conditions mutualist diversity is neutral, costly, or beneficial.

Acknowledgments

Financial support for this work came from the Sevilleta Long-Term Ecological Research Program (NSF DEB awards 1655499, 1748133, and 2425290). We acknowledge the many students and technicians who have helped collect annual census data for this project, including M. Donald, J. Fowler, M. Saucedo, K. Dickens, T. Jordan-Millet, J. Moutouama, C. Oxley, K. Schraeder, B. Scherick, A. Sears, M. Tucker, and J. Xiong. J. Moutouama, A. Bradley, B. Scherick, A. Gough, J. Martin, and V. Rudolf provided helpful feedback on the manuscript. We also thank three reviewers for thoughtful comments and suggestions on earlier versions of this manuscript.

Author Contribution Statement

A.M.C. and T.E.X.M. conceptualized the project and designed the analytical methods. T.E.X.M. initiated the long-term study. A.M.C. wrote the simulation code, analyzed and visualized data, an-

alyzed the models, and wrote the original draft. A.M.C. and T.E.X.M. obtained funding, collected data, and participated in review and editing of the manuscript.

Data and Code Availability

Most of the data that support the findings of this study (through 2019) are openly available as a data package (Miller, 2020) on the Environmental Data Initiative website, package id: knb-lter-sev.323.1. An updated data package, including recent years, is presently under review and we expect will be available in early 2026. For now, all of our data and code are available in our public zenodo repo: <https://zenodo.org/records/20030584>.

Literature Cited

- Afkhami, M. E., Friesen, M. L., and Stinchcombe, J. R. (2021). Multiple mutualism effects generate synergistic selection and strengthen fitness alignment in the interaction between legumes, rhizobia and mycorrhizal fungi. *Ecology Letters*, 24(9):1824–1834.
- Afkhami, M. E., Rudgers, J. A., and Stachowicz, J. J. (2014). Multiple mutualist effects: conflict and synergy in multispecies mutualisms. *Ecology*, 95(4):833–844.
- Afkhami, M. E. and Stinchcombe, J. R. (2016). Multiple mutualist effects on genomewide expression in the tripartite association between *Medicago truncatula*, nitrogen-fixing bacteria and mycorrhizal fungi. *Molecular Ecology*, 25:4946–4962.
- Agrawal, A. A. and Rutter, M. T. (1998). Dynamic anti-herbivore defense in ant-plants: the role of induced responses. *Oikos*, 83(2):227–236.
- Alarcón, R., Waser, N. M., and Ollerton, J. (2008). Year-to-year variation in the topology of a plant-pollinator interaction network. *Oikos*, 117(11):1796–1807.

- Alonso, L. E. (1998). Spatial and temporal variation in the ant occupants of a facultative ant-plant. *Biotropica*, 30:201–213.
- Barrett, L. G., Bever, J. D., Bissett, A., and Thrall, P. H. (2015). Partner diversity and identity impacts on plant productivity in *Acacia*-rhizobial interactions. *Journal of Ecology*, 103(1):130–142.
- Barton, K. E. and Koricheva, J. (2010). The ontogeny of plant defense and herbivory: characterizing general patterns using meta-analysis. *The American Naturalist*, 175(4):481–493.
- Bascompte, J. (2009). Disentangling the web of life. *Science*, 325(5939):414–416.
- Batstone, R. T., Carscadden, K. A., Afkhami, M. E., and Frederickson, M. E. (2018). Using niche breadth theory to explain generalization in mutualisms. *Ecology*, 99(5):1039–1050.
- Beattie, A. (1985). *The Evolutionary Ecology of Ant-Plant Mutualisms*. Cambridge University Press, Cambridge, UK.
- Benson, L. (1982). *Cacti of the United States and Canada*. Stanford University Press, Stanford, CA.
- Boege, K. and Marquis, R. J. (2005). Facing herbivory as you grow up: The ontogeny of resistance in plants. *Trends in Ecology and Evolution*, 20(7):441–448.
- Bronstein, J. L. (1998). The contribution of ant-plant protection studies to our understanding of mutualism. *BioTropica*, 30(2):150–161.
- Bronstein, J. L., Alarcón, R., and Geber, M. (2006). The evolution of plant–insect mutualisms. *New Phytologist*, 172(1):412–428.
- Bruna, E. M., Izzo, T. J., Inouye, B. D., and Vasconcelos, H. L. (2014). Effect of mutualist partner identity on plant demography. *Ecology*, 95(12):3237–3243.

- Byk, J. and Del-Claro, K. (2011). Ant-plant interaction in the Neotropical savanna: direct beneficial effects of extrafloral nectar on ant colony fitness. *Population Ecology*, 53(3):327–332.
- Báez, S., Donoso, D. A., Queenborough, S. A., Jaramillo, L., Valencia, R., and Dangles, O. (2016). Ant mutualism increases long-term growth and survival of a common Amazonian tree. *The American Naturalist*, 188(5):1–9.
- Dallas, T. A. and Kramer, A. M. (2022). Temporal variability in population and community dynamics. *Ecology*, 103(2):e03577.
- Dattilo, W., Marquitti, F. M. D., Guimaraes, P. R., and Izzo, T. J. (2014). The structure of ant-plant ecological networks: Is abundance enough? *Ecology*, 95(2):475–485.
- Dejean, A., Djieto-Lordon, C., Cereghino, R., and Leponce, M. (2008). Ontogenetic succession and the ant mosaic: an empirical approach using pioneer trees. *Basic and Applied Ecology*, 9(3):316–323.
- Djiéto-Lordon, C., Dejean, A., Ring, R. A., Nkongmeneck, B. A., Lauga, J., and McKey, D. (2005). Ecology of an improbable association: The pseudomyrmecine plant-ant *Tetraponera tessmanni* and the myrmecophytic liana *Vitex thyrsoiflora* (lamiaceae) in Cameroon. *Biotropica*, 37(3):421–430.
- Donald, M. L. and Miller, T. E. X. (2022). Does ant-plant mutualism have spillover effects on the non-partner ant community? *Ecology and Evolution*, 12(1):1–12.
- Elder, B. D. and Miller, T. E. X. (2016). Quantifying demographic uncertainty: Bayesian methods for integral projection models. *Ecological Monographs*, 86(1):125–144.
- Ellner, S. P., Childs, D. Z., and Rees, M. (2016). *Data-driven Modeling of Structured Populations: A Practical Guide to the Integral Projection Model*. Springer, Ithaca, NY.
- Fehling, L. S. (2022). Reward complementarity and context dependency in multispecies mutualist

- interactions in partridge pea (*Chamaecrista fasciculata*). Master's thesis, Miami University, Miami, FL.
- Fiala, B., Grunsky, H., Maschwitz, U., and Linsenmair, K. E. (1994). Diversity of ant-plant interactions: protective efficacy in macaranga species with different degrees of ant association. *Oecologia*, 97(2):186–192.
- Fonseca, C. R. and Benson, W. W. (2003). Ontogenetic succession in Amazonian ant trees. *Oikos*, 102(2):407–412.
- Ford, K. R., Ness, J. H., Bronstein, J. L., and Morris, W. F. (2015). The demographic consequences of mutualism: ants increase host-plant fruit production but not population growth. *Oecologia*, 179:435–446.
- Franco, M. and Silvertown, J. (2004). A comparative demography of plants based upon elasticities of vital rates. *Ecology*, 85(2):531–538.
- Fraser, A. M., Axen, A. H., and Pierce, N. E. (2001). Assessing the quality of different ant species as partners of a myrmecophilous butterfly. *Oecologia*, 129(4):452–460.
- Frederickson, M. E. (2005). Ant species confer different partner benefits on two Neotropical myrmecophytes. *Oecologia*, 143(3):387–395.
- Frederickson, M. E. (2013). Rethinking mutualism stability: Cheaters and the evolution of sanctions. *Quarterly Review of Biology*, 88(4):269–295.
- Gaume, L., Mckey, D., and Terrin, S. (1998). Ant-plant-homopteran mutualism: how the third partner affects the interaction between a plant-specialist ant and its myrmecophyte host. *Ecological Entomology*, 26:569–575.
- Gustafson, D. and Casper, B. (2005). Differential host plant performance as a function of soil arbuscular mycorrhizal fungal communities: experimentally manipulating co-occurring glomus species. *Plant Ecology*, 183:257–263.

- Hautier, Y., Isbell, F., Borer, E. T., Seabloom, E. W., Harpole, W. S., Lind, E. M., MacDougall, A. S., Stevens, C. J., and Tilman, D. (2014). Eutrophication weakens stabilizing effects of diversity in natural grasslands. *Nature*, 508:521–525.
- Hernandez, D., Kieseewetter, K., Palakurty, S., Stinchcombe, J., and Afkhami, M. (2019). Synergism and symbioses: unpacking complex mutualistic species interactions using transcriptomic approaches. In de Bruijn, F. J., editor, *The Model Legume Medicago truncatula*, pages 1045–1054. John Wiley & Sons, Hoboken, New Jersey, USA.
- Horvitz, C. C. and Schemske, D. W. (1986). Seed dispersal of a Neotropical myrmecochore: variation in removal rates and dispersal distance. *bioTropica*, 18(4):319–323.
- Horvitz, C. C. and Schemske, D. W. (1990). Spatiotemporal variation in insect mutualists of a Neotropical herb. *Ecology*, 71:1085–1097.
- Keller, K. R., Carabajal, S., Navarro, F., and Lau, J. A. (2018). Effects of multiple mutualists on plants and their associated arthropod communities. *Oecologia*, 186:185–194.
- Larimer, A. L., Clay, K., and Bever, J. D. (2014). Synergism and context dependency of interactions between arbuscular mycorrhizal fungi and rhizobia with a prairie legume. *Ecology*, 95:1045–1054.
- Loreau, M. (1999). Biodiversity and ecosystem functioning: Effects of species diversity on productivity and stability in plant communities. *Proceedings of the Royal Society B: Biological Sciences*, 264(1381):127–130.
- Ludka, J., Levan, K. E., and Holway, D. A. (2015). Infiltration of a facultative ant–plant mutualism by the introduced Argentine ant: effects on mutualist diversity and mutualism benefits. *Ecological Entomology*, 40(4):437–443.
- Lázaro, A., Gómez-Martínez, C., González-Estévez, M. A., and Hidalgo, M. (2022). Portfolio effect and asynchrony as drivers of stability in plant–pollinator communities along a gradient of landscape heterogeneity. *Ecography*, 45(3):1–14.

- Mann, J. (1969). *Cactus-feeding insects and mites*. United States National Museum Bulletin. Smithsonian Institution Press, Washington, D.C.
- McKeon, C. S., Stier, A. C., McIlroy, S. E., and Bolker, B. M. (2012). Multiple defender effects: synergistic coral defense by mutualist crustaceans. *Oecologia*, 169(4):1095–1103.
- Metcalf, C. J. E., Ellner, S. P., Childs, D. Z., Salguero-Gómez, R., Merow, C., McMahon, S. M., Jongejans, E., and Rees, M. (2015). Statistical modelling of annual variation for inference on stochastic population dynamics using integral projection models. *Methods in Ecology and Evolution*, 6(9):1007–1017.
- Miller, T. (2020). Long-term study of tree cholla demography in the Los Piños mountains, Sevilleta National Wildlife Refuge.
- Miller, T. E. and Ellner, S. P. (2025). My, how you’ve grown: A practical guide to modeling size transitions for integral projection model (ipm) applications. *Ecology*, 106(5):e70088.
- Miller, T. E., Louda, S. M., Rose, K. A., and Eckberg, J. O. (2009). Impacts of insect herbivory on cactus population dynamics: experimental demography across an environmental gradient. *Ecological Monographs*, 79(1):155–172.
- Miller, T. E. X. (2007). Does having multiple partners weaken the benefits of facultative mutualism? a test with cacti and cactus-tending ants. *Oikos*, 116(3):500–512.
- Miller, T. E. X. (2014). Plant size and reproductive state affect the quantity and quality of rewards to animal mutualists. *Journal of Ecology*, 102(4):496–507.
- Miller, T. E. X., Tyre, A. J., and Louda, S. M. (2006). Plant reproductive allocation predicts herbivore dynamics across spatial and temporal scales. *The American Naturalist*, 168(5):608–616.
- Moraes, S. C. and Vasconcelos, H. L. (2009). Long-term persistence of a neotropical ant-plant population in the absence of obligate plant-ants. *Ecology*, 90(8):2375–2383.

- Nakazawa, T. (2020). Ontogenetic shifts in mutualistic interactions: stage-structured models of host–symbiont dynamics. *Oikos*, 129(6):857–869.
- Ness, J. H., Morris, W. F., and Bronstein, J. L. (2006). Integrating quality and quantity of mutualistic service to contrast ant species protecting *Ferocactus wislizeni*. *Ecology*, 87(4):912–921.
- Ness, J. H., Morris, W. F., and Bronstein, J. L. (2009). For ant-protected plants, the best defense is a hungry offense. *Ecology*, 90(10):2823–2831.
- Noe, R. and Hammerstein, P. (1994). Biological markets: supply and demand determine the effect of partner choice in cooperation, mutualism and mating. *Behavioral Ecology and Sociobiology*, 35(1):1–11.
- Ohm, J. R. and Miller, T. E. X. (2014). Balancing anti-herbivore benefits and anti-pollinator costs of defensive mutualists. *Ecology*, 95(10):2924–2935.
- Oliveira, P. S., Rico-Gray, V., Díaz-Castelazo, C., and Castillo-Guevara, C. (1999). Interaction between ants, extrafloral nectaries and insect herbivores in neotropical coastal sand dunes: herbivore deterrence by visiting ants increases fruit set in *Opuntia stricta* (cactaceae). *Functional Ecology*, 13(5):623–631.
- Ollerton, J., Johnson, S. D., and Hingston, A. B. (2006). Geographical variation in diversity and specificity of pollination systems. In Waser, N. M. and Ollerton, J., editors, *Plant-Pollinator Interactions: From Specialization to Generalization*, pages 283–308. University of Chicago Press, Chicago, IL.
- Palmer, T. M., Doak, D. F., Stanton, M. L., Bronstein, J. L., Kiers, T. E., Young, T. P., Goheen, J. R., and Pringle, R. M. (2010). Synergy of multiple partners, including freeloaders, increases host fitness in a multispecies mutualism. *Proceedings of the National Academy of Sciences*, 107(40):17234–17239.

- R Core Team (2023). *R: A Language and Environment for Statistical Computing*. R Foundation for Statistical Computing, Vienna, Austria. Version 4.3.3.
- Rees, M. and Ellner, S. P. (2009). Integral projection models for populations in temporally varying environments. *Ecological Monographs*, 79(4):575–594.
- Rogers, S. R., Tarpy, D. R., and Burrack, H. J. (2014). Bee species diversity enhances productivity and stability in a perennial crop. *PloS One*, 9(5):e97307.
- Rudgers, J. A. (2004). Enemies of herbivores can shape plant traits: selection in a facultative ant-plant mutualism. *Ecology*, 85(1):192–205.
- Sachs, J. L., Mueller, U. G., Wilcox, T. P., and Bull, J. J. (2004). The evolution of cooperation. *Quarterly Review of Biology*, 79(2):135–160.
- Stachowicz, J. J. and Whitlatch, R. B. (2005). Multiple mutualists provide complementary benefits to their seaweed host. *Ecology*, 86(9):2418–2427.
- Stan Development Team (2023). RStan: the R interface to Stan. R package version 2.32.3.
- Stanton, M. L. (2003). Interacting guilds: moving beyond the pairwise perspective on mutualisms. *The American Naturalist*, 162:S10–S23.
- Stevens, B. L. F., Howard, K., Bogar, L. M., et al. (2024). Environmental fluctuations promote host reward strategies that maintain partner diversity in multispecies mutualisms. *The American Naturalist*, 202(6):E000.
- Thibaut, L. M., Connolly, S. R., and Sweatman, H. P. (2012). Diversity and stability of herbivorous fishes on coral reefs. *Ecology*, 93(4):891–901.
- Tilman, D., Wedin, D., and Knops, J. (1994). Biodiversity and stability in grasslands. *Nature*, 367:363–365.

- Trager, M. D., Bhotika, S., Hostetler, J. A., Andrade, G. V., Rodriguez-Cabal, M. A., Mckeen, C. S., Osenberg, C. W., and Bolker, B. M. (2010). Benefits for plants in ant-plant protective mutualisms: a meta-analysis. *PLoS ONE*, 5(12):e14308.
- Trojelsgaard, K., Jordano, P., Carstensen, D. W., and Olesen, J. M. (2015). Geographical variation in mutualistic networks: Similarity, turnover and partner fidelity. *Proceedings of the Royal Society B: Biological Sciences*, 282(18-2):1–9.
- Ushio, M. (2020). Interaction capacity as a potential driver of community diversity. *Proceedings of the Royal Society B*, 289(1969):20212690.
- Ventre-Lespiauq, A., Jacquemyn, H., Ramirez, S., and Bidartondo, M. I. (2021). Temporal dynamics of orchid–mycorrhizal interactions: turnover and persistence of fungal symbionts across life stages. *New Phytologist*, 230(2):697–708.
- Wetzel, W. C., Inouye, B. D., Hahn, P. G., Whitehead, S. R., and Underwood, N. (2023). Variability in plant-herbivore interactions. *Annual Review of Ecology, Evolution, and Systematics*, 54:451–474.
- Williams, J. L., Miller, T. E. X., and Ellner, S. P. (2012). Avoiding unintentional eviction from integral projection models. *Ecology*, 93(9):2008–2014.
- Winfree, R. (2020). How does biodiversity relate to ecosystem functioning in natural ecosystems? In Dobson, A., Holt, R. D., and Tilman, D., editors, *Unsolved Problems in Ecology*, pages 302–318. Princeton University Press, Princeton, NJ.
- Yang, N., Røder, H. L., Wicaksono, W. A., Wassermann, B., Russel, J., Li, X., Nesme, J., Berg, G., Sørensen, S. J., and Burmølle, M. (2024). Interspecific interactions facilitate keystone species in a multispecies biofilm that promotes plant growth. *International Society for Microbial Ecology Journal*, 18(1):wrae012.

Figure Legends

Figure 1 The next predicted size of tree cholla based on previous size in relation to ant partner (a, *C. opuntiae*; b, *L. apiculatum*; c, Other; d, Vacant, e, all states combined). Points show observed size transitions, colored lines are the mean next size, and contours show probability density of the skewed normal growth model. Dashed lines indicate extrapolation while solid lines indicate the range of observed data. They gray dashed line indicates no change in size (areas above this line show growth while areas under it show shrinkage).

alt. text: A multipaneled figure. The first four panels are line plots wit 95% confidence intervals overlaid. Each panel shows the probability of growing from one size to the next based on ant partner identity. The fifth panel shows all four of these mean lines overlaid.

Figure 2 Probability of survival in relation to size and ant partner. Layout as in Fig. 1, except here points show data as binned means, where point size is proportional to number of observations. Grey areas show the 90% confidence interval for the mean.

alt. text: A multipaneled figure. The first four panels are line plots wit 95% confidence intervals overlaid. Each panel shows the probability of surviving from one year to the next based on size and ant partner identity. The fifth panel shows all four of these mean lines overlaid.

Figure 3 Flowerbud viability in relation to ant partner status. Points show observed data (fraction of initiated buds that were viable). Points and bars show the means and 95% credible intervals for each group.

alt. text: A single paneled figure with the mean estimated flowers produced by size. The 95% confidence interval and real data are overlaid.

Figure 4 The correlation coefficients (Pearson's r) of ant-specific year-random effects from statistical models for growth (a), survival (b), and floral viability (c). Dark red indicates negative correlation, dark blue indicates positive correlation, white indicates no correlation. The year-specific random effects for each ant state from the statistical models are shown in a time series

for growth (d), survival (e), and floral viability (f). All random effects have mean zero such that positive values indicate better-than-average years and negative values indicate worse-than-average years. Values are missing where there were no data available for that year.

alt. text: A multipaneled figure. The top 3 panels show correlation plots with color indicating the correlation of ant-year effects. The bottom three plots show the ant random effects across time. These bottom three are line plots.

Figure 5 The probability of ant state based on plant size and previous ant state. Each panel shows size-dependent probabilities of the next ant state conditional on the previous ant state and size. Solid lines represent predictions of the multinomial statistical model and points show data binned over size intervals, where point size is proportional to the number of observations.

alt. text: A multipaneled figure. Each panel is a series of lines representing the probability of being visited by each ant species or vacant with the data overlaid. Each panel represents a previous ant state.

Figure 6 Tree cholla stochastic growth rate (λ_S) corresponding to number (a) or number and identity (b-c) of ant partners for synchronous and non-synchronous inter-annual fluctuations. Points and intervals show mean and 95% credible intervals from the posterior distributions of λ_S . In b-c letters in the legend correspond to ant partners as follows (V = Vacant, C = *C. opuntiae*, L = *L. apiculatum*, O = other).

alt. text: A multipaneled figure. Each panel shows a series of means as points with 95% confidence intervals represented as lines extending.

Figure A1 The posterior distributions of seeds produced per fruit on reproducing plants. These estimates are ant specific with the yellow representing vacant plant seed production, purple representing *C. opuntiae* tended plant seed production, and teal representing *L. apiculatum* tended plant seed production.

Figure A2 The probability of a cholla plant reproducing based on its size. The green line

shows the average probability, the gray surrounding area shows the 90% confidence interval, and the black points show the actual data.

Figure A3 The number of flowers produced by a cholla plant based on its size. The green line shows the average number of flowers produced, the gray surrounding area shows the 90% confidence interval, and the green points show the binned data.

Figure B4 The proportion of reproducing plants which are visited by herbivores. Each bar represents the observed proportion of cacti in each ant state with observed herbivores.

Figure C5 Panel a) the mean estimated λ_{NS} and λ_S for different numbers of partners (0-3) for the synchronous IPM (black circle) and the non-synchronous IPM (red diamond). The lines show the posterior distribution spread of the estimated λ values. Panels b-c) the mean estimated λ_{NS} and λ_S respectively, for each simulated combination of ant partner as the filled in circles. The lines show the posterior distribution spread of estimated λ values. The letters in the legend correspond to what ant partners are present (V = Vacant, C = *C. opuntiae*, L = *L. apiculatum*, O = other).

Figure C6 Panel a) the mean estimated λ_{NS} and λ_S for different numbers of partners (0-3) for the synchronous IPM (black circle) and the non-synchronous IPM (red diamond). The lines show the posterior distribution spread of the estimated λ values. Panels b-c) the mean estimated λ_{NS} and λ_S respectively, for each simulated combination of ant partner as the filled in circles. The lines show the posterior distribution spread of estimated λ values. The letters in the legend correspond to what ant partners are present (V = Vacant, C = *C. opuntiae*, L = *L. apiculatum*, O = other).

Figure D7 The distributions of $\lambda_S^4 - \lambda_S^0$ for the non-synchronous and synchronous models are shown in pink and green respectively. The vertical dashed line shows where the effect of partners on the fitness of the cholla is 0 (to the left the partners have a negative effect, to the right the partners are beneficial).

Figure D8 A comparison of synchronous and non-synchronous cactus responses to partner diversity across inter-annual fluctuations. If the non-synchronous effects of partner diversity are

greater than the synchronous ones under fluctuation environments, evidence of the portfolio effect is found. Panel a) shows the mean effects of partner diversity ($\lambda_S^4 - \lambda_S^0$) under realistic levels of non-synchronicity (what was estimated from the field data), while panels b) and c) shows the mean effects of partner diversity under no synchronicity (where we artificially forced a correlation of 0 for the ant-specific year random effects) and negative synchronicity (where we artificially forced a correlation close to -1 for the ant-specific year random effects). The mean estimations of $\lambda_S^4 - \lambda_S^0$ for the non-synchronous and synchronous models are shown in green and pink respectively.