

Functions, Vectors, Operators, Matrices, and Green Functions

Andrew Carnes

October 11, 2018

Contents

1	Introduction	3
2	Functions As Vectors	3

1 Introduction

This document looks at functions as vectors and reinterprets derivatives, Green functions, and the Dirac delta function as matrices. We also look at dot products, integrals, and transformations (like the Fourier transform) in this formalism. We find that the Green function for a differential operator is the inverse matrix of that differential operator. Looking at Green functions this way really helped me with electromagnetism and QFT in graduate school. Plus it's pretty cool. Nothing here is super rigorous, but it gets the point across and provides a new way to look at things.

2 Functions As Vectors

What is a vector? A vector \vec{v} is an ordered 1D storage device. A vector of length N has N bins to put numbers in, and you keep track of the bins by labeling them with an integer. Note that you can only put one number in each bin. If we label the bins with i, v_i is the information in the i^{th} bin. \vec{v} usually written like so,

$$\vec{v} = \begin{pmatrix} v_1 \\ v_2 \\ \dots \\ v_N \end{pmatrix}. \quad (1)$$

Now consider a function. A function $f(x)$ is an ordered 1D storage device with a bunch of bins. The bins are labeled by x, and $f(x)$ is the number in the x^{th} bin. Each bin only has one number. In this case, x is a real number so $f(x)$ is much more dense than \vec{v} . $f(x)$ can be written like so,

$$f(x) = \begin{pmatrix} f(-\infty) \\ \dots \\ f(-3.1111111128) \\ \dots \\ f(-3.111111112) \\ \dots \\ f(0) \\ \dots \\ f(912.99) \\ \dots \\ f(999999999) \\ \dots \\ f(\infty) \end{pmatrix}. \quad (2)$$

Between any two bins there are an infinite number of bins (welcome to the real numbers), which makes things tricky, but aside from the storage density, functions and vectors work the same way. Give them a location and they spit out the info you want.

Let's discretize the real number line into chunks of width dx. We can then label the

entries in $f(x)$ by $f(x_i)$, where $x_{i+1} - x_i = dx$. Then we have

$$f(x) = \begin{pmatrix} f(x_{-\infty}) \\ \dots \\ \dots \\ f(x_{-2}) \\ \dots \\ f(x_{-1}) \\ \dots \\ f(x_0) \\ \dots \\ f(x_1) \\ \dots \\ f(x_2) \\ \dots \\ \dots \\ f(x_{\infty}) \end{pmatrix}. \quad (3)$$

When dx is infinitely small we can approximate all $f(x)$ between $f(x_i)$ and $f(x_{i+1})$ by $f(x_i)$. Now let's take a dot product. With two vectors, \vec{v} and \vec{u} , the dot product is

$$\vec{v} \cdot \vec{u} = v_1 u_1 + v_2 u_2 + \dots + v_N u_N = \sum_{i=1}^N v_i u_i. \quad (4)$$

And as usual, when $\vec{v} \cdot \vec{u} = 0$ the vectors are orthogonal and when $\frac{\vec{v} \cdot \vec{u}}{\sqrt{v^2 u^2}} = 1$ the vectors are parallel. When all v_i and u_i are 1, $\vec{v} \cdot \vec{u} = N$, the number of components. If we consider only a subset of components with all $v_i = u_i = 1$ then $\sum_{i=1}^{N_c} v_i u_i = N_c$, the number of components summed over.

Now take a dot product of functions $f(x)$ and $g(x)$,

$$f(x) \cdot g(x) = \dots + f(x_{-2})g(x_{-2}) + \dots + f(x_{-1})g(x_{-1}) + \dots + f(x_0)g(x_0) + \dots + f(x_1)g(x_1) + \dots + f(x_2)g(x_2) + \dots \quad (5)$$

Let's look at the sum from x_1 to x_2 when $f(x) = g(x) = 1$ in all bins

$$f(x) \cdot g(x) = f(x_1)g(x_1) + \dots + \dots \text{up to } f(x_2)g(x_2) = 1 + \dots + 1 = N_c. \quad (6)$$

This gives us the number of components summed over, N_c , which is infinite. That's true, but it's not a helpful normalization. Let's redefine the dot product between functions to yield the size of the real interval summed over, $x_2 - x_1 = dx$.

$$f(x) \cdot g(x) = f(x_1)g(x_1)\epsilon + \dots + \dots \text{up to } f(x_2)g(x_2) = 1\epsilon + \dots + 1\epsilon = N_c\epsilon = dx. \quad (7)$$

The full dot product is now

$$f(x) \cdot g(x) = \dots + f(x_{-2})g(x_{-2})\epsilon + \dots + f(x_{-1})g(x_{-1})\epsilon + \dots + f(x_0)g(x_0)\epsilon + \dots + f(x_1)g(x_1)\epsilon + \dots + f(x_2)g(x_2)\epsilon + \dots \quad (8)$$

Finally, let's represent all N_c terms between $f(x_i)g(x_i)$ and $f(x_{i+1})g(x_{i+1})$ by $f(x_i)g(x_i)$ using the assumption that $f(x)$ and $g(x)$ change very little throughout the interval dx . This yields

$$f(x) \cdot g(x) = \dots + f(x_{-2})g(x_{-2})N_c\epsilon + f(x_{-1})g(x_{-1})N_c\epsilon + f(x_0)g(x_0)N_c\epsilon + f(x_1)g(x_1)N_c\epsilon + f(x_2)g(x_2)N_c\epsilon + \dots =$$

(9)