

A SEARCH FOR THE STANDARD MODEL HIGGS DECAYING TO TWO MUONS AT THE  
CMS EXPERIMENT

By  
ANDREW CARNES

A DISSERTATION PRESENTED TO THE GRADUATE SCHOOL  
OF THE UNIVERSITY OF FLORIDA IN PARTIAL FULFILLMENT  
OF THE REQUIREMENTS FOR THE DEGREE OF  
DOCTOR OF PHILOSOPHY

UNIVERSITY OF FLORIDA

2017

© 2017 Andrew Carnes

## ACKNOWLEDGMENTS

I would like to thank my Mom and Dad for just being all around great parents. Without all of their support, the completion of this Ph.D would never have been possible. I would also like to thank Professors Paul Avery and Darin Acosta for their help. In addition, I would like to thank Andrew Brinkerhoff, Pierluigi Bortignon, and Andrea Marini for all of their support and hard work on the project. I'd also like to thank my friends Stany, Leo, Sean, Sophie, and David for making Geneva a great place to live.

# TABLE OF CONTENTS

	<u>page</u>
ACKNOWLEDGMENTS . . . . .	3
LIST OF TABLES . . . . .	6
LIST OF FIGURES . . . . .	7
ABSTRACT . . . . .	8
CHAPTER	
1 INTRODUCTION . . . . .	9
2 THE LARGE HADRON COLLIDER AND THE CMS EXPERIMENT . . . . .	11
2.1 Large Hadron Collider . . . . .	11
2.2 Compact Muon Solenoid Detector . . . . .	13
2.2.1 Silicon Tracker . . . . .	14
2.2.2 Calorimeters . . . . .	15
2.2.3 Muon System . . . . .	17
2.2.3.1 Drift Tubes . . . . .	19
2.2.3.2 Cathode Strip Chambers . . . . .	19
2.2.3.3 Resistive Plate Chambers . . . . .	20
2.2.4 Trigger System . . . . .	20
2.2.4.1 L1 Trigger . . . . .	21
3 THE STANDARD MODEL . . . . .	23
3.1 Quantum Field Theory . . . . .	25
3.1.1 Particles, Symmetries, and Labels . . . . .	26
3.1.2 The Lagrangian Formalism . . . . .	27
3.1.3 QFT From Symmetry . . . . .	31
3.1.3.1 Rotations . . . . .	31
3.1.3.2 The Lorentz Group . . . . .	35
3.1.3.3 The Poincare Group and Particle Labels . . . . .	43
3.1.3.4 Building the Lagrangian for a Free Scalar Particle . . . . .	44
3.1.3.5 Building the Lagrangian for a Free Spin $\frac{1}{2}$ Particle . . . . .	46
3.1.3.6 Building the Lagrangian for a Free Spin 1 Particle . . . . .	52
3.1.4 Lagrangians in Quantum Mechanics and QFT . . . . .	53
3.1.5 The Higgs Mechanism . . . . .	55
3.1.6 Perturbation Theory and Feynman Rules . . . . .	55
3.2 The Standard Model Higgs . . . . .	55
3.2.1 SM Higgs Production and Decay Modes . . . . .	55

4	RESULTS . . . . .	59
4.1	Fusce Eget Tempus Lectus, . . . . .	59
5	SUMMARY AND CONCLUSIONS . . . . .	64
5.1	Non Porttitor Tellus . . . . .	64
5.1.1	Nam Arcu Magna . . . . .	64
5.1.1.1	Ut pellentesque velit sede . . . . .	64
APPENDIX		
A	THIS IS THE FIRST APPENDIX . . . . .	65
B	AN EXAMPLE OF A HALF TITLE PAGE . . . . .	66
C	DERIVATION OF THE $\Upsilon$ FUNCTION . . . . .	70
D	DERIVATION OF THE $\Upsilon$ FUNCTION . . . . .	74
REFERENCES . . . . .		78
BIOGRAPHICAL SKETCH . . . . .		79

## LIST OF TABLES

Table

page

## LIST OF FIGURES

<u>Figure</u>	<u>page</u>
2-1 The CERN Accelerator Complex ? . . . . .	12
2-2 The CMS detector ? . . . . .	13
2-3 The sagitta measurement . . . . .	15
2-4 A slice of the CMS detector ? . . . . .	16
2-5 A Look at the Muon System ? . . . . .	18
2-6 A Drift Tube ? . . . . .	19
2-7 A Cathode Strip Chamber ? . . . . .	20
2-8 The L1 Trigger Architecture ? . . . . .	22
3-1 The Standard Model Particles . . . . .	24
3-2 The Feynman diagram for two gluons fusing into a Higgs. There are three vertices and this is a third order diagram. Two vertices involve the strong force and one vertex involves the Higgs coupling. The matrix element for this diagram would have two factors of the strong force coupling and one factor for the Higgs coupling which involves the mass of the top quark. . . . .	55
3-3 The highest production mode cross sections for the SM Higgs ? . . . . .	56
3-4 The graphic on the top left presents the SM Higgs branching fractions as functions of mass while the table on the bottom right displays the branching fractions for a 125 GeV SM Higgs ? . . . . .	57
3-5 The SM production modes with the highest cross sections. a) Gluon Gluon Fusion (GF) b) Vector Boson Fusion (VBF) c) Associated Production with a Vector Boson (VH) d) $t\bar{t}H$ . . . . .	57
3-6 Quark gluon scattering creates many two jet events. This background looks very similar to GF when the Higgs decays to two jets. The colliding protons are made of quarks and gluons so this process is extremely common. . . . .	58
B-1 L <sup>A</sup> T <sub>E</sub> X2 <sub>ε</sub> . logo . . . . .	67

Abstract of Dissertation Presented to the Graduate School  
of the University of Florida in Partial Fulfillment of the  
Requirements for the Degree of Doctor of Philosophy

A SEARCH FOR THE STANDARD MODEL HIGGS DECAYING TO TWO MUONS AT THE  
CMS EXPERIMENT

By

Andrew Carnes

December 2017

Chair: Paul Avery

Major: Physics

In 2012 two collaborations at the Large Hadron Collider announced the discovery of a new particle with properties similar to the Standard Model Higgs Boson. In order to determine whether the boson discovered with a mass of 125 GeV is actually the Standard Model Higgs, all of the different ways the particle can decay need to be investigated. If the probabilities for the different decays do not match the predictions of the Standard Model then this would imply new physics.

This dissertation presents the search for the Standard Model Higgs Boson decaying to  $\mu^+\mu^-$ . The search uses the  $35.9 \pm 0.9 \text{ fb}^{-1}$  of  $\sqrt{s} = 13 \text{ TeV}$  proton-proton collision data recorded by the CMS detector in 2016. The observed and expected upper limits on the rate at a 95 % confidence level are presented for Higgs masses in the range 120 to 130 GeV. The expected and observed upper limits at a mass of 125 GeV are  $x.xx$  and  $1.98^{+0.81}_{-0.57} \times \text{SM}$  respectively. These results provide the best results to date on the Higgs coupling to second generation fermions. No deviations from the Standard Model are observed.



## CHAPTER 1 INTRODUCTION

The Standard Model of particle physics is an extremely successful theory shown to correctly predict the behavior of the particles and forces which make up the most basic constituents of the universe. In fact, it correctly describes all of the forces known except for gravity. All of the elementary particles detailed in the Standard Model have been discovered except for one, the Higgs boson. As explained by special relativity, potential energy can be realized as mass and it is the potential from the interaction with the nonzero Higgs field that bestows mass onto the massive elementary particles in the Standard Model. On July 4, 2012 two collaborations at the Large Hadron Collider (LHC), the ATLAS and Compact Muon Solenoid (CMS), announced the discovery of a new boson at 125 GeV with properties similar to the Standard Model Higgs  $???$ . This discovery was fueled by the investigation into the Higgs decays to the vector bosons  $ZZ$  and  $\gamma\gamma$ . Soon after, evidence for the Higgs coupling to matter was found through the  $\tau^+\tau^-$  and  $b\bar{b}$  decays  $????$ . Whether the newly discovered boson is indeed the expected Standard Model Higgs remains to be determined. Insofar, all of the different decay modes will be investigated to search for deviations from the Standard Model predictions.

This leads to the study of the Higgs decay to  $\mu^+\mu^-$ . Although this decay is the smallest branching fraction expected to be detected  $??$ , the dimuon decay offers high efficiency and excellent momentum resolution, which should lead to a narrow peak over the falling background, mostly Drell Yan events. The tiny branching fraction enables greater sensitivity to small deviations from the predicted decay rate and in this respect offers an advantage over other channels where a miniscule deviation could be drowned out. Furthermore, the Higgs coupling to second generation fermions remains to be determined.

This dissertation presents the search for the Standard Model Higgs Boson decaying to  $\mu^+\mu^-$  using the proton-proton collision data recorded by the CMS experiment in 2016. In order to maximize the data available for the search, the first machine learning in the L1 Trigger

system at the LHC was developed and deployed for 2016 data collection. To further maximize the sensitivity of the search, an additional machine learning technique is invented to categorize events based upon the detector resolution and the event kinematics. The search looks for a Higgs boson with a mass between 120 and 130 GeV and presents the expected and observed upper limits in this range.

The dissertation first presents an overview of the Standard Model and the symmetry breaking mechanism within the theory responsible for the Higgs. After covering the theoretical basis for the Standard Model Higgs, the accelerating apparatus responsible for accelerating and colliding the protons, the LHC, is covered. Then the CMS detector responsible for measuring the paths, momentum, and energy of the particles emerging from every collision is reviewed. Next the machine learning implementation in the L1 trigger that enables the detector to save more of the relevant collisions data is detailed. After, the search for  $H \rightarrow \mu^+ \mu^-$  is presented and then finally the conclusions.

## CHAPTER 2 THE LARGE HADRON COLLIDER AND THE CMS EXPERIMENT

### 2.1 Large Hadron Collider

The Large Hadron Collider is a particle collider near Geneva, Switzerland run by the European Organization for Nuclear Research (CERN). The LHC is the largest and most powerful particle collider ever built, designed to collide protons with a center of mass energy of 14 TeV and a luminosity of  $10^{34} \text{cm}^{-2} \text{s}^{-1}$  ?. The luminosity is given by

$$L = \frac{n_b f N_p^2 \gamma}{4\pi \epsilon_n \beta^*} \quad (2-1)$$

where  $n_b$  is the number of bunches in each ring,  $f$  is the frequency for a bunch to circle the ring,  $N_p$  gives the number of protons in a bunch, and  $\gamma$  is the Lorentz factor.  $\epsilon_n$  is the normalized transverse emittance, a measure of the spread of the beam in momentum and position space.  $\beta^*$  measures the focus of the beam at the interaction point.  $\epsilon_n \beta^*$  represents the transverse area at the point of interaction. A large luminosity is characterized by a high frequency of bunch crossings with lots of protons in each bunch packed as densely as possible, and a large luminosity results in a high rate of collisions. With many collisions at high energy, the detectors can collect enough events from yet unexplored regimes of physics to discover new physics or to verify or discard the predictions different hypothesis, so these parameters are very important.

The collider itself is 26.7 km in circumference 45-170 m underground. 8.3 T supercooled superconducting magnets operating at 2 K steer the high energy proton beams. In order to save money the LHC not only reuses the tunnels of a previous collider, the Large Electron Positron Collider (LEP), but also reuses older accelerators which were state of the art at their time. These older accelerators ramp up the energy of the protons and inject them into the LHC. All of this together makes up the CERN accelerator complex.

First, the protons are created from a source of Hydrogen gas. The hydrogen atoms of the gas are placed into a large electric field that separates the atoms into unbound protons

## CERN's Accelerator Complex



Figure 2-1. The CERN Accelerator Complex ?

and electrons. The protons are then sent to a radio frequency quadrupole which focuses the protons and accelerates them. The radio frequency field is stronger for the protons in the back than in the front and consequently squeezes them into a tighter bunch. The protons then proceed to a linear accelerator, LINAC2, where they are accelerated to 50 MeV or 5% of the speed of light ( $c$ ). The protons then enter a series of synchrotrons. A synchrotron is a device that accelerates particles by guiding them around a fixed circular path with a magnetic field and boosting their speed with an electric field as they pass a certain point. Since a faster particle bends less in the same magnetic field, the magnetic field strength is synchronized with the speed of the accelerating particles to keep them in the fixed circular path.

After LINAC2 the protons enter the first of the synchrotrons, the Proton Synchrotron Booster (PSB) accelerating the protons to 1.4 GeV ( $0.81c$ ). From here the protons are injected into the Proton Synchrotron (PS) and accelerate to 25 GeV ( $0.999c$ ). The PS then injects

the protons into the Super Proton Synchrotron (SPS) further accelerating them to 450 GeV (0.999999c). Finally the protons are injected into the LHC where they accelerate up to 6.5 TeV (0.999999999c). Once accelerated to the appropriate collision energy, the proton beams are made to collide in the different detectors located around the ring. By colliding enough protons at large enough energies it is possible to probe corners of physics that have never been seen before. The two general purpose detectors at the LHC, ATLAS and CMS, are used to look for signs of new physics like the Higgs boson, dark matter, and extra dimensions by measuring the energy, the momentum, and the paths of the particles coming out of the collisions.

## 2.2 Compact Muon Solenoid Detector

The CMS detector, located in Cessy, France, is 21.6 m long, 15 m in diameter, and weighs more than the Eiffel Tower. Not only is the detector a massive and complex device it's also run by a huge collaboration involving approximately 3,800 people from 200 institutes spanning 43 different countries ?. The greatest achievement of the collaboration to date is the discovery of a Higgs like particle in 2012, a feat shared with ATLAS.



Figure 2-2. The CMS detector ?

CMS was built primarily to look for the Standard Model Higgs and signs of Beyond Standard Model (BSM) physics like Supersymmetry, extra dimensions, or new heavy weak bosons ?. Because BSM and Higgs decays to muons and electrons often have the highest

signal to background ratio, CMS is designed to identify and measure these particles with a high accuracy. In layman's terms a high signal to background ratio just means that these events have fewer look-alikes. Jets <sup>1</sup> and photons are measured to a high degree of accuracy as well. In order to measure the energy, momentum, and location of the different types of particles CMS deploys a variety of subdetectors working in concert. The defining feature of the detector is an extremely powerful solenoid which enables the accurate measurement of momentum for charged particles. The tracker and calorimeters fit snugly within the 6 m diameter solenoid. The muon detectors reside outside the magnet but within the return yoke.

### 2.2.1 Silicon Tracker

The 3.8T magnetic field inside the solenoid enables the tracker to measure the transverse momentum of charged particles based upon the curvature of the track. Charged particles with lower transverse momentum ( $p_t$ ) bend more in a magnetic field than high  $p_t$  particles. As such, a measurement of the deviation of a curved track from a straight line, the sagitta, can be used to measure the curvature and determine the momentum ?.

$$p_t \cong \frac{L^2 q B}{8s} \quad (2-2)$$

Here L is the length of the straight line between the first and last position measurements, q is the charge of the particle, B is the magnetic field, and s is the sagitta.

The equation for the error in the momentum measurement shows that a higher magnetic field enables better  $p_t$  resolution, illuminating the design choice for a powerful magnet.

$$\frac{\delta p_t}{p_t} \propto \frac{p_t}{L^2 B} \quad (2-3)$$

---

<sup>1</sup> When a quark or gluon is created it can't exist alone, since it has color charge, and pulls other quarks from the vacuum creating a tight cone of composite colorless particles as well as their decays. This cone of particles is called a jet.



Figure 2-3. The sagitta measurement

The silicon tracker is made of tiny reverse biased bipolar diodes. When a charged particle travels through one of these diodes the ionization force of the particle releases electron hole pairs beyond the electrostatic equilibrium, inciting a current to flow. The tracker needs to be small enough such that the particles flowing through it don't deposit much energy. Energy deposition in the tracker would throw off energy measurements in the calorimeters. This means that the tracker needs to be smaller than a few radiation lengths <sup>2</sup>. The tracker at the thickest part is one radiation length. The tracker is placed nearest the collision point in order to identify primary and secondary vertices and to measure the momentum of particles before they are tainted by interactions with other detectors. <sup>3</sup> Being so near the collision point, the silicon tracker is bombarded by a constant flux of high intensity radiation. As such, the tracker is carefully designed to be robust to this radiation rich environment.

### 2.2.2 Calorimeters

The Electromagnetic Calorimeter (ECAL) is right outside the tracker and its main goal is to measure the energy of electrons and photons. It's designed to contain entire electromagnetic

---

<sup>2</sup> the length scale over which an electron deposits a substantial amount of energy into the material

<sup>3</sup> Vertex is shorthand for the location of the collision or decay that produced a set of particles.

showers for these particles and is consequently many radiation lengths thick. The ECAL is made of lead tungstenate scintillating crystals which release an amount of light proportional to the energy deposition. The light is collected and the total energy is calculated. The separation into individual crystals allows some spatial resolution as well. Particles with larger mass deposit less energy per unit distance into a solid. Many of the hadronic particles make it through the ECAL for this reason.



Figure 2-4. A slice of the CMS detector ?

The HCAL is placed outside the ECAL to collect the energy of the particles that survived the other subsystems, mostly strongly interacting hadronic particles from jets. The HCAL works in a similar manner to the ECAL except that layers of plastic scintillating material are interspersed with layers of a dense passive absorber like brass or steel. The density of the passive absorber increases the chance of interaction and shower production thus reducing the total length of the hadronic shower and enabling the measurement of the total energy for most of the cascades. Again the scintillation light is collected to determine the energy. If the ECAL and HCAL were placed outside the magnet the particles would interact with the solenoid material before entering the calorimeters, throwing off their measurements. The showers in the calorimeters are a consequence of the electromagnetic and strong forces, which means that particles without these interactions pass through the materials undetected, e.g. neutrinos or BSM weakly interacting particles. Since the momentum in an interaction is conserved any imbalance means that some particles escaped the detector. If there is an excess of missing



momentum beyond the amount expected due to neutrinos this may indicate the existence of dark matter or some other BSM particle. In order to measure the missing energy correctly it's important that the HCAL is built without any gaps and that it is dense enough to collect the energy of the strongly and electrically interacting particles.

While the momentum resolution in the tracker is proportional to the momentum, the energy resolution in the calorimeters decreases with increasing energy ?.

$$\frac{\delta E}{E} = \sqrt{\left(\frac{S}{\sqrt{E}}\right)^2 + \left(\frac{N}{E}\right)^2 + C} \quad (2-4)$$

The first term in the square root describes statistical fluctuations. The energy measured is proportional to the number of photons captured which has poissonian fluctuations and the error ( $\delta E$ ) for this term is  $\propto \sqrt{E}$ . The second term describes noise in the electronics whose error is energy independent, and the last term describes the errors in energy calibration which are proportional to energy.

### 2.2.3 Muon System

Neutrinos aren't the only Standard Model particles that make it through the tracker, ECAL, and HCAL. Muons have a relatively long lifetime  $\sim 10^{-6}$ s with  $c\tau \sim 100$ m. The large gamma factor in combination with their long lifetime enables them to travel hundreds of kilometers on average, well through the entire CMS detector before decaying. Muons are charged so their tracks show up in the tracker and some energy is deposited in the calorimeters but, muons are so much more massive than electrons that the energy deposition in the ECAL is minimal. Making it through the ECAL the muons enter the HCAL. The HCAL is designed to stop strongly interacting hadronic particles and collect their energy. But muons don't interact with the strong force and make it through the HCAL as well. This enables the muon system to be placed outside the magnet.

The muon system consists of a few different types of detectors which all involve the same basic principle. The charged muon ionizes some gas and the ionized particles are attracted to charged surfaces initiating a current in the surfaces. With a large enough voltage differential

between the charged surfaces the the ionized particles may gain enough kinetic energy to further ionize other atoms in the gas initiating an avalanche effect and reducing the need for signal amplification later. The muon system uses this strategy in the different detectors. The types of detectors in the muon system are the Cathode Strip Chambers (CSC), the Drift Tubes (DT), and the Resistive Plate Chambers (RPC) ?.



Figure 2-5. A Look at the Muon System ?

### 2.2.3.1 Drift Tubes

The drift tubes are located in the barrel portion of CMS. Throughout the majority of the barrel the magnetic field is basically uniform. The drift tubes have aluminum plates on the top and bottom separated by aluminum I-beams shown in Figure 2-6. A wire acts as the anode and the I-beams are the cathodes. The tubes are designed to provide a constant drift velocity throughout each tube.



Figure 2-6. A Drift Tube ?

When a charged particle flies through the tube it ionizes the gas inside. The electrons drift at constant velocity to the anode. The distance from the anode is deduced from the drift time, utilizing the fact that the ionized electrons drift with a constant velocity. This calculation does however require a reference time. In each chamber the drift tubes are placed in layers and the average crossing time in the chamber is used as the reference time.

### 2.2.3.2 Cathode Strip Chambers

The CSCs are located in the endcaps of the detector which range in  $|\eta|$  from 0.8 to 2.4. One of the reasons the endcaps use CSCs instead of DTs is the nonuniform magnetic field which would adversely affect the drift times in the DT system. In this system there are oppositely charged strips and wires running roughly perpendicular to each other.

When a muon flies through the CSC it induces charge on the wires and the strips and ionizes gas in the chamber. The ionized particles in the gas float to the charged strips and wires initiating a current in the nearby wires and strips. The induced charge from the muon itself also contributes to the currents. The most intense currents should be those associated with the location of the muon. The position resolution in the phi direction is roughly  $100 \mu\text{m}$ .



Figure 2-7. A Cathode Strip Chamber ?

### 2.2.3.3 Resistive Plate Chambers

The RPCs are located both in the barrel and in the end caps. The RPCs have excellent timing resolution on the order of 1 ns. The RPCs use their excellent timing resolution to determine each particle's bunch crossing of origin. The accurate and rapid timing information helps with the online selection of muons, a huge priority for CMS considering that many interesting collisions produce muons. For this reason the RPCs focus on efficient online selection of muons instead of accurate offline reconstruction ?.

In this way the RPCs complement the DTs and CSCs. The RPCs consist of two high resistance parallel plates surrounding a volume of gas. The outsides of the plates are painted with graphite paint forming the electrodes. A large voltage differential is kept between the electrodes. When a charged particle crosses the plates it induces an electrical discharge in the plates which remains localized in time and space due to the large resistivity.

### 2.2.4 Trigger System

Collision events come at a rate of 10 MHz with each event taking up roughly a MB of information. If the detector had to store all of the information from each event this would amount to pushing terabytes of information into a storage system every second, which is remarkably infeasible. To deal with this issue CMS utilizes a trigger system, which selects only

interesting events cutting the rate down from 10 MHz to 1 KHz ?. Since bunch crossings happen every 25 ns the trigger needs to operate at an incredibly high rate.

CMS tackled this issue by dividing the trigger into different tiers. The Level 1 Trigger is the first stage of the trigger system made from custom hardware which can operate at fantastic speed. The Level 1 Trigger reduces the rate from 10 MHz to 100 KHz and the events passing the L1 Trigger go onto the High Level Trigger (HLT) which further reduces the rate to 1 KHz. Due to the lower input rate the HLT can operate in software.

#### **2.2.4.1 L1 Trigger**

The L1 Trigger is made of up different subsystems that work together to decide whether to keep the data from a beam crossing for further processing. The University of Florida works with the Level 1 muon trigger system, the Endcap Muon Track Finder (EMTF) in particular. The muon system needs to determine the transverse momenta of muons and their location and choose the best candidates. Each of the different muon detectors have their own local triggers which send their best muon tracks to the Global Muon Trigger (GMT). The GMT chooses the best muon candidates from that set and passes these on to the Global Trigger (GT). The GT combines the information from the calorimeter triggering system. The GT uses this combined information to check whether the bunch crossing should be sent to the HLT or discarded. The L1 Trigger has many different trigger criteria defining separate triggers which are the trigger bits. If an event passes any of the triggers then it is forwarded for further processing.

The Track Finders (TF) play an important role in the L1 Trigger system. The EMTF combines the location and direction information from the different CSC stations into muon tracks and calculates the transverse momenta for the different tracks. The EMTF chooses the best candidates (highest momentum and highest quality) to send to the GMT. The Drift Tube Track Finder (DTTF) performs a similar process for muons in the DT system. The RPC system calculates the location and direction and forms tracks in the same stage. In the process the RPC trigger system assigns transverse momenta and quality, and like the others chooses the best tracks to send to the GMT.



Figure 2-8. The L1 Trigger Architecture ?

## CHAPTER 3

### THE STANDARD MODEL

The Standard Model (SM) of particle physics is an incredibly successful theory that correctly describes the physics of all known particles and forces that make up the universe, excluding gravity<sup>1</sup>. The particles of the SM come in two types: fermions and bosons. Fermions are the spin  $\frac{1}{2}$  particles and make up the different types of matter, and bosons are the integer spin particles responsible for the different forces. Electrons are the most familiar type of fermion, but there are more exotic kinds like the up and down quarks that make up protons and neutrons. Electrons combine with the quark based nuclei to make atoms and account for nearly all of the matter in our day to day experience, but there are actually many other fermions. In fact, there are three generations of quarks and leptons<sup>1</sup> with each generation heavier than the next. The up and down quarks are the first generation of quarks, charmed and strange are the next, and top and bottom are the third generation. For the leptons, the electron and electron neutrino are the first generation, the muon and muon neutrino are the second, and the tau and the tau neutrino the third. Each fermion also has a corresponding antiparticle. As an example, the positron is the antiparticle for the electron.

The universe would be pretty boring if particles couldn't interact to form more complex objects like atoms, molecules, and even people. Luckily the different forces exist as well, and these forces are described by the spin 1 bosons. Fermions attract and repel by exchanging bosons, which is why the bosons are often called force carriers. The force mediating bosons are the gluons, photons, and the W and Z particles. Gluons mediate the strong force, photons the electromagnetic force, and the W and Z bosons mediate the weak force. Every force has an associated charge: particles with electric charge can interact through the electromagnetic force, those with color charge may interact via the strong force, and those with isospin may interact

---

<sup>1</sup> Leptons are fermions like the electron that aren't quarks. The quarks interact with the strong force that binds nuclei together and the leptons do not.



Figure 3-1. The Standard Model Particles

through the weak force. The fundamental forces and particles interact to make the familiar composite objects that surround us in our daily lives. The strong force binds quarks to form protons and neutrons, and even binds the protons and neutrons together to form nuclei, and the electromagnetic force binds electrons and nuclei to form atoms. The size of the composite objects gives an idea of the relative strength of the forces. A proton is  $10^{-15}$  meters in size while an atom is  $10^{-10}$  meters and a solar system is  $10^{12}$  meters. The more tightly bound the stronger the force. In fact the ratio of the strength of the forces is like so  $1:10^{-3}:10^{-16}:10^{-41}$ , strong : electromagnetic : weak : gravitational <sup>2</sup> .

Of all the particles predicted by the SM, only the Higgs boson remains to be found. The Higgs is the only spin 0 particle of the SM, and plays a special role. As the universe cooled from the Big Bang the Higgs field went through a phase transition and settled into a nonzero ground state forming a condensate. The potential energy from the interactions with this nonzero ground state give the massive fundamental particles their mass. With such a large

<sup>2</sup> Gravity is just included for perspective. The Standard Model does not describe this force and reconciling gravity with quantum mechanics is an open problem.



role in the SM, finding this particle or a BSM Higgs has been a huge priority for the CMS collaboration ?. In 2012 a Higgs particle with a mass of 125 GeV was found and to date remains consistent with the Standard Model. However, the properties need to be investigated further before declaring the discovered Higgs the Higgs of the Standard Model.

In order to lay out the Standard Model in mathematical terms, a bunch of background information needs to be covered, and since the Standard Model is described by Quantum Field Theory (QFT), the mathematical formulation of QFT will be covered first. After developing the necessary mathematical formalism, the Standard Model and the Higgs mechanism will be derived and explained. Afterwards the experimental search for Higgs to dimuons will be described.

In this dissertation  $\hbar$  and the speed of light,  $c$ , are set to 1. Moreover, 0,1,2,3 and  $t,x,y,z$  are used interchangeably to label the components of a four vector. When relevant 0 represents time and 1,2,3 represent  $x,y,z$  respectively. Einstein summation notation is used indicating that repeated indices are summed over, so  $x_i x_i y_j y_j$  is shorthand for  $\sum_i \sum_j x_i x_i y_j y_j$ . Repeated Greek indices assume sums over all four space and time components, while repeated Roman indices assume sums over only the spatial components.

### 3.1 Quantum Field Theory

The mathematical framework used to describe the physics of the SM as well as other Beyond Standard Model (BSM) field theories is called Quantum Field Theory (QFT). QFT enables the predictions of different measurable probabilities. One of the most important is the probability for a set of particles to come out of a collision of an initial set of particles. Another important one is the probability for a single particle to decay into another set of particles. These probabilities are encompassed in the cross sections and branching fractions. For example, the theory of the SM predicts the cross section for two protons to collide and create a Higgs. As another example, the SM also predicts the branching fraction for a Z boson to decay into two muons. These probabilities can be measured simply by colliding particles and counting the outcomes which in turn means that the theory can be tested. In fact, any

QFT model can be tested in this manner. Quantum Field Theory is written down in terms of a Lagrangian, and the math for the various predictions follows from there, but first a brief aside about particles.

### **3.1.1 Particles, Symmetries, and Labels**

Since QFT makes quantitative predictions in terms of particle collisions and particle decays it's interesting to contemplate what a particle really is. Consider an observer in a frame  $x$  with particle  $p$  and an observer in another frame  $x'$ . If the observer in  $x'$  can't identify particle  $p$  as well then it doesn't make sense to call  $p$  a particle. More concretely, consider a world where in frame  $x$  an observer sees a neatly stacked deck of cards, but in  $x'$  the observer sees the cards scattered all over the place. Calling the deck of cards a particle doesn't really make sense. On the other hand, both parties can still agree on the individual cards which kept the same suit and value. The suit and value are conserved quantities identifiable between frames. If the two observers get together later and compare notes they can see what happened to each card upon transforming from  $x$  to  $x'$  and work out a set of rules. The king of hearts may do one thing and the 10 of clubs another. They can then add the different forces into play repeat the process and compare again. Figuring this all out allows the two to determine the laws of physics for the fundamental pieces called particles.

This idea leads to Wigner's view: a particle is an object with conserved quantities that observers can agree on between frames. In our universe these labels are the mass, charge, spin, color, and isospin, with each label attached to a specific symmetry or group of symmetries. And because different observers can agree on these quantities they can compare notes and work out the laws of physics for the different types of particles. The deep connection between conserved quantities, symmetries, and labels for particles will be seen later. For now the focus is to work out laws that people in different frames can confirm, and this is where the Lagrangian formalism comes into play.

### 3.1.2 The Lagrangian Formalism

The Lagrangian formalism is a mathematical device that allows physicists to describe the evolution of a physical system over time, and it's within this formalism that QFT can be built. But before building the full mathematics of QFT, a simple example describing the free Newtonian particle is covered. The Newtonian example serves to illuminate the fact that the laws of physics are indistinguishable in different inertial frames. Moving to the full relativistic theory reveals that the symmetry goes deeper. Investigating the symmetries of the relativistic theory end up leading the way to QFT, a relativistic description of quantum mechanics.

So let's get into the framework. The goal of physics is to describe how a physical system evolves over time, and this evolution is usually given by some differential equation describing the state the system will take in the next interval of time given the current time. Moving from state to state from one interval of time to the next, the system traces out a path in some abstract space of possible states. As a concrete example, the differential equations describing the motion of a Newtonian particle determine the position and velocity of the particle at the next instant of time based upon the values in the previous instant of time. So how does one get the appropriate differential equation? At classical scales, nature tries to minimize the difference between the energy spent <sup>3</sup> and the energy available to spend and it minimizes the action  $S$ . At the more fundamental level of quantum mechanics, all possible paths contribute with different phases, but those close to the minimum add constructively. As the action gets larger and the paths away from the minimum count less and less and the quantum rule agrees with the classical rule. The classical case is analyzed first to derive certain symmetries. In the equation below,  $L$  is the Lagrangian,  $T$  is the kinetic energy and  $U$  is the potential energy.

$$S = \int L dt = \int T - U dt \quad (3-1)$$

---

<sup>3</sup> Spent here just means used as kinetic energy.

At an extremum of  $S$ ,  $\delta S = 0$  if  $S$  is smooth:  $S$  must decrease, go through a slope of zero, and then increase (or vice versa). For a true extremum,  $\delta S$  must be 0 in all directions. So to get the equations of motion vary the parameters of  $L$  and solve the system of equations for the values that yield zero change in the action.

$$\begin{aligned}\frac{\delta S}{\delta z_1} &= \int L(z_1 + dz_1, z_2, \dots) dt - \int L(z_1, z_2, \dots) dt = 0 \\ \frac{\delta S}{\delta z_2} &= \int L(z_1, z_2 + dz_2, \dots) dt - \int L(z_1, z_2, \dots) dt = 0 \\ &\dots\end{aligned}\tag{3-2}$$

Following this process provides the Euler-Lagrange equations, describing how the parameters  $z$  evolve over time. The  $z$ 's may be the position and velocity, or the quantum fields, or the temperature and volume or some other set of parameters that describe the system. The Lagrangian for a Newtonian free particle in one dimension is pretty simple and gets the point across.

$$S = \frac{1}{2} \int m \dot{x}^2 dt \tag{3-3}$$

If the action is at an extremum, perturbing the path  $x(t)$  by adding the infinitesimal  $\epsilon(t)$  leaves the action unchanged.

$$S' = \frac{1}{2} \int m(\dot{x} + \dot{\epsilon})^2 dt = \frac{1}{2} \int m(\dot{x}^2 + 2\dot{x}\dot{\epsilon} + \dot{\epsilon}^2) dt = \frac{1}{2} \int m(\dot{x}^2 + 2\dot{x}\dot{\epsilon}) dt \tag{3-4}$$

$$\delta S = S' - S = 0 = \frac{1}{2} \int m(\dot{x}^2 + 2\dot{x}\dot{\epsilon}) dt - \frac{1}{2} \int m\dot{x}^2 dt = \int m\dot{x}\dot{\epsilon} dt \tag{3-5}$$

Since  $x(t)$  is fixed at the boundaries of the integral,  $\epsilon$  must be zero at  $t_o$  and  $t_f$ , so integrating by parts yields the following equation.

$$\delta S = 0 = \epsilon(t_f)\dot{x}(t_f) - \epsilon(t_o)\dot{x}(t_o) + \int m\ddot{x}\epsilon dt = 0\dot{x}(t_f) - 0\dot{x}(t_o) + \int m\ddot{x}\epsilon dt = \int m\ddot{x}\epsilon dt \quad (3-6)$$

And this equation must be zero for any infinitesimal deviation  $\epsilon$ .

$$\delta S = 0 \rightarrow m\ddot{x} = 0 \quad (3-7)$$

So a free particle keeps the same velocity over time. Note that a Newtonian boost by constant velocity  $v \rightarrow v' = v + u$ <sup>4</sup> leaves the equations of motion consistent. In the unprimed frame the particle has velocity  $v$  with 0 acceleration. In the primed frame the particle has velocity  $v + u$  with 0 acceleration. Both observers see the particle act as if there are zero forces.

$$S = \frac{1}{2} \int m(v + u)^2 dt = \frac{1}{2} \int m(v')^2 dt \rightarrow \delta S = 0 \rightarrow m \frac{d}{dt}(v + u) = m \frac{d}{dt}(v') = 0 \quad (3-8)$$

If  $u$  is not constant but a function of time  $u(t)$  then the equations of motion do not describe the same time evolution.

$$m \frac{d}{dt}(v + u) = m\dot{v} + m\dot{u} = 0 \rightarrow \dot{v} = -\dot{u} \quad (3-9)$$

In the case where  $u(t)$  depends upon time, the difference between the primed and unprimed frames' equations of motion is then  $\delta F$ .

$$\delta F = m \frac{d}{dt}(v + u) - m \frac{dv}{dt} = m\dot{v} + m\dot{u} - m\dot{v} = m\dot{u} \quad (3-10)$$

In the unprimed frame, the particle identified by the mass moves with constant velocity,  $\dot{v} = 0$ . The observer in the primed frame looks at the particle with the same mass and sees

---

<sup>4</sup> Renaming  $\dot{x}$  as  $v$ .

it change velocity given by the equation  $\dot{v} = -\dot{u}$ . As an example, set  $v$  and  $u, \dot{u}$  to zero for all times before  $t=0$ , and let  $u, \dot{u}$  turn on after time 0. Both observers will agree that the particle is stationary up until time 0. After which, the observer in the primed frame will see the particle accelerate in strange ways. Meanwhile, the unprimed frame will continue to observe a stationary particle.

In general, every inertial frame finds  $\delta F = 0$  and every accelerating frame finds an extra force  $\delta F$  unique to its acceleration. In this way no observer in an inertial frame can perform an experiment and determine which inertial frame he or she is in. On the other hand, each accelerating frame is identified by its  $\delta F$ . Put another way, in every inertial frame a ball released at rest remains at rest. In an accelerating frame the ball will accelerate according to the motion of the frame  $\delta F$  and this change in the laws of physics identifies the frame in a unique way. Conversely, the laws of physics remain the same boosting between inertial frames, and this invariance is a symmetry of physics. Of course this example is Newtonian and the correct way to boost is given by the Lorentz transformation from Special Relativity, but this gets the point across.

Delving further along the path of symmetry, the fundamental forces depend only on the distance from the charge and not the direction implying that rotations are also a symmetry. This can be seen by looking at the Lagrangian.

$$L = \frac{1}{2}m\dot{\vec{x}}^2 - U((\vec{x} - \vec{x}')^2) \quad (3-11)$$

Rotations leave dot products and consequently the magnitude of vectors unchanged so the Lagrangian is invariant under this transformation. Naturally if the Lagrangian is invariant the equations of motion will be as well.

$$m \frac{d\vec{v}}{dt} = \vec{\nabla} U \quad (3-12)$$

In the equations of motion above, both sides are vectors and vectors transform the same way under rotations so the equations of motion are invariant. Note that in the case of

rotations both the Lagrangian and the equations of motion are invariant. While for Newtonian boosts only the equations of motion were invariant. This is due to the fact that Newtonian mechanics is the low velocity limit of relativistic mechanics. In the theory of Special Relativity the action for a massive free particle is written like so.

$$S = \int \frac{m}{2} u^\mu u_\mu d\tau \quad (3-13)$$

Just as rotations preserve the dot product, Lorentz transformations (boosts and rotations) preserve the four vector product. Insofar, both the relativistic Lagrangian and the resulting equations of motion remain invariant under a boost or rotation to a new inertial frame. Building a Lagrangian out of four vector products guarantees that it will remain invariant under Lorentz transformations. However, four vectors aren't the most fundamental objects with invariant products. By studying the properties of the Lorentz group, it's possible to find even more fundamental building blocks called spinors.

### 3.1.3 QFT From Symmetry

The laws of physics are invariant under boosts and rotations, and the Lagrangian provides a mathematical framework for physical predictions. These facts together imply that there's a good shot at building a proper QFT by creating the appropriate invariant Lagrangian. Four vector products remain invariant under Lorentz transformations so they are a natural ingredient, but there are other mathematical objects that could be used as well. In this vein, the symmetries under rotations and boosts are investigated in order to look for other building blocks. The goal is to find three different representations of the Lorentz group and use one representation for scalar spin 0 particles, another for the spin  $\frac{1}{2}$  fermions, and another for the spin 1 bosons.

#### 3.1.3.1 Rotations

Rotations in three dimensions are described by the  $SO(3)$  group. Rotations preserve the lengths of vectors and the angles between them, which means that dot products between

vectors remain invariant as well. In three dimensions one can rotate about any of the three axes. The rotations about the x, y, and z axes may be characterized by the matrices below.

$$R_x = \begin{pmatrix} 1 & 0 & 0 \\ 0 & \cos \theta_x & -\sin \theta_x \\ 0 & \sin \theta_x & \cos \theta_x \end{pmatrix} \quad (3-14)$$

$$R_y = \begin{pmatrix} \cos \theta_y & 0 & \sin \theta_y \\ 0 & 1 & 0 \\ -\sin \theta_y & 0 & \cos \theta_y \end{pmatrix} \quad (3-15)$$

$$R_z = \begin{pmatrix} \cos \theta_z & -\sin \theta_z & 0 \\ \sin \theta_z & \cos \theta_z & 0 \\ 0 & 0 & 1 \end{pmatrix} \quad (3-16)$$

These rotations may be built up from repeated rotations by an infinitesimally small angle  $d\theta$ . The matrices characterizing an infinitesimal rotation are given by taking the limit as  $\theta$  goes to zero.

$$dR_x = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & -d\theta_x \\ 0 & d\theta_x & 1 \end{pmatrix} = 1 - id\theta_x \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & -i \\ 0 & i & 0 \end{pmatrix} = 1 - id\theta_x J_x \quad (3-17)$$

$$dR_y = \begin{pmatrix} 1 & 0 & d\theta_y \\ 0 & 1 & 0 \\ -d\theta_y & 0 & 1 \end{pmatrix} = 1 - id\theta_y \begin{pmatrix} 0 & 0 & i \\ 0 & 0 & 0 \\ -i & 0 & 0 \end{pmatrix} = 1 - id\theta_y J_y \quad (3-18)$$

$$dR_z = \begin{pmatrix} 1 & -d\theta_z & 0 \\ d\theta_z & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} = 1 - id\theta_z \begin{pmatrix} 0 & -i & 0 \\ i & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} = 1 - id\theta_z J_z \quad (3-19)$$



Repeating an infinitesimal rotation many times builds the finite rotation, and in this way, the  $J$  matrices generate rotations along their respective axes. As such, they are aptly referred to as the generators of the group. Doing some algebra reveals this to be the case. This point is confirmed below.

$$R = (1 - i\frac{\theta}{N}J)^N = 1 + (-id\theta J) + \frac{1}{2!}(-id\theta J)^2 + \frac{1}{3!}(-id\theta J)^3 + \dots = e^{-i\theta J} \quad (3-20)$$

Notice that even powers of  $J$  yield  $J^2$  and that odd powers of  $J$  return  $J$ .

$$= 1 - J^2 + J^2(1 + \frac{i^2}{2!}d\theta^2 + \frac{i^4}{4!}d\theta^4 + \dots) - iJ(d\theta + \frac{i^2}{3!}d\theta^3 + \frac{i^4}{5!}d\theta^5 + \dots) = (1 - J^2) + J^2 \cos \theta - iJ \sin \theta \quad (3-21)$$

Plugging in  $J_z$  reveals that this process does in fact rebuild the rotation matrix  $R_z$ .

$$\begin{aligned} R_z &= \left( \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} - \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 0 \end{pmatrix} \right) + \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 0 \end{pmatrix} \cos \theta_z + \begin{pmatrix} 0 & -1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} \sin \theta_z \\ &= \begin{pmatrix} \cos \theta_z & -\sin \theta_z & 0 \\ \sin \theta_z & \cos \theta_z & 0 \\ 0 & 0 & 1 \end{pmatrix} \end{aligned} \quad (3-22)$$

Similarly, the other generators rebuild their respective rotation matrices. The generators of the group are actually more fundamental than the rotation matrices. The multiplication table for the generators describes the algebra of the group, which describes the behavior of rotations at a local level. In fact the  $SO(3)$  rotation matrices are just one of the groups with this local algebra, and a specific group obeying the local algebra is analogous to a specific solution of a differential equation: each solution has a different global behavior yet each obeys the same

physics. Moreover the multiplication table can be specified without declaring any particular representation for the generators.

$$\begin{aligned}
J_x * J_y &= iJ_z + J_y * J_x \\
J_y * J_z &= iJ_x + J_z * J_y \\
J_z * J_x &= iJ_y + J_x * J_z
\end{aligned}
\tag{3-23}$$

The multiplication table can be specified in a more compact notation using the commutator <sup>5</sup>,  $[a, b] = ab - ba$ , and the antisymmetric tensor  $\epsilon$ .

$$[J_k, J_l] = i\epsilon_{klm}J_m \tag{3-24}$$

Finding 3x3 generators that obey the algebra and then repeatedly applying the infinitesimal transformations builds the  $SO(3)$  rotation group. The group acts on real 3x1 objects called vectors, and these 3x1 vectors are a suitable candidate for a Newtonian Lagrangian. Finding another representation obeying this algebra will provide a more fundamental ingredient for the Lagrangian and allow the construction of a proper QFT. Similar to the way real numbers are built from the squares of imaginary numbers, vectors are built from spinors.

Looking for the lowest order nontrivial  $n \times n$  matrices satisfying the algebra gives the 2x2 Pauli matrices. The 1x1 matrices are the trivial solution: 1x1 matrices are simply scalar complex numbers, which commute and therefore fail to satisfy the algebra unless all of the  $J$  matrices are 0. The objects these 1x1 operators act on are 1x1 numbers called scalars which remain invariant under rotations and correspond to spin 0. The solution for the 2x2 case, the Pauli matrices are given by

---

<sup>5</sup> The Lie Bracket defines the multiplication for a Lie Algebra and this reduces to the commutator for Lie groups of matrices like  $SO(3)$ . Notice that using the commutator below returns another member of the group and thus the group is closed under commutation.

$$\sigma_x = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \sigma_y = \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}, \sigma_z = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \quad (3-25)$$

However, plugging these into the commutator reveals a factor of two difference.

$$[\sigma_k, \sigma_l] = 2i\epsilon_{klm}\sigma_m \quad (3-26)$$

Defining  $J_k = \frac{1}{2}\sigma_k$  fixes this. These matrices act on an array of 2x1 complex numbers called spinors, and this new rotation group is called SU(2). Note that by starting with vectors and analyzing the SO(3) rotation group along with its underlying algebra, new mathematical objects have been discovered. The 1x1 matrices satisfying the rotation algebra make up the spin 0 representation of SU(2), the complex 2x2 matrices acting on complex 2x1 objects satisfying the algebra make up the spin  $\frac{1}{2}$  representation, and the complex 3x3 matrices acting on complex 3x1 objects satisfying the algebra make up the spin 1 representation. The pattern continues on. There are in fact many representations of SU(2). It's now possible to use these representations to build rotationally invariant Lagrangians, using the 2x1 spinors of SU(2) to model fermions and the 3x1 representation to model bosons. While rotationally invariant Lagrangians are important for nonrelativistic theories, the real goal is to break down four vectors in the same way to find the most fundamental ingredients for relativistic Lagrangians.

$$[J_k, J_l] = i\epsilon_{klm}J_m \quad (3-27)$$

### 3.1.3.2 The Lorentz Group

Four vector products are invariant with regards to rotations and boosts. This statement is defined by the mathematical equation below, where the  $\Lambda$  matrices represent the rotation/boost matrices of the Lorentz Group <sup>6</sup> and the  $\eta$  matrix is the Minkowski metric  $\begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 \\ 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & -1 \end{pmatrix}$ .

---

<sup>6</sup> The Lorentz group dealt with here is the proper orthochronous Lorentz Group SO(1,3).

$$x'_\mu x'^\mu = x_\mu x^\mu \rightarrow \eta_{\sigma\rho} \Lambda_\mu^\sigma \Lambda_\nu^\rho x^\mu x^\nu = \eta_{\mu\nu} x^\mu x^\nu \rightarrow \eta_{\sigma\rho} \Lambda_\mu^\sigma \Lambda_\nu^\rho = \eta_{\mu\nu} \quad (3-28)$$

In this 3 + 1 dimensional space the rotations and their corresponding generators are now given by the following R and J matrices, where the R matrices satisfy the equation above.

$$R_x = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & \cos \theta_x & -\sin \theta_x \\ 0 & 0 & \sin \theta_x & \cos \theta_x \end{pmatrix}, J_x = \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & -i \\ 0 & 0 & i & 0 \end{pmatrix} \quad (3-29)$$

$$R_y = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & \cos \theta_y & 0 & \sin \theta_y \\ 0 & 0 & 1 & 0 \\ 0 & -\sin \theta_y & 0 & \cos \theta_y \end{pmatrix}, J_y = \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & i \\ 0 & 0 & 0 & 0 \\ 0 & -i & 0 & 0 \end{pmatrix} \quad (3-30)$$

$$R_z = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & \cos \theta_z & -\sin \theta_z & 0 \\ 0 & \sin \theta_z & \cos \theta_z & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}, J_z = \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & -i & 0 \\ 0 & i & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix} \quad (3-31)$$

And the boosts are given by the following B matrices. These also leave the four vector product invariant.

$$B_x = \begin{pmatrix} \cosh \omega_x & \sinh \omega_x & 0 & 0 \\ \sinh \omega_x & \cosh \omega_x & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix} \quad (3-32)$$

$$B_y = \begin{pmatrix} \cosh \omega_y & 0 & \sinh \omega_y & 0 \\ 0 & 1 & 0 & 0 \\ \sinh \omega_y & 0 & \cosh \omega_y & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix} \quad (3-33)$$

$$B_z = \begin{pmatrix} \cosh \omega_z & 0 & 0 & \sinh \omega_z \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ \sinh \omega_z & 0 & 0 & \cosh \omega_z \end{pmatrix} \quad (3-34)$$

Looking at the differential boosts yields the generators K.

$$dB_x = \begin{pmatrix} 1 & d\omega_x & 0 & 0 \\ d\omega_x & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix} = 1 + d\omega_x \begin{pmatrix} 0 & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix} = 1 + d\omega_x K_x \quad (3-35)$$

$$dB_y = \begin{pmatrix} 1 & 0 & d\omega_y & 0 \\ 0 & 1 & 0 & 0 \\ d\omega_y & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix} = 1 + d\omega_y \begin{pmatrix} 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix} = 1 + d\omega_y K_y \quad (3-36)$$

$$dB_z = \begin{pmatrix} 1 & 0 & 0 & d\omega_z \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ d\omega_z & 0 & 0 & 1 \end{pmatrix} = 1 + d\omega_z \begin{pmatrix} 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 \end{pmatrix} = 1 + d\omega_z K_z \quad (3-37)$$

As before with the algebra for rotations in three dimensions the Lorentz algebra is defined by its multiplication table, but now there are rotations and boosts. The multiplication table is given by the commutation relations below, which can be confirmed by brute force computation.

$$[J_i, J_j] = i\epsilon_{ijk}J_k \quad (3-38)$$

$$[J_i, K_j] = i\epsilon_{ijk}K_k \quad (3-39)$$

$$[K_i, K_j] = -i\epsilon_{ijk}J_k \quad (3-40)$$

Notice that the commutator between J matrices returns another J matrix, but that the commutator between K matrices returns a J matrix. This means that the J operators form their own subgroup, but the K operators don't. On the other hand, mixing the Js and Ks up by defining the  $Y^\pm$  operators allows the Lorentz algebra to be represented by two independent subgroups.

$$Y^\pm = \frac{1}{2}(J_i \pm iK_i) \quad (3-41)$$

$$[Y_i^\pm, Y_j^\pm] = i\epsilon_{ijk}Y_k^\pm \quad (3-42)$$

$$[Y_i^\pm, Y_j^\mp] = 0 \quad (3-43)$$

Both of the Y groups have the same commutation relations as SU(2). In this way, the Lorentz algebra can be viewed as if two SU(2) rotation algebras have been glued together. This is similar to the way in which orthogonal basis vectors are stuck together to create a larger dimensional space. Now, in order to figure out how to use this space to build the appropriate Lagrangians, the individual subspaces must be investigated. So looking at  $Y^+$  alone is akin to looking along the  $Y^+$  axis by setting  $Y^-$  to zero. Using  $(y_+, y_-)$  to label the representation, the simplest nontrivial case along the  $Y^+$  axis is spin  $\frac{1}{2}$  x spin 0, given by  $(\frac{1}{2}, 0)$ .

Using  $Y_i^- = \frac{1}{2}(J_i - iK_i) = 0$  implies that  $J_i = iK_i$ , and since  $Y_i^+$  is the 2x2 representation obeying the SU(2) algebra,  $Y_i^+ = \frac{\sigma_i}{2}$ . Putting this together,

$$\begin{aligned}
J_i &= iK_i \\
\rightarrow Y_i^+ &= \frac{1}{2}(J_i + iK_i) = \frac{\sigma_i}{2} = \frac{1}{2}(J_i + J_i) = J_i \\
\rightarrow J_i &= \frac{1}{2}\sigma_i \\
\rightarrow K_i &= \frac{-i\sigma_i}{2}
\end{aligned} \tag{3-44}$$

Finally, the finite Lorentz transformations for the  $(\frac{1}{2}, 0)$  representation are given by

$$R^{(L)} = e^{i\theta_i J_i} = e^{i\theta_i \frac{\sigma_i}{2}} \tag{3-45}$$

for rotations, and

$$B^{(L)} = e^{i\phi_i K_i} = e^{\phi_i \frac{\sigma_i}{2}} \tag{3-46}$$

for boosts. These act on 2x1 objects called left-chiral spinors,  $\mathcal{L}$ . A general Lorentz transformation on a left-chiral spinor can be written

$$\Lambda^{(L)} = e^{\frac{i}{2}\theta_i \sigma_i + \frac{1}{2}\phi_i \sigma_i} \tag{3-47}$$

.

Replacing  $K_i$  with  $-K_i$  takes  $Y_i^+$  to  $Y_i^-$ , and gives the finite Lorentz transformations for the  $(0, \frac{1}{2})$  representation

$$R^{(R)} = e^{i\theta_i J_i} = e^{\frac{i}{2}\theta_i \sigma_i} \tag{3-48}$$

for rotations, and

$$B^{(R)} = e^{i\phi_i K_i} = e^{-\phi_i \frac{\sigma_i}{2}} \tag{3-49}$$

for boosts. These act on 2x1 objects called right-chiral spinors,  $\mathcal{R}$ . The general Lorentz transformation on a right-chiral spinor can be written

$$\Lambda^{(R)} = e^{\frac{i}{2}\theta_i\sigma_i - \frac{1}{2}\phi_i\sigma_i} \quad (3-50)$$

Last but not least, rank 2 spinors are given by the  $(\frac{1}{2}, \frac{1}{2})$  representation. This representation is a tensor combining two spinors via outer product, and as will be shown later, these objects are actually four vectors.

$$\alpha = \mathcal{L}\mathcal{R}^T \quad (3-51)$$

In order to transform  $\alpha$ , both  $\mathcal{L}$  and  $\mathcal{R}$  must be transformed.

$$\alpha' = \Lambda^{(L)}\mathcal{L}\mathcal{R}^T\Lambda^{(R)T} = e^{\frac{i}{2}\theta_i\sigma_i + \frac{1}{2}\phi_i\sigma_i}\mathcal{L}\mathcal{R}^Te^{\frac{i}{2}\theta_i\sigma_i^T - \frac{1}{2}\phi_i\sigma_i^T} \quad (3-52)$$

However it would be nice if the transformation term on the right side was the Hermitian conjugate of the transformation on the left side. This would be the case if  $\sigma_i^T$  was  $-\sigma_i^\dagger = -(\sigma_i^*)^T$ , and this requires a transformation that turns  $\sigma$  into  $-\sigma^*$ . So  $\mathcal{R}$  is rearranged such that  $\mathcal{R} \rightarrow \tilde{\mathcal{R}} = t\mathcal{R}$  where  $t\sigma_i t^{-1} = -\sigma_i^*$ . The matrix  $t = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$  satisfies the requirements. This change of basis for  $\mathcal{R}$  redefines  $\alpha$ ,

$$\alpha = \mathcal{L}\tilde{\mathcal{R}}^T. \quad (3-53)$$

Defined in this manner, the  $(\frac{1}{2}, \frac{1}{2})$  representation now transforms like so,

$$\alpha' = e^{\frac{i}{2}\theta_i\sigma_i + \frac{1}{2}\phi_i\sigma_i}\mathcal{L}\tilde{\mathcal{R}}^Te^{-\frac{i}{2}\theta_i\sigma_i^\dagger + \frac{1}{2}\phi_i\sigma_i^\dagger}. \quad (3-54)$$

Considering the fact that  $\sigma_i^\dagger = \sigma_i$ , the transformation reduces further,

$$\alpha' = e^{\frac{i}{2}\theta_i\sigma_i + \frac{1}{2}\phi_i\sigma_i}\alpha e^{-\frac{i}{2}\theta_i\sigma_i + \frac{1}{2}\phi_i\sigma_i}. \quad (3-55)$$



A transformation of the form  $M' = HMH^\dagger$  where  $H$  is a Hermitian matrix, preserves the Hermitivity of the matrix  $M$ . Namely,  $M'$  will be Hermitian if  $M$  is Hermitian,

$$M'^\dagger = (MH^\dagger H)^\dagger = HM^\dagger H^\dagger = HMH^\dagger = M'. \quad (3-56)$$

On the other hand,  $M'$  will be anti-Hermitian if  $M$  is anti-Hermitian,

$$M'^\dagger = (MH^\dagger H)^\dagger = HM^\dagger H^\dagger = H(-M)H^\dagger = -MH^\dagger H = -M'. \quad (3-57)$$

The transformation of the  $(\frac{1}{2}, \frac{1}{2})$  object  $\alpha$  is of this type. With this in mind, note that any complex matrix can be broken up into a Hermitian piece and an anti-Hermitian piece, and that these pieces remain independent under the  $(\frac{1}{2}, \frac{1}{2})$  Lorentz transformations constructed here. Then note that a general complex  $2 \times 2$  matrix has 8 free parameters, 4 from the Hermitian part and 4 from the anti-Hermitian part. Meanwhile,  $\alpha$  has only 4, two from each spinor<sup>7</sup>. This means that  $\alpha$  may be represented by the Hermitian space alone. An appropriate basis for this space is the collection of Pauli matrices plus the identity.

$$\sigma_0 = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \sigma_1 = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \sigma_2 = \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix} \sigma_3 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \quad (3-58)$$

Thus any  $(\frac{1}{2}, \frac{1}{2})$  object  $\alpha$  may be written in terms of its four independent parameters like so,

---

<sup>7</sup> A spinor has 2 complex components yielding 4 parameters. Two constraints reduce the number of free parameters to 2. One constraint requires a magnitude of 1, and the other requires that the overall phase doesn't matter.

$$\alpha = \alpha_0 \sigma_0 + \alpha_1 \sigma_1 + \alpha_2 \sigma_2 + \alpha_3 \sigma_3$$

$$\alpha = \begin{pmatrix} \alpha_0 + \alpha_3 & \alpha_1 - i\alpha_2 \\ \alpha_1 + i\alpha_2 & \alpha_0 - \alpha_3 \end{pmatrix} \quad (3-59)$$

Boosting  $\alpha$  along the z direction hints that the four components transform like a four vector.

$$\alpha' = e^{\frac{1}{2}\phi_3 \sigma_3} \begin{pmatrix} \alpha_0 + \alpha_3 & \alpha_1 - i\alpha_2 \\ \alpha_1 + i\alpha_2 & \alpha_0 - \alpha_3 \end{pmatrix} e^{\frac{1}{2}\phi_3 \sigma_3} \quad (3-60)$$

Exponentiating the  $\sigma_3$  matrix and multiplying everything yields,

$$\begin{pmatrix} \alpha'_0 + \alpha'_3 & \alpha'_1 - i\alpha'_2 \\ \alpha'_1 + i\alpha'_2 & \alpha'_0 - \alpha'_3 \end{pmatrix} = \begin{pmatrix} e^{\frac{1}{2}\phi_3} & 0 \\ 0 & e^{-\frac{1}{2}\phi_3} \end{pmatrix} \begin{pmatrix} \alpha_0 + \alpha_3 & \alpha_1 - i\alpha_2 \\ \alpha_1 + i\alpha_2 & \alpha_0 - \alpha_3 \end{pmatrix} \begin{pmatrix} e^{\frac{1}{2}\phi_3} & 0 \\ 0 & e^{-\frac{1}{2}\phi_3} \end{pmatrix} \quad (3-61)$$

$$= \begin{pmatrix} e^{\phi_3}(\alpha_0 + \alpha_3) & (\alpha_1 - i\alpha_2) \\ (\alpha_1 + i\alpha_2) & e^{-\phi_3}(\alpha_0 - \alpha_3) \end{pmatrix}.$$

Then solving the systems of equations makes the transformation clearer,

$$\alpha'_1 = \alpha_1$$

$$\alpha'_2 = \alpha_2$$

$$\alpha'_0 = (\cosh \phi_3) \alpha_0 + (\sinh \phi_3) \alpha_3$$

$$\alpha'_3 = (\sinh \phi_3) \alpha_0 + (\cosh \phi_3) \alpha_3. \quad (3-62)$$

Finally the transformation can be written as a 4x4 matrix acting on the 4x1 four vector,

$$\alpha' = \begin{pmatrix} \cosh \phi_3 & 0 & 0 & \sinh \phi_3 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ \sinh \phi_3 & 0 & 0 & \cosh \phi_3 \end{pmatrix} \begin{pmatrix} \alpha_0 \\ \alpha_1 \\ \alpha_2 \\ \alpha_3 \end{pmatrix} \quad (3-63)$$

The other transformations reproduce the usual 4x4 Lorentz transformations as well. In this way, four vectors are just rearranged versions of the  $(\frac{1}{2}, \frac{1}{2})$  rank 2 spinors. Similar to the way complex numbers are the square root of real numbers, spinors are the square root of four vectors. In practice it's easier to work with 4 vectors as 4x1 column vectors, though one could use the  $(\frac{1}{2}, \frac{1}{2})$  rank 2 spinor representation.

### 3.1.3.3 The Poincare Group and Particle Labels

The Poincare group is the Lorentz Group plus translations. Physics experiments should be the same with a rotated apparatus, an apparatus moving at a different constant velocity, and also at another location. Adding translations to the group amounts to adding another the set of translation generators  $P_i$ . In the "Particles, Symmetries, and Labels" section the argument was made that particles are things that observers agree on between frames. Distinguishable particles have labels that remain invariant. A spin  $\frac{1}{2}$  particle with mass  $m$  remains a spin  $\frac{1}{2}$  particle with mass  $m$  for all observers. As it turns out these labels relate to different subspaces in the representation of the group.

Consider a group with certain generators. When the operators of the group are represented a certain way, say as some NxN matrices there is a likelihood that there are some invariant subspaces. As an example, a representation of a group may act on a 5 dimensional space spanned by vectors  $e_1, e_2, e_3, e_4, e_5$  the group may always transform vectors in the  $e_1, e_2, e_3$  subspace into one another, and those in  $e_4, e_5$  into one another, but never mix up  $e_1, e_2, e_3$  with  $e_4, e_5$ . In this example the 5x5 operators could be decomposed into a 3x3 operator acting only on the  $e_1, e_2, e_3$  subspace and a 2x2 operator acting only on the  $e_4, e_5$  subspace. Each 5x5 member of the group could be written as a block diagonal matrix with a

3x3 block and a 2x2 block. The same goes for the generators. Since these subspaces retain their identity under the transformations of the group<sup>8</sup>, they may be considered different particles. The question is whether there are labels for the subspaces.

Labels in physics need to be measurable, and by the current understanding of quantum mechanics these must be eigenvalues of Hermitian operators. The operators that label these subspaces are called the Casimir operators and must be built from the generators of the group. The operators should give the same values for  $e_1, e_2$ , and  $e_3$  since they transform into one another and represent the same particle, which implies that the Casimir labeling operator must be proportional to the identity in that subspace. The same goes for the operator in the  $e_4, e_5$  subspace. The Casimir operator for SU(2) is the  $J^2$  operator which labels the spin of the particle. The Poincare group has two Casimir operators corresponding to two labels the mass,  $m$ , and the spin,  $j$ . Looking at the irreducible representations<sup>9</sup> of the Poincare group, the mass and spin arise naturally as labels for particles.

#### 3.1.3.4 Building the Lagrangian for a Free Scalar Particle

Analyzing the symmetries of the SO(3) rotation group revealed the SU(2) group, the 2x2 complex Pauli matrices, and the complex 2x1 spinors. Analyzing the symmetries of the Lorentz group revealed the  $(\frac{1}{2}, 0)$  left-chiral spinors, the  $(0, \frac{1}{2})$  right chiral spinors, and the four vectors encoded in the  $(\frac{1}{2}, \frac{1}{2})$  representation. Now, these pieces are put together to form relativistically invariant Lagrangians for free particles.

The  $(0, 0)$  representation of the Lorentz group is a scalar, which means that it remains the same under Lorentz transformations. This representation is used for spin 0 particles. The

---

<sup>8</sup> This assumes that the 3x3 and 2x2 pieces have no invariant subspaces besides themselves and 0.

<sup>9</sup> This is the group theory term for the block diagonal pieces and the corresponding invariant subspaces. The irreducible representations, irreps, are those that can't be broken down in terms of smaller invariant subspaces and smaller block diagonal matrices. An arbitrary representation of the group is built from these irreps.

equations of motion must relate the change in the scalar field  $\Phi$  at one moment in space and time to the value of  $\Phi$  at the next moment in space and time, so the Lagrangian must include both the field itself and the four vector derivative,  $\partial_\mu$ . Note that both Newton's equations of motion,  $F = m\ddot{x}$ , and the Schrodinger equation for the free particle,  $i\partial_t\psi = \frac{1}{2m}(-i\partial_x)(-i\partial_x)\psi$ <sup>10</sup> have at most second order derivatives. The same holds for Maxwell's equations of electromagnetism. Hence, as an assumption, the derivative term in the Lagrangian will be the lowest order possible. As a Lorentz invariant scalar, any power of  $\Phi$  can be included. On the other hand,  $\partial_\mu$  is a four vector, and must be paired with a  $\partial^\mu$  to form the invariant four vector product. Cross terms between these invariant pieces are also invariant, but lead to feedback between the derivatives and the value of the function. This means that the derivatives  $i\partial_t = E$  and  $-i\partial_i = P_i$  will change over time, but  $E$  and  $\vec{P}$  should remain constant for a free particle. The cross terms are thrown out to prevent this. The  $\Phi$  terms with an order different than  $\Phi^2$  cause the same problem, and these are thrown out too. All of these choices lead to the following action,

$$S = \int d^4x (c_0 + c_1\Phi^2 + c_2\partial_\mu\Phi\partial^\mu\Phi + c_3\partial_\mu\partial^\mu\Phi). \quad (3-64)$$

Note that the  $c_3$  term is a total derivative and by the divergence theorem depends only on the values at the boundary, which are fixed. This implies that the contribution to the action from the  $c_3$  term is the same regardless of how  $\Phi$  changes in the volume. Because the Euler-Lagrange equations depend only on the variation in the volume, this term cannot contribute to  $\delta S$  and  $c_3$  may be set to zero. The  $c_0$  term is a more obvious constant and does not affect  $\delta S$  either, so it may be set to zero as well.

$$S = \int d^4x (c_1\Phi^2 + c_2\partial_\mu\Phi\partial^\mu\Phi) \quad (3-65)$$

---

<sup>10</sup> Here the one dimensional case is presented for simplicity

Finding  $\Phi$  such that  $\delta S = 0$  amounts to applying the Euler-Lagrange equations  $\partial_\mu \frac{\partial \mathcal{L}}{\partial(\partial_\mu \Phi)} = \frac{\partial \mathcal{L}}{\partial \Phi}$ . These yield

$$\begin{aligned} 2c_2 \partial_\mu \partial^\mu \Phi &= -2c_1 \Phi \\ \rightarrow (-E^2 + \vec{P}^2) \Phi &= \frac{c_1}{c_2} \Phi = -m^2 \Phi \\ \rightarrow \frac{c_1}{c_2} &= -m^2 \end{aligned} \quad (3-66)$$

In order to get the correct dispersion relation for a relativistic particle,  $c_1$  is set to  $-\frac{1}{2}m^2$  and  $c_2$  is set to  $\frac{1}{2}$ . Notice that including any  $\Phi^n$  with  $n \neq 2$  in the Lagrangian would have contributed to the differential equation via  $\frac{\partial \mathcal{L}}{\partial \Phi}$  and that  $E, \vec{P}$  wouldn't be constant. Thus the equation wouldn't work for a free particle. The resulting equation of motion for the scalar particle,  $\partial_\mu \partial^\mu \Phi = m^2 \Phi$ , is called the Klein-Gordon equation, and provides the correct description for spin 0 particles. This was all derived using symmetry, a reasonable assumption about the order of the derivatives, and the fact that the Energy shouldn't change over time for a free particle. The final Lagrangian for the scalar particle is

$$S = \int d^4x \frac{1}{2} (\partial_\mu \Phi \partial^\mu \Phi - m^2 \Phi^2). \quad (3-67)$$

### 3.1.3.5 Building the Lagrangian for a Free Spin $\frac{1}{2}$ Particle

With the action for the free scalar particle in hand, the free spin  $\frac{1}{2}$  particle is up next. The spin  $\frac{1}{2}$  action must combine the  $\mathcal{L}$  and  $\mathcal{R}$  spinors of the  $(\frac{1}{2}, 0)$  and  $(0, \frac{1}{2})$  representations and the four vector,  $\partial_\mu$ , in a Lorentz invariant way. Moving forward in this regard, the  $\mathcal{L}$ ,  $\mathcal{R}$ , and  $\partial_\mu$  transformations are now analyzed to find the lowest order invariant combinations. The transformation for the left-chiral spinor is

$$\Lambda^{(L)} = e^{\frac{i}{2}\theta_i \sigma_i + \frac{1}{2}\phi_i \sigma_i} \quad (3-68)$$

,

and the transformation for the right-chiral spinor is,

$$\Lambda^{(R)} = e^{\frac{i}{2}\theta_i\sigma_i - \frac{1}{2}\phi_i\sigma_i} \quad (3-69)$$

Taking the Hermitian conjugate of the left-chiral transformation gives,

$$(\Lambda^{(L)})^\dagger = e^{-\frac{i}{2}\theta_i\sigma_i^\dagger + \frac{1}{2}\phi_i\sigma_i^\dagger} = e^{-\frac{i}{2}\theta_i\sigma_i + \frac{1}{2}\phi_i\sigma_i} \quad (3-70)$$

This reveals that  $(\Lambda^{(L)})^\dagger$  is the inverse of  $\Lambda^{(R)}$ . Similarly,  $(\Lambda^{(R)})^\dagger$  is the inverse of  $\Lambda^{(L)}$ .

Thus,  $\mathcal{L}^\dagger\mathcal{R}$  and  $\mathcal{R}^\dagger\mathcal{L}$  are Lorentz invariants, which can be seen below,

$$(\mathcal{L}^\dagger\mathcal{R})' = \mathcal{L}^\dagger(\Lambda^{(L)})^\dagger\Lambda^{(R)}\mathcal{R} = \mathcal{L}^\dagger(\Lambda^{(R)})^{-1}\Lambda^{(R)}\mathcal{R} = \mathcal{L}^\dagger\mathcal{R} \quad (3-71)$$

$$(\mathcal{R}^\dagger\mathcal{L})' = \mathcal{R}^\dagger(\Lambda^{(R)})^\dagger\Lambda^{(L)}\mathcal{L} = \mathcal{R}^\dagger(\Lambda^{(L)})^{-1}\Lambda^{(L)}\mathcal{L} = \mathcal{R}^\dagger\mathcal{L}. \quad (3-72)$$

These are the lowest order invariant pieces involving the field alone. In order to couple the derivative to the field the  $\mathcal{L}$  and  $\mathcal{R}$  spinors must attach to  $\partial_\mu$  in an invariant way. Recall that a four vector may be expressed as an outer product of left and right spinors,

$$\alpha_{L\tilde{R}} = \mathcal{L}'\tilde{\mathcal{R}}'^T. \quad (3-73)$$

The following also works,

$$\alpha_{R\tilde{L}} = \mathcal{R}'\tilde{\mathcal{L}}'^T. \quad (3-74)$$

Recall that the  $\sim$  transformation sent  $\sigma_i$  to  $-\sigma_i^*$ . So the transformations for the  $\sim$  transposed spinors are,

$$(\Lambda^{(\tilde{L})})^T = e^{\frac{i}{2}\theta_i(-\sigma_i^*)^T + \frac{1}{2}\phi_i(-\sigma_i^*)^T} = e^{-\frac{i}{2}\theta_i\sigma_i^\dagger - \frac{1}{2}\phi_i\sigma_i^\dagger} = e^{-\frac{i}{2}\theta_i\sigma_i - \frac{1}{2}\phi_i\sigma_i} \quad (3-75)$$

and

$$(\Lambda^{(\tilde{R})})^T = e^{\frac{i}{2}\theta_i(-\sigma_i^*)^T - \frac{1}{2}\phi_i(-\sigma_i^*)^T} = e^{-\frac{i}{2}\theta_i\sigma_i^\dagger + \frac{1}{2}\phi_i\sigma_i^\dagger} = e^{-\frac{i}{2}\theta_i\sigma_i + \frac{1}{2}\phi_i\sigma_i}. \quad (3-76)$$

The equations above show that the  $(\Lambda^{(\tilde{L})})^T$  transformation is the inverse of  $\Lambda^{(L)}$ , and that the  $(\Lambda^{(\tilde{R})})^T$  transformation is the inverse of  $\Lambda^{(R)}$ . So the invariant pieces coupling the spinor to the four vector are,

$$\mathcal{R}^\dagger \alpha_{L\tilde{R}} \mathcal{R} = \mathcal{R}^\dagger \mathcal{L}' \tilde{\mathcal{R}}'^T \mathcal{R} \quad (3-77)$$

and

$$\mathcal{L}^\dagger \alpha_{R\tilde{L}} \mathcal{L} = \mathcal{L}^\dagger \mathcal{R}' \tilde{\mathcal{L}}'^T \mathcal{L}. \quad (3-78)$$

The two types of four vectors have slightly different transformations

$$\alpha'_{L\tilde{R}} = e^{\frac{i}{2}\theta_i\sigma_i + \frac{1}{2}\phi_i\sigma_i} \alpha_{L\tilde{R}} e^{-\frac{i}{2}\theta_i\sigma_i + \frac{1}{2}\phi_i\sigma_i} \quad (3-79)$$

$$\alpha'_{R\tilde{L}} = e^{\frac{i}{2}\theta_i\sigma_i - \frac{1}{2}\phi_i\sigma_i} \alpha_{R\tilde{L}} e^{-\frac{i}{2}\theta_i\sigma_i - \frac{1}{2}\phi_i\sigma_i}. \quad (3-80)$$

Both transform with positive  $\theta$  under rotations, but the boosts are a different story. The  $L\tilde{R}$  four vector transforms with positive  $\phi$  while the  $R\tilde{L}$  four vector transforms with negative  $\phi$ . Behaving the same way under rotations implies that the x,y,z components mix up the same way in both types of four vector. In this respect, rotating x,y,z or -x,-y,-z are both valid. Exemplifying this, an infinitesimal rotation around z by  $d\theta_z$  gives



$$\begin{aligned}x' &= x - y d\theta_z \\ y' &= y + x d\theta_z\end{aligned}\tag{3-81}$$

for positive  $x, y, z$  and

$$\begin{aligned}-x' &= -x - (-y) d\theta_z \\ -y' &= -y' + (-x) d\theta_z\end{aligned}\tag{3-82}$$

for  $-x, -y, -z$ . These transformations are equivalent. Moving onto boosts, the opposite  $\phi$  sign naively implies that the  $x, y, z$  terms mix up with  $t$  in the opposite way for the two types of rank 2 spinors. However, if both  $L\tilde{R}$  and  $R\tilde{L}$  represent a four vector, then the boost must yield the same transformation on the components. Therefore, to get from  $L\tilde{R}$  to  $R\tilde{L}$  take  $t, x, y, z$  to  $t, -x, -y, -z$ . Illuminating this, an infinitesimal boost along  $x$  by  $d\phi_x$  gives

$$\begin{aligned}t' &= t + x d\phi_x \\ x' &= x + t d\phi_x\end{aligned}\tag{3-83}$$

for the  $L\tilde{R}$  representation and

$$\begin{aligned}t' &= t + (-x)(-d\phi_x) \\ -x' &= -x + t(-d\phi_x)\end{aligned}\tag{3-84}$$

for  $R\tilde{L}$ . The negative  $\phi$  and the negative spatial components counteract to transform the four vector correctly. The  $R\tilde{L}$  rank 2 spinor encodes a four vector in the same way as a  $L\tilde{R}$  rank 2 spinor, except that the spatial components are coded into the rank 2 spinor with the opposite sign. Writing the representations out in terms of the four vector components yields

$$\begin{aligned}
\alpha_{\tilde{L}\tilde{R}} &= \alpha_0\sigma_0 + \alpha_1\sigma_1 + \alpha_2\sigma_2 + \alpha_3\sigma_3 \\
\alpha_{\tilde{L}\tilde{R}} &= \alpha_\mu\sigma^\mu \\
\alpha_{\tilde{L}\tilde{R}} &= \begin{pmatrix} \alpha_0 + \alpha_3 & \alpha_1 - i\alpha_2 \\ \alpha_1 + i\alpha_2 & \alpha_0 - \alpha_3 \end{pmatrix}
\end{aligned} \tag{3-85}$$

and

$$\begin{aligned}
\alpha_{\tilde{R}\tilde{L}} &= \alpha_0\sigma_0 - \alpha_1\sigma_1 - \alpha_2\sigma_2 - \alpha_3\sigma_3 \\
\alpha_{\tilde{R}\tilde{L}} &= \alpha_\mu\bar{\sigma}^\mu \\
\alpha_{\tilde{R}\tilde{L}} &= \begin{pmatrix} \alpha_0 - \alpha_3 & \alpha_1 + i\alpha_2 \\ \alpha_1 - i\alpha_2 & \alpha_0 + \alpha_3 \end{pmatrix}.
\end{aligned} \tag{3-86}$$

The action for the spin  $\frac{1}{2}$  particle is built from the lowest order invariant combinations of the spinors themselves and the lowest order invariant term coupling  $\partial_\mu$  to the spinors. These ingredients are  $\mathcal{L}^\dagger\mathcal{R}$ ,  $\mathcal{R}^\dagger\mathcal{L}$ ,  $\mathcal{R}^\dagger\sigma^\mu\partial_\mu\mathcal{R}$ , and  $\mathcal{L}^\dagger\bar{\sigma}^\mu\partial_\mu\mathcal{L}$ . In the scalar particle case m coupled the field to itself as potential energy <sup>11</sup> with the same power as  $\partial_\mu$  <sup>12</sup>. Following that example and throwing an i into the derivative terms<sup>13</sup> to make them Hermitian gives the following Lagrangian,

$$S = \int d^4x \left( \mathcal{R}^\dagger\sigma^\mu i\partial_\mu\mathcal{R} + i\mathcal{L}^\dagger\bar{\sigma}^\mu i\partial_\mu\mathcal{L} - m\mathcal{L}^\dagger\mathcal{R} - m\mathcal{R}^\dagger\mathcal{L} \right). \tag{3-87}$$

The Euler-Lagrange equations reveal how the fields change over time,

---

<sup>11</sup> The mass term had a negative sign in the Lagrangian.

<sup>12</sup> E, P, and m must have the same power to get  $E^2 = m^2 + \vec{P}^2$  right, and  $E \sim \partial_0$ ,  $P_i \sim \partial_i$  so m must have the same power as  $\partial_\mu$ .

<sup>13</sup> This gives the  $P_\mu$  operator which equals  $i\partial_\mu$ .

$$\begin{aligned}
\sigma^\mu i \partial_\mu \mathcal{R} &= m \mathcal{L} \\
\bar{\sigma}^\mu i \partial_\mu \mathcal{L} &= m \mathcal{R}. \\
\sigma^\mu i \partial_\mu \mathcal{R}^\dagger &= -m \mathcal{L}^\dagger \\
\bar{\sigma}^\mu i \partial_\mu \mathcal{L}^\dagger &= -m \mathcal{R}^\dagger.
\end{aligned} \tag{3-88}$$

Notice that the mass term couples the left and right chiral spinors. At rest,  $P=0$  and the equations reduce to  $i \partial_0 \mathcal{R} = m \mathcal{L}$  and  $i \partial_0 \mathcal{L} = m \mathcal{R}$  showing that  $\partial_0^2 \mathcal{R} = -m^2 \mathcal{R}$  and  $\partial_0^2 \mathcal{L} = -m^2 \mathcal{L}$ . The mass is actually a frequency determining how quickly a particle oscillates between its right-chiral and left-chiral states. Another interesting point is that swapping the sign of  $m$  here swaps the roles of the fields and the conjugate fields.

Plugging  $\mathcal{R} = \frac{1}{m} \bar{\sigma}^\nu i \partial_\nu \mathcal{L}$  into  $\sigma^\mu i \partial_\mu \mathcal{R} = m \mathcal{L}$  shows that the constants chosen for the terms in the Lagrangian provide the correct dispersion relation,

$$\begin{aligned}
\sigma^\mu i \partial_\mu \frac{1}{m} \bar{\sigma}^\nu i \partial_\nu \mathcal{L} &= m \mathcal{L} \\
\rightarrow \sigma^\mu \bar{\sigma}^\nu P_\mu P_\nu &= m^2 \\
\rightarrow \eta_\nu^\mu P_\mu P_\nu &= m^2 \\
\rightarrow P^\mu P_\mu &= m^2 \\
\rightarrow E^2 - \vec{P}^2 &= m^2
\end{aligned} \tag{3-89}$$

The action for the spin  $\frac{1}{2}$  particle can be rewritten into its more compact, canonical form,

$$S = \int d^4x \bar{\psi} (i \gamma^\mu \partial_\mu - m) \psi, \tag{3-90}$$

after defining

$$\gamma^\mu = \begin{pmatrix} 0 & \sigma^\mu \\ \bar{\sigma}^\mu & 0 \end{pmatrix}, \psi = \begin{pmatrix} \mathcal{L} \\ \mathcal{R} \end{pmatrix}, \text{ and } \bar{\psi} = \psi^\dagger \gamma_0. \tag{3-91}$$

The spin  $\frac{1}{2}$  Lagrangian is called the Dirac Lagrangian and the resulting equations of motion are the Dirac equation.

### 3.1.3.6 Building the Lagrangian for a Free Spin 1 Particle

Last but not least is the Lagrangian for a spin 1 force carrying particle. The main ingredients are the  $A_\mu$  and  $\partial_\mu$  four vectors. The lowest order possible invariants are  $\partial^\mu A^\nu \partial_\mu A_\nu$ ,  $\partial^\mu A^\nu \partial_\nu A_\mu$ ,  $A^\mu A_\mu$ , and  $\partial^\mu A_\mu$ . However the last term is a total derivative and won't be included. The resulting action and equations of motion are

$$S = \int d^4x (c_0 \partial^\mu A^\nu \partial_\mu A_\nu + c_1 \partial^\mu A^\nu \partial_\nu A_\mu + c_2 A^\mu A_\mu). \quad (3-92)$$

$$c_2 A^\nu = -\partial_\mu (c_0 \partial^\mu A^\nu + c_1 \partial^\nu A^\mu). \quad (3-93)$$

This is the same form as the equation for the four vector potential in electromagnetism, which has  $c_0 = \frac{1}{2}$  and  $c_1 = -\frac{1}{2}$ . The remaining term,  $c_2$  looks like a mass term, so  $c_2$  is set to  $\frac{1}{2}m^2$ <sup>14</sup>. The action and equations of motion with the appropriate constants become,

$$S = \int d^4x \left( \frac{-1}{2} \partial^\mu A^\nu \partial_\mu A_\nu + \frac{1}{2} \partial^\mu A^\nu \partial_\nu A_\mu + \frac{m^2}{2} A^\mu A_\mu \right) \quad (3-94)$$

$$-m^2 A^\nu = \partial_\mu (\partial^\mu A^\nu - \partial^\nu A^\mu). \quad (3-95)$$

Taking another derivative reveals that the equations reduce further and take on the correct dispersion relation, confirming the choice of constants,

---

<sup>14</sup> The order of the mass should match the order of the derivatives.

$$\begin{aligned}
-m^2 \partial_\nu A^\nu &= \partial_\nu \partial_\mu \partial^\mu A^\nu - \partial_\nu \partial_\mu \partial^\nu A^\mu = 0 \\
\rightarrow \partial_\nu A^\nu &= 0 \\
\rightarrow -m^2 A^\nu &= \partial_\mu \partial^\mu A^\nu \\
\rightarrow m^2 A^\nu &= i \partial_\mu i \partial^\mu A^\nu \\
\rightarrow m^2 &= E^2 - \vec{P}^2.
\end{aligned} \tag{3-96}$$

The action for the spin 1 particle is called the Proca action. It's normally written in an equivalent form in terms of the tensor  $F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu$

$$S = \int d^4x \left( \frac{-1}{4} F_{\mu\nu} F^{\mu\nu} + \frac{m^2}{2} A^\mu A_\mu \right). \tag{3-97}$$

### 3.1.4 Lagrangians in Quantum Mechanics and QFT

Quantum behavior can only be predicted in a strange probabilistic way. Classically, each particle takes a single well defined path, but in the quantum regime particles behave as if they take all possible paths from some initial point to some final point. Analogously, the fields from which particles arise behave as if they take all possible configurations between states. Each possibility contributes as a unit complex number with a phase given by the action. These contributions are called probability amplitudes.

$$\mathcal{A}_j = e^{iS_j} \tag{3-98}$$

Adding the amplitudes for every possibility gives the net amplitude, and squaring the net amplitude gives the probability.

$$P = \left| \sum_j \mathcal{A}_j \right|^2 \tag{3-99}$$

Consider a free non-relativistic quantum particle in one dimension with the Lagrangian  $\mathcal{L} = \frac{1}{2} m \dot{x}^2$ . The amplitude for the particle to start at  $x_0, t_0$  and end at  $x_f, t_f$  can be calculated

by discretizing time,  $t \in t_0, t_1, \dots, t_f$ . A path is then determined by specifying  $x$  at each point in time,  $x(t) \in x(t_0), x(t_1), \dots, x(t_f)$ . In this way, the action for a path  $x$ ,  $S(x)$ , can be written

$$S(x) = \int dt L = \frac{m}{2} \int_{t_0}^{t_f} dt \dot{x}^2 = \frac{m}{2} \sum_i \frac{(x(t_i) - x(t_{i-1}))^2}{(t_i - t_{i-1})^2} (t_i - t_{i-1}) = \frac{m}{2} \sum_i \frac{(x(t_i) - x(t_{i-1}))^2}{\Delta t}. \quad (3-100)$$

The amplitude for the path is then given by,

$$\mathcal{A}(x) = e^{i\frac{m}{2} \frac{(x(t_f) - x(t_{f-1}))^2}{\Delta t}} e^{i\frac{m}{2} \frac{(x(t_{f-1}) - x(t_{f-2}))^2}{\Delta t}} \dots e^{i\frac{m}{2} \frac{(x(t_1) - x(t_0))^2}{\Delta t}}. \quad (3-101)$$

Summing over all paths,  $x(t)$ , with fixed endpoints,  $x(t_0) = x_0$  and  $x(t_f) = x_f$ , provides the net amplitude

$$\begin{aligned} \mathcal{A} &= \sum_x \mathcal{A}(x) = \sum_{x(t_{f-1})} \sum_{x(t_{f-2})} \dots \sum_{x(t_1)} e^{i\frac{m}{2} \frac{(x(t_f) - x(t_{f-1}))^2}{\Delta t}} e^{i\frac{m}{2} \frac{(x(t_{f-1}) - x(t_{f-2}))^2}{\Delta t}} \dots e^{i\frac{m}{2} \frac{(x(t_1) - x(t_0))^2}{\Delta t}} \\ &= \mathcal{N} \int dx_{t_{f-1}} dx_{t_{f-2}} \dots dx_{t_1} e^{i\frac{m}{2} \frac{(x(t_f) - x(t_{f-1}))^2}{\Delta t}} e^{i\frac{m}{2} \frac{(x(t_{f-1}) - x(t_{f-2}))^2}{\Delta t}} \dots e^{i\frac{m}{2} \frac{(x(t_1) - x(t_0))^2}{\Delta t}} \\ &= \int \mathcal{D}[x(t)] e^{i\frac{m}{2} \int_{t_0}^{t_f} dt \partial_t x \partial_t x}. \end{aligned} \quad (3-102)$$

Integrating by parts yields,

$$\int \mathcal{D}[x(t)] e^{i\frac{m}{2} \int_{t_0}^{t_f} dt (x \partial_t^2 x)}. \quad (3-103)$$

In general the amplitude to go from  $\phi(t_0) = \phi_0$  to  $\phi(t_f) = \phi_f$  is determined by the path integral,

$$\mathcal{A}(\phi_f, t_f; \phi_0, t_0) = \int \mathcal{D}[\phi(t)] e^{i \int_{t_0}^{t_f} dt L}. \quad (3-104)$$




Figure 3-2. The Feynman diagram for two gluons fusing into a Higgs. There are three vertices and this is a third order diagram. Two vertices involve the strong force and one vertex involves the Higgs coupling. The matrix element for this diagram would have two factors of the strong force coupling and one factor for the Higgs coupling which involves the mass of the top quark.

### 3.1.5 The Higgs Mechanism

### 3.1.6 Perturbation Theory and Feynman Rules

## 3.2 The Standard Model Higgs

### 3.2.1 SM Higgs Production and Decay Modes

The SM Higgs can be created in a variety of ways. Some of these cross sections are shown below for 14 TeV collisions. The production cross sections are functions of the mass of the Higgs as well as the energy of the collisions. For a given collision energy the cross sections decrease as the Higgs mass increases: there are fewer kinematic possibilities for a heavier particle since more of the energy was used to create the particle. For a given mass, say 125 GeV, the cross section grows with collision energy. This contrasts with cross sections involving collisions of fundamental particles, e.g. electron antielectron collisions. This is an artifact of the fact that the LHC collides protons together.

Protons behave like a collection of an infinite number of quark-antiquarks, an infinite number of gluons, and the usual uud. The total momentum of the proton is divided up amongst them with lots of particles having little of the total momentum. The actual scattering events are between these more fundamental particles. The larger the total energy the smaller the fraction of energy needed to make a Higgs and since there are more particles with a lower fraction, this results in a growth of the cross section with collision energy.

The SM Higgs is unstable and decays with a width of  $\sim 5\text{MeV}$  at 125 GeV. The probability of each decay changes depending upon the mass of the Higgs. In general the Higgs




Figure 3-3. The highest production mode cross sections for the SM Higgs ?

couples more strongly to particles with higher mass making the decays to heavier particles more likely.

The muon has the lowest mass – excluding the photon and gluon – of the particles in Figure 3-4 and consequently  $H \rightarrow \mu^+ \mu^-$  has the lowest branching fraction in the set.<sup>15</sup> The gluons and photons are massless and do not couple to the Higgs at leading order. These massless vector bosons interact with the Higgs through a loop of top quarks. The extremely heavy mass of the top quark, about 173 GeV, balances the fact that the loop production is a higher order mechanism.

The Higgs to massless vector boson coupling via the top loop is seen in the GF Feynman diagram in Figure 3-5. At  $M_h = 125$  GeV,  $\sqrt{s} = 13$  TeV, the GF channel comprises 87% of the total Higgs production cross section, VBF 7%, VH 4%, and  $t\bar{t}H$  1% ?. Besides  $t\bar{t}H$ , the

<sup>15</sup> The Higgs also couples to the electron and the first generation quarks but the masses are so light that CMS does not expect to see the SM Higgs in those modes.






Figure 3-4. The graphic on the top left presents the SM Higgs branching fractions as functions of mass while the table on the bottom right displays the branching fractions for a 125 GeV SM Higgs ?.




Figure 3-5. The SM production modes with the highest cross sections. a) Gluon Gluon Fusion (GF) b) Vector Boson Fusion (VBF) c) Associated Production with a Vector Boson (VH) d)  $t\bar{t}H$

process  $q + \bar{q} \rightarrow H$  isn't considered due to its low cross section. The low masses of the other quarks suppress the process.

Quark gluon (qg) scattering is a major background for the Higgs to two jets decays since the process closely resembles GF in this mode.




Figure 3-6. Quark gluon scattering creates many two jet events. This background looks very similar to GF when the Higgs decays to two jets. The colliding protons are made of quarks and gluons so this process is extremely common.

## CHAPTER 4 RESULTS

### 4.1 Fusce Eget Tempus Lectus,

**Algorithm 4.1.** *Euclids algorithm*

1: <b>procedure</b> EUCLID( $a, b$ )	▷ <i>The g.c.d. of <math>a</math> and <math>b</math></i>
2: $r \leftarrow a \bmod b$	
3: <b>while</b> $r \neq 0$ <b>do</b>	▷ <i>We have the answer if <math>r</math> is 0</i>
4: $a \leftarrow b$	
5: $b \leftarrow r$	
6: $r \leftarrow a \bmod b$	
7: <b>end while</b>	
8: <b>return</b> $b$	▷ <i>The gcd is <math>b</math></i>
9: <b>end procedure</b>	

**Proposition 4.1.** *The Upsilon Function*

(1) *If  $\beta > 0$  and  $\alpha \neq 0$ , then for all  $n \geq -1$ ,*

$$I_n(c; \alpha; \beta; \delta) = -\frac{e^{\alpha c}}{\alpha} \sum_{i=0}^n \left(\frac{\beta}{\alpha}\right)^{n-i} \text{Hh}_i(\beta c - \delta) \\ + \left(\frac{\beta}{\alpha}\right)^{n+1} \frac{\sqrt{2\pi}}{\beta} e^{\frac{\alpha\delta}{\beta} + \frac{\alpha^2}{2\beta^2}} \phi\left(-\beta c + \delta + \frac{\alpha}{\beta}\right)$$

(2) *If  $\beta < 0$  and  $\alpha < 0$ , then for all  $x \geq -1$*

$$I_n(c; \alpha; \beta; \delta) = -\frac{e^{\alpha c}}{\alpha} \sum_{i=0}^n \left(\frac{\beta}{\alpha}\right)^{n-i} \text{Hh}_i(\beta c - \delta) \\ - \left(\frac{\beta}{\alpha}\right)^{n+1} \frac{\sqrt{2\pi}}{\beta} e^{\frac{\alpha\delta}{\beta} + \frac{\alpha^2}{2\beta^2}} \phi\left(\beta c - \delta - \frac{\alpha}{\beta}\right)$$

*Proof.* Case 1.

$\beta > 0$  and  $\alpha \neq 0$ . Since, for any constant  $\alpha$  and  $n \geq 0$ ,  $e^{\alpha x} \text{Hh}_n(\beta x - \delta) \rightarrow 0$  as  $x \rightarrow \infty$

thanks to (B4), integration by parts leads to

$$I_n = -\frac{1}{\alpha} \text{Hh}(\beta c - \delta) e^{\alpha c} + \frac{\beta}{\alpha} \int_c^\infty e^{\alpha x} \text{Hh}_{n-1}(\beta x - \delta) dx$$

In other words, we have a recursion, for  $n \geq 0$ ,  $I_n = -(e^{\alpha c} \alpha) \text{Hh}_n(\beta c - \delta) + (\frac{\beta}{\alpha}) I_{n-1}$  with

$$\begin{aligned} I_{-1} &= \sqrt{2\pi} \int_c^\infty e^{\alpha x} \varphi(-\beta x + \delta) dx \\ &= \frac{\sqrt{2\pi}}{\beta} e^{\frac{\alpha \delta}{\beta} + \frac{\alpha^2}{2\beta^2}} \phi(-\beta c + \delta + \frac{\alpha}{\beta}) \end{aligned}$$

Solving it yields, for  $n \geq -1$ ,

$$\begin{aligned} I_n &= -\frac{e^{\alpha c}}{\alpha} \sum_{i=0}^n \left(\frac{\beta}{\alpha}\right)^i \text{Hh}_{n-i}(\beta c + \delta) + \left(\frac{\beta}{\alpha}\right)^{n+1} I_{-1} \\ &= -\frac{e^{\alpha c}}{\alpha} \sum_{i=0}^n \left(\frac{\beta}{\alpha}\right)^{n-i} \text{Hh}_i(\beta c + \delta) \\ &\quad + \left(\frac{\beta}{\alpha}\right)^{n+1} \frac{\sqrt{2\pi}}{\beta} e^{\frac{\alpha \delta}{\beta} + \frac{\alpha^2}{2\beta^2}} \phi(-\beta c + \delta + \frac{\alpha}{\beta}) \end{aligned}$$

where the sum over an empty set is defined to be zero. □

*Proof. Case2.*  $\beta < 0$  and  $\alpha < 0$ . In this case, we must also have, for  $n \geq 0$  and any constant  $\alpha < 0$ ,  $e^{\alpha x} \text{Hh}_n(\beta x - \delta) \rightarrow 0$  as

$x \rightarrow \infty$ , thanks to (B5). Using integration by parts, we again have the same recursion, for  $n \geq 0$ ,  $I_n = -(e^{\alpha c} / \alpha) \text{Hh}_n(\beta c - \delta) + (\beta / \alpha) I_{n-1}$ , but with a different initial condition

$$\begin{aligned} I_{-1} &= \sqrt{2\pi} \int_c^\infty e^{\alpha x} \varphi(-\beta x + \delta) dx \\ &= -\frac{\sqrt{2\pi}}{\beta} \exp\left\{\frac{\alpha \delta}{\beta} + \frac{\alpha^2}{2\beta^2}\right\} \phi(\beta c - \delta - \frac{\alpha}{\beta}) \end{aligned}$$

Solving it yields (B8), for  $n \geq -1$ . □

*Finally, we sum the double exponential and the normal random variables*

*Proposition B.3.*

Suppose  $\{\xi_1, \xi_2, \dots\}$  is a sequence of i.i.d. exponential random variables with rate  $\eta > 0$ , and  $Z$  is a normal variable with distribution  $N(0, \sigma^2)$ . Then for every  $n \geq 1$ , we have: (1) The density functions are given by:

$$f_{Z+\sum_{i=1}^n \xi_i}(t) = (\sigma\eta)^n \frac{e^{(\sigma\eta)^2/2}}{\sigma\sqrt{2\pi}} e^{-t\eta} \text{Hh}_{n-1}\left(-\frac{t}{\sigma} + \sigma\eta\right)$$

$$f_{Z-\sum_{i=1}^n \xi_i}(t) = (\sigma\eta)^n \frac{e^{(\sigma\eta)^2/2}}{\sigma\sqrt{2\pi}} e^{-t\eta} \text{Hh}_{n-1}\left(\frac{t}{\sigma} + \sigma\eta\right)$$

(2) The tail probabilities are given by

$$P(Z + \sum_{i=1}^n \xi_i \geq x) = (\sigma\eta)^n \frac{e^{(\sigma\eta)^2/2}}{\sigma\sqrt{2\pi}} e^{-t\eta} I_{n-1}\left(x; -\eta, -\frac{1}{\sigma}, -\sigma\eta\right)$$

$$P(Z - \sum_{i=1}^n \xi_i \geq x) = (\sigma\eta)^n \frac{e^{(\sigma\eta)^2/2}}{\sigma\sqrt{2\pi}} e^{-t\eta} I_{n-1}\left(x; \eta, \frac{1}{\sigma}, -\sigma\eta\right)$$

*Proof. Case 1. The densities of  $Z + \sum_{i=1}^n \xi_i$ , and  $Z - \sum_{i=1}^n \xi_i$ . We have*

$$\begin{aligned} f_{Z+\sum_{i=1}^n \xi_i}(t) &= \int_{-\infty}^{\infty} f_{\sum_{i=1}^n \xi_i}(t-x) f_Z(x) dx \\ &= e^{-t\eta} (\eta^n) \int_{-\infty}^{\infty} t \frac{e^{x\eta} (t-x)^{n-1}}{(n-1)!} \frac{1}{\sigma\sqrt{2\pi}} e^{-x^2/(2\sigma^2)} dx \\ &= e^{-t\eta} (\eta^n) e^{(\sigma\eta)^2/(2)} \int_{-\infty}^{\infty} t \frac{(t-x)^{n-1}}{(n-1)!} \frac{1}{\sigma\sqrt{2\pi}} e^{-(x-\sigma^2\eta)^2/(2\sigma^2)} dx \end{aligned}$$

*Letting  $y = (x - \sigma^2\eta)/\sigma$  yields*

$$\begin{aligned} f_{Z+\sum_{i=1}^n \xi_i}(t) &= e^{-t\eta} (\eta^n) e^{(\sigma\eta)^2/(2)} \sigma^{n-1} \\ &\times \int_{-\infty}^{t/\sigma - \sigma\eta} \frac{(t/\sigma - y - \sigma\eta)^{n-1}}{(n-1)!} \frac{1}{\sqrt{2\pi}} e^{-y^2/2} dy \end{aligned}$$

$$= \frac{e^{(\sigma\eta)^2/2}}{\sqrt{2\pi}} (\sigma^{n-1} \eta^n) e^{-t\eta} \text{Hh}_{n-1}(-t/\sigma + \sigma\eta)$$

because  $(1/(n-1)! \int_{-\infty}^{\infty} a(a-y)^{n-1} e^{-y^2/2} dy = \text{Hh}_{n-1}(a)$ . The derivation of  $f_{Z+\sum_{i=1}^n \xi_i}(t)$  is similar.

Case 2.  $P(Z + \sum_{i=1}^n \xi_i \geq x)$  and  $P(Z - \sum_{i=1}^n \xi_i \geq x)$ . From (B9), it is clear that

$$\begin{aligned} P(Z + \sum_{i=1}^n \xi_i \geq x) &= \frac{(\sigma\eta)^n e^{(\sigma\eta)^2/2}}{\sigma\sqrt{2\pi}} \int_x^{\infty} e^{(-i\eta)} \text{Hh}_{n-1}\left(-\frac{t}{\sigma} + \sigma\eta\right) dt \\ &= \frac{(\sigma\eta)^n e^{(\sigma\eta)^2/2}}{\sigma\sqrt{2\pi}} I_{n-1}\left(x; -\eta, -\frac{1}{\sigma}, -\sigma\eta\right) \end{aligned}$$

by (B6). We can compute  $P(Z - \sum_{i=1}^n \xi_i \geq x)$  similarly.

**Theorem 4.1.** Theorem With  $\pi_n := P(N(t) = n) = e^{-\lambda T} (\lambda T)^n / n!$  and  $I_n$  in Proposition ?? , we have

$$\begin{aligned} P(Z(T) \geq a) &= \frac{e^{(\sigma\eta_1)^2 T/2}}{\sigma\sqrt{2\pi T}} \sum_{n=1}^{\infty} \pi_n \sum_{k=1}^n P_{n,k}(\sigma\sqrt{T}\eta_1)^k \times I_{k-1}\left(a - \mu T; -\eta_1, -\frac{1}{\sigma\sqrt{T}}, -\sigma\eta_1\sqrt{T}\right) \\ &\quad + \frac{e^{(\sigma\eta_2)^2 T/2}}{\sigma\sqrt{2\pi T}} \sum_{n=1}^{\infty} \pi_n \sum_{k=1}^n Q_{n,k}(\sigma\sqrt{T}\eta_2)^k \\ &\quad \times I_{k-1}\left(a - \mu T; \eta_2, \frac{1}{\sigma\sqrt{T}}, -\sigma\eta_2\sqrt{T}\right) \\ &\quad + \pi_0 \phi\left(-\frac{a - \mu T}{\sigma\sqrt{T}}\right) \end{aligned}$$

Proof by the decomposition (B2)

$$P(Z(T) \geq a) = \sum_{n=0}^{\infty} \pi_n P(\mu T + \sigma\sqrt{T}Z + \sum_{j=1}^n Y_j \geq a)$$

$$\begin{aligned}
&= \pi_0 P(\mu T + \sigma \sqrt{T} Z \geq a) \\
&+ \sum_{n=1}^{\infty} \pi_n \sum_{k=1}^n P_{n,k} P(\mu T + \sigma \sqrt{T} Z + \sum_{j=1}^n \xi_j^+ \geq a) \\
&+ \sum_{n=1}^{\infty} \pi_n \sum_{k=1}^n Q_{n,k} P(\mu T + \sigma \sqrt{T} Z - \sum_{j=1}^n \xi_j^- \geq a)
\end{aligned}$$

*The result now follows via (B11) and (B12) for  $\eta_1 > 1$  and  $\eta_2 > 0$ .*

## CHAPTER 5 SUMMARY AND CONCLUSIONS

### 5.1 Non Porttitor Tellus

Aliquam molestie sed urna quis convallis. Aenean nibh eros, aliquam non eros in, tempus lacinia justo. In magna sapien, blandit a faucibus ac, scelerisque nec purus. Praesent fermentum felis nec massa interdum, vel dapibus mi luctus. Cras id fringilla mauris. Ut molestie eros mi, ut hendrerit nulla tempor et. Pellentesque tortor quam, mattis a scelerisque nec, euismod et odio. Mauris rhoncus metus sit amet risus mattis, eu mattis sem interdum.

#### 5.1.1 Nam Arcu Magna

Semper vel lorem eu, venenatis ultrices est. Nam aliquet ut erat ac scelerisque. Maecenas ut molestie mi. Phasellus ipsum magna, sollicitudin eu ipsum quis, imperdiet cursus turpis. Etiam pretium enim a fermentum accumsan. Morbi vel vehicula enim.

##### 5.1.1.1 Ut pellentesque velit sede

Placerat cursus. Integer congue urna non massa dictum, a pellentesque arcu accumsan. Nulla posuere, elit accumsan eleifend elementum, ipsum massa tristique metus, in ornare neque nisl sed odio. Nullam eget elementum nisi. Duis a consectetur erat, sit amet malesuada sapien. Aliquam nec sapien et leo sagittis porttitor at ut lacus. Vivamus vulputate elit vitae libero condimentum dictum. Nulla facilisi. Quisque non nibh et massa ullamcorper iaculis.



## APPENDIX A

### THIS IS THE FIRST APPENDIX

Lorem ipsum dolor sit amet, consectetur adipiscing elit. Maecenas eget magna. Aenean et lorem. Ut dignissim neque at nisi. In hac habitasse platea dictumst. In porta ornare eros. Nunc eu ante. In non est vehicula tellus cursus suscipit. Proin sed libero. Sed risus enim, eleifend in, pellentesque ac, nonummy quis, nulla. Phasellus imperdiet libero nec massa. Ut sapien libero, adipiscing eu, volutpat porttitor, ultricies eget, nisi. Sed odio. Suspendisse potenti. Duis dolor augue, viverra id, porta in, dignissim id, nisl. Vivamus blandit cursus eros. Maecenas sit amet urna sit amet orci nonummy pharetra.

Praesent cursus nibh et mauris. In aliquam felis sit amet ligula. Nulla faucibus nisl eget nisl. Aliquam tincidunt. Mauris eget elit sed massa luctus posuere. Pellentesque suscipit. In odio urna, semper ut, convallis ut, porta et, nibh. Nulla sodales metus nec velit posuere gravida. Cras tristique. Etiam urna risus, accumsan ut, placerat sed, iaculis id, est.

Nullam mi. Pellentesque habitant morbi tristique senectus et netus et malesuada fames ac turpis egestas. Duis vitae metus in massa hendrerit rhoncus. Fusce tortor justo, laoreet eu, facilisis at, gravida et, felis. Donec imperdiet mollis erat. Integer tempus nulla ac lorem. Fusce porttitor. Aenean quis arcu. Morbi consectetur, leo eu mollis elementum, urna massa malesuada risus, euismod tempor lorem elit ut mauris. Cras elit orci, facilisis ac, mattis iaculis, cursus ac, augue. Donec eget nisl. Pellentesque fermentum sodales nibh. Vivamus non risus. Donec est libero, tincidunt sit amet, pretium vitae, blandit sed, tellus. Nunc diam risus, interdum sed, laoreet quis, varius ac, turpis. In et purus eget nibh vehicula rhoncus. Aenean et neque. Praesent nisl nisi, tempus quis, nonummy ac, auctor a, neque. Suspendisse et metus. Suspendisse non metus eu mauris auctor sagittis.

APPENDIX B  
AN EXAMPLE OF A HALF TITLE PAGE

L<sup>A</sup>T<sub>E</sub>X 2<sub>ε</sub>

Figure B-1. L<sup>A</sup>T<sub>E</sub>X 2<sub>ε</sub>. logo

This is how a section should look if the first page is a landscape page. Lorem ipsum dolor sit amet, consectetur adipiscing elit. Ut sit amet nulla. Integer mauris turpis, dapibus ac, auctor non, vehicula sit amet, magna. Suspendisse eu tellus. Etiam porta. Donec magna. Donec ut dui. In hac habitasse platea dictumst. Nullam suscipit, mi at adipiscing commodo, lorem erat scelerisque erat, non pulvinar leo mi eu metus. Phasellus id felis. Sed quam purus, molestie quis, ultrices nec, dictum at, magna. Proin viverra viverra ante.

Maecenas sagittis magna quis ligula. Duis vestibulum mi a felis. Aenean accumsan mattis massa. Nullam lacus sem, consectetur non, condimentum sit amet, pharetra ac, odio. Morbi nisi magna, tincidunt sed, placerat nec, tincidunt id, lectus. Donec ac dui non mauris vulputate aliquam. Nullam scelerisque congue pede. Integer ipsum. Vestibulum auctor. Suspendisse eget leo id libero cursus dictum. Sed malesuada. Aliquam imperdiet. Donec dui metus, porta eu, aliquet vel, vulputate vitae, lacus.

Nulla quis purus id turpis luctus feugiat. Fusce feugiat. Proin felis. Morbi elit est, fermentum in, tincidunt vitae, convallis vel, orci. Vestibulum justo. Suspendisse non nisl. Pellentesque pretium adipiscing elit. Phasellus fermentum consequat augue. Sed pede nisl, fermentum vel, vulputate id, sollicitudin sed, ligula. Cras suscipit, quam et euismod sagittis, nisl felis gravida felis, quis pulvinar purus est vel pede. Suspendisse mattis est ac nunc. Curabitur rutrum, turpis sit amet commodo tempus, metus lorem commodo lectus, eget fringilla justo nisi et purus. Ut quam sapien, vehicula quis, rhoncus non, sagittis nec, risus.

Donec eget augue ac lacus adipiscing porta. Maecenas pede. Vivamus molestie. Duis condimentum ligula auctor pede. Nullam ullamcorper rhoncus erat. Ut ornare interdum urna. Suspendisse potenti. Curabitur mattis mauris nec risus. Aenean iaculis turpis eu tortor. Donec nec ante non mauris pellentesque fringilla.

Phasellus vitae dui id orci sodales cursus. Curabitur sed nulla quis mauris tincidunt iaculis. Vivamus semper semper orci. Phasellus suscipit ante vitae leo. Sed arcu ipsum, condimentum id, luctus in, sodales eu, magna. In dictum, arcu quis pharetra vestibulum, ante enim placerat lacus, vitae placerat est leo vitae elit. Pellentesque bibendum enim vulputate eros. Nunc

laoreet. Pellentesque habitant morbi tristique senectus et netus et malesuada fames ac turpis egestas. Praesent purus odio, euismod sit amet, aliquam a, volutpat in, augue. Phasellus id massa. Suspendisse suscipit ligula pharetra dolor. Pellentesque vel pede.

Aliquam pharetra est sit amet magna. Aliquam varius. Donec eu lectus et nisl iaculis porttitor. Morbi mattis, mauris sed luctus hendrerit, nulla velit molestie dolor, ac volutpat urna augue vel quam. Maecenas pellentesque libero et massa. Integer vestibulum, lacus at mattis euismod, nisl arcu commodo lectus, ut euismod dolor ligula sit amet libero. Nam in ligula sit amet ante eleifend aliquet. Phasellus feugiat erat at nulla. Proin in lectus. Proin laoreet leo laoreet leo congue lacinia. Quisque non diam sit amet enim ultrices commodo. Praesent fermentum lectus sed ligula. Integer pulvinar accumsan pede. Quisque molestie ligula eget odio. Vestibulum ante ipsum primis in faucibus orci luctus et ultrices posuere cubilia Curae;

## APPENDIX C DERIVATION OF THE $\Upsilon$ FUNCTION

**Proposition C.1.** *The Upsilon Function*

(1) If  $\beta > 0$  and  $\alpha \neq 0$ , then for all  $n \geq -1$ ,

$$\begin{aligned} I_n(c; \alpha; \beta; \delta) &= -\frac{e^{\alpha c}}{\alpha} \sum_{i=0}^n \left(\frac{\beta}{\alpha}\right)^{n-i} \text{Hh}_i(\beta c - \delta) \\ &\quad + \left(\frac{\beta}{\alpha}\right)^{n+1} \frac{\sqrt{2\pi}}{\beta} e^{\frac{\alpha\delta}{\beta} + \frac{\alpha^2}{2\beta^2}} \phi\left(-\beta c + \delta + \frac{\alpha}{\beta}\right) \end{aligned}$$

(2) If  $\beta < 0$  and  $\alpha < 0$ , then for all  $x \geq -1$

$$\begin{aligned} I_n(c; \alpha; \beta; \delta) &= -\frac{e^{\alpha c}}{\alpha} \sum_{i=0}^n \left(\frac{\beta}{\alpha}\right)^{n-i} \text{Hh}_i(\beta c - \delta) \\ &\quad - \left(\frac{\beta}{\alpha}\right)^{n+1} \frac{\sqrt{2\pi}}{\beta} e^{\frac{\alpha\delta}{\beta} + \frac{\alpha^2}{2\beta^2}} \phi\left(\beta c - \delta - \frac{\alpha}{\beta}\right) \end{aligned}$$

*Proof.* Case 1.

$\beta > 0$  and  $\alpha \neq 0$ . Since, for any constant  $\alpha$  and  $n \geq 0$ ,  $e^{\alpha x} \text{Hh}_n(\beta x - \delta) \rightarrow 0$  as  $x \rightarrow \infty$  thanks to (B4), integration by parts leads to

$$I_n = -\frac{1}{\alpha} \text{Hh}(\beta c - \delta) e^{\alpha c} + \frac{\beta}{\alpha} \int_c^\infty e^{\alpha x} \text{Hh}_{n-1}(\beta x - \delta) dx$$

In other words, we have a recursion, for  $n \geq 0$ ,  $I_n = -(e^{\alpha c} \alpha) \text{Hh}_n(\beta c - \delta) + \left(\frac{\beta}{\alpha}\right) I_{n-1}$  with

$$\begin{aligned} I_{-1} &= \sqrt{2\pi} \int_c^\infty e^{\alpha x} \phi(-\beta x + \delta) dx \\ &= \frac{\sqrt{2\pi}}{\beta} e^{\frac{\alpha\delta}{\beta} + \frac{\alpha^2}{2\beta^2}} \phi\left(-\beta c + \delta + \frac{\alpha}{\beta}\right) \end{aligned}$$

Solving it yields, for  $n \geq -1$ ,

$$\begin{aligned}
I_n &= -\frac{e^{\alpha c}}{\alpha} \sum_{i=0}^n \left(\frac{\beta}{\alpha}\right)^i \text{Hh}_{n-i}(\beta c + \delta) + \left(\frac{\beta}{\alpha}\right)^{n+1} I_{-1} \\
&= -\frac{e^{\alpha c}}{\alpha} \sum_{i=0}^n \left(\frac{\beta}{\alpha}\right)^{n-i} \text{Hh}_i(\beta c + \delta) \\
&\quad + \left(\frac{\beta}{\alpha}\right)^{n+1} \frac{\sqrt{2\pi}}{\beta} e^{\frac{\alpha\delta}{\beta} + \frac{\alpha^2}{2\beta^2}} \phi\left(-\beta c + \delta + \frac{\alpha}{\beta}\right)
\end{aligned}$$

where the sum over an empty set is defined to be zero.  $\square$

*Case2.*  $\beta < 0$  and  $\alpha < 0$ . In this case, we must also have, for  $n \geq 0$  and any constant  $\alpha < 0$ ,  $e^{\alpha x} \text{Hh}_n(\beta x - \delta) \rightarrow 0$  as

$x \rightarrow \infty$ , thanks to (B5). Using integration by parts, we again have the same recursion, for  $n \geq 0$ ,  $I_n = -(e^{\alpha c}/\alpha) \text{Hh}_n(\beta c - \delta) + (\beta/\alpha) I_{n-1}$ , but with a different initial condition

$$\begin{aligned}
I_{-1} &= \sqrt{2\pi} \int_c^\infty e^{\alpha x} \varphi(-\beta x + \delta) dx \\
&= -\frac{\sqrt{2\pi}}{\beta} \exp\left\{\frac{\alpha\delta}{\beta} + \frac{\alpha^2}{2\beta^2}\right\} \phi\left(\beta c - \delta - \frac{\alpha}{\beta}\right)
\end{aligned}$$

Solving it yields (B8), for  $n \geq -1$ .

Finally, we sum the double exponential and the normal random variables

*Proposition B.3.*

Suppose  $\{\xi_1, \xi_2, \dots\}$  is a sequence of i.i.d. exponential random variables with rate  $\eta > 0$ , and  $Z$  is a normal variable with distribution  $N(0, \sigma^2)$ . Then for every  $n \geq 1$ , we have: (1) The density functions are given by:

$$f_{Z+\sum_{i=1}^n \xi_i}(t) = (\sigma\eta)^n \frac{e^{(\sigma\eta)^2/2}}{\sigma\sqrt{2\pi}} e^{-t\eta} \text{Hh}_{n-1}\left(-\frac{t}{\sigma} + \sigma\eta\right)$$

$$f_{Z-\sum_{i=1}^n \xi_i}(t) = (\sigma\eta)^n \frac{e^{(\sigma\eta)^2/2}}{\sigma\sqrt{2\pi}} e^{-t\eta} \text{Hh}_{n-1}\left(\frac{t}{\sigma} + \sigma\eta\right)$$

(2) The tail probabilities are given by

$$P(Z + \sum_{i=1}^n \xi_i \geq x) = (\sigma\eta)^n \frac{e^{(\sigma\eta)^2/2}}{\sigma\sqrt{2\pi}} e^{-t\eta} I_{n-1}(x; -\eta, -\frac{1}{\sigma}, -\sigma\eta)$$

$$P(Z - \sum_{i=1}^n \xi_i \geq x) = (\sigma\eta)^n \frac{e^{(\sigma\eta)^2/2}}{\sigma\sqrt{2\pi}} e^{-t\eta} I_{n-1}(x; \eta, \frac{1}{\sigma}, -\sigma\eta)$$

*Proof. Case 1. The densities of  $Z + \sum_{i=1}^n \xi_i$ , and  $Z - \sum_{i=1}^n \xi_i$ . We have*

$$\begin{aligned} f_{Z+\sum_{i=1}^n \xi_i}(t) &= \int_{-\infty}^{\infty} f_{\sum_{i=1}^n \xi_i}(t-x) f_Z(x) dx \\ &= e^{-t\eta} (\eta^n) \int_{-\infty}^{\infty} t \frac{e^{x\eta} (t-x)^{n-1}}{(n-1)!} \frac{1}{\sigma\sqrt{2\pi}} e^{-x^2/(2\sigma^2)} dx \\ &= e^{-t\eta} (\eta^n) e^{(\sigma\eta)^2/(2)} \int_{-\infty}^{\infty} t \frac{(t-x)^{n-1}}{(n-1)!} \frac{1}{\sigma\sqrt{2\pi}} e^{-(x-\sigma^2\eta)^2/(2\sigma^2)} dx \end{aligned}$$

*Letting  $y = (x - \sigma^2\eta)/\sigma$  yields*

$$\begin{aligned} f_{Z+\sum_{i=1}^n \xi_i}(t) &= e^{-t\eta} (\eta^n) e^{(\sigma\eta)^2/(2)} \sigma^{n-1} \\ &\times \int_{-\infty}^{t/\sigma - \sigma\eta} \frac{(t/\sigma - y - \sigma\eta)^{n-1}}{(n-1)!} \frac{1}{\sqrt{2\pi}} e^{-y^2/2} dy \\ &= \frac{e^{(\sigma\eta)^2/2}}{\sqrt{2\pi}} (\sigma^{n-1} \eta^n) e^{-t\eta} Hh_{n-1}(-t/\sigma + \sigma\eta) \end{aligned}$$

*because  $(1/(n-1)!) \int_{-\infty}^a (a-y)^{n-1} e^{-y^2/2} dy = Hh_{n-1}(a)$ . The derivation of  $f_{Z+\sum_{i=1}^n \xi_i}(t)$  is similar.*

*Case 2.  $P(Z + \sum_{i=1}^n \xi_i \geq x)$  and  $P(Z - \sum_{i=1}^n \xi_i \geq x)$ . From (B9), it is clear that*

$$P(Z + \sum_{i=1}^n \xi_i \geq x) = \frac{(\sigma\eta)^n e^{(\sigma\eta)^2/2}}{\sigma\sqrt{2\pi}} \int_x^{\infty} e^{(-i\eta)} Hh_{n-1}\left(-\frac{t}{\sigma} + \sigma\eta\right) dt$$



$$= \frac{(\sigma\eta)^n e^{(\sigma\eta)^2/2}}{\sigma\sqrt{2\pi}} I_{n-1}(x; -\eta, -\frac{1}{\sigma}, -\sigma\eta) dt$$

by (B6). We can compute  $P(Z - \sum_{i=1}^n \xi_i \geq x)$  similarly.

**Theorem C.1.** *Theorem With  $\pi_n := P(N(t) = n) = e^{-\lambda T}(\lambda T)^n/n!$  and  $I_n$  in Proposition ?? , we have*

$$\begin{aligned} P(Z(T) \geq a) &= \frac{e^{(\sigma\eta_1)^2 T/2}}{\sigma\sqrt{2\pi T}} \sum_{n=1}^{\infty} \pi_n \sum_{k=1}^n P_{n,k}(\sigma\sqrt{T}\eta_1)^k \times I_{k-1}(a - \mu T; -\eta_1, -\frac{1}{\sigma\sqrt{T}}, -\sigma\eta_1\sqrt{T}) \\ &\quad + \frac{e^{(\sigma\eta_2)^2 T/2}}{\sigma\sqrt{2\pi T}} \sum_{n=1}^{\infty} \pi_n \sum_{k=1}^n Q_{n,k}(\sigma\sqrt{T}\eta_2)^k \\ &\quad \times I_{k-1}(a - \mu T; \eta_2, \frac{1}{\sigma\sqrt{T}}, -\sigma\eta_2\sqrt{T}) \\ &\quad + \pi_0 \phi\left(-\frac{a - \mu T}{\sigma\sqrt{T}}\right) \end{aligned}$$

*Proof by the decomposition (B2)*

$$\begin{aligned} P(Z(T) \geq a) &= \sum_{n=0}^{\infty} \pi_n P(\mu T + \sigma\sqrt{T}Z + \sum_{j=1}^n Y_j \geq a) \\ &= \pi_0 P(\mu T + \sigma\sqrt{T}Z \geq a) \\ &\quad + \sum_{n=1}^{\infty} \pi_n \sum_{k=1}^n P_{n,k} P(\mu T + \sigma\sqrt{T}Z + \sum_{j=1}^n \xi_j^+ \geq a) \\ &\quad + \sum_{n=1}^{\infty} \pi_n \sum_{k=1}^n Q_{n,k} P(\mu T + \sigma\sqrt{T}Z - \sum_{j=1}^n \xi_j^- \geq a) \end{aligned}$$

*The result now follows via (B11) and (B12) for  $\eta_1 > 1$  and  $\eta_2 > 0$ .*

## APPENDIX D DERIVATION OF THE $\Upsilon$ FUNCTION

We first decompose the sum of the double exponential random variables.

The memoryless property of exponential random variables yields  $(\xi^+ - \xi^- | \xi^+ > \xi^-) =^d \xi^+$  and  $(\xi^+ - \xi^- | \xi^+ < \xi^-) =^d -\xi^-$ , thus leading to the conclusion that

$$\xi^+ - \xi^- = \begin{cases} \xi^+ & \text{with probability } \eta_2/(\eta_1 + \eta_2) \\ -\xi^- & \text{with probability } \eta_1/(\eta_1 + \eta_2) \end{cases}.$$

because the probabilities of the events  $\xi^+ > \xi^-$  and  $\xi^+ < \xi^-$  are  $\eta_2/(\eta_1 + \eta_2)$  and  $\eta_1/(\eta_1 + \eta_2)$ , respectively. The following proposition extends (B.1.)

**Proposition B.1.** For every  $n \geq 1$ , we have the following decomposition

$$\sum_{i=1}^n Y_i =^d \begin{cases} \sum_{i=1}^k \xi_i^+ & \text{with probability } P_{n,k}, k = 1, 2, \dots, n \\ -\sum_{i=1}^k \xi_i^- & \text{with probability } Q_{n,k}, k = 1, 2, \dots, n \end{cases}.$$

where  $P_{n,k}$  and  $Q_{n,k}$  are given by

$$P_{n,k} = \sum_{i=k}^{n-1} \binom{n-k-1}{i-k} \binom{n}{i} \left(\frac{\eta_1}{\eta_1 + \eta_2}\right)^{i-k} \left(\frac{\eta_2}{\eta_1 + \eta_2}\right)^{n-i} p^i q^{n-i}$$

$$1 \leq k \leq n-1$$

$$Q_{n,k} = \sum_{i=k}^{n-1} \binom{n-k-1}{i-k} \binom{n}{i} \left(\frac{\eta_1}{\eta_1 + \eta_2}\right)^{n-i} \left(\frac{\eta_2}{\eta_1 + \eta_2}\right)^{i-k} p^{n-i} q^i$$

$$1 \leq k \leq n-1, P_{n,n} = p^n, Q_{n,n} = q^n$$

and  $\binom{0}{0}$  is defined to be one. Hence  $\xi_i^+$  and  $\xi_i^-$  are i.i.d. exponential random variables with rates  $\eta_1$  and  $\eta_2$ , respectively.

As a key step in deriving closed-form solutions for call and put options, this proposition indicates that the sum of the i.i.d. double exponential random variable can be written, in

distribution, as a randomly mixed gamma random variable. To prove Proposition B.1, the following lemma is needed.

Lemma B.1.

$$\sum_{i=1}^n \xi_i^+ - \sum_{i=1}^n \xi_i^-$$

$$=^d \left\{ \begin{array}{ll} \sum_{i=1}^k \xi_i & \text{with probability } \binom{n-k+m-1}{m-1} \left(\frac{\eta_1}{\eta_1+\eta_2}\right)^{n-k} \left(\frac{\eta_2}{\eta_1+\eta_2}\right)^m, k = 1, \dots, n \\ -\sum_{i=1}^l \xi_i & \text{with probability } \binom{n-l+m-1}{n-1} \left(\frac{\eta_1}{\eta_1+\eta_2}\right)^n \left(\frac{\eta_2}{\eta_1+\eta_2}\right)^{m-1}, l = 1, \dots, m \end{array} \right\}.$$

We prove it by introducing the random variables  $A(n, m) = \sum_{i=1}^n \xi_i - \sum_{j=1}^m \tilde{\xi}_j$ . Then

$$A(n, m) =^d \left\{ \begin{array}{ll} A(n-1, m-1) + \xi^+ & \text{with probability } \eta_2/(\eta_1 + \eta_2) \\ A(n-1, m-1) - \xi^- & \text{with probability } \eta_1/(\eta_1 + \eta_2) \end{array} \right\}.$$

$$=^d \left\{ \begin{array}{ll} A(n, m-1) & \text{with probability } \eta_2/(\eta_1 + \eta_2) \\ A(n-1, m) & \text{with probability } \eta_1/(\eta_1 + \eta_2) \end{array} \right\}.$$

via B.1.. Now suppose horizontal axis that are representing the number of  $\{\zeta_i^+\}$  and vertical axis representing the number of  $\{\zeta_i^-\}$ . Suppose we have a random walk on the integer lattice points. Starting from any point  $(n, m)$ ,  $n, m \geq 1$ , the random walk goes either one step to the left with probability  $\eta_1/(\eta_1 + \eta_2)$  or one step down with probability  $\eta_2/(\eta_1 + \eta_2)$ , and the random walks stops once it reaches the horizontal or vertical axis. For any path from  $(n, m)$  to  $(k, 0)$ ,  $1 \geq k \geq n$ , it must reach  $(k, 1)$  first before it makes a final move to  $(k, 0)$ . Furthermore, all the paths going from  $(n, m)$  to  $(k, 1)$  must have exactly  $n-k$  lefts and  $m-1$  downs, whence the total number of such paths is  $\binom{n-k+m-1}{m-1}$ . Similarly the total number of paths from  $(n, m)$  to  $(0, l)$ ,  $1 \geq l \geq m$ , is  $\binom{n-l+m-1}{n-1}$ . Thus

$$A(n, m) =^d \left\{ \begin{array}{l} \sum_{i=1}^k \xi_i \quad \text{with probability } \binom{n-k+m-1}{m-1} \left(\frac{\eta_1}{\eta_1+\eta_2}\right)^{n-k} \left(\frac{\eta_2}{\eta_1+\eta_2}\right)^m, k = 1, \dots, n \\ - \sum_{i=1}^l \xi_i \quad \text{with probability } \binom{n-l+m-1}{n-1} \left(\frac{\eta_1}{\eta_1+\eta_2}\right)^n \left(\frac{\eta_2}{\eta_1+\eta_2}\right)^{m-1}, l = 1, \dots, m \end{array} \right\}.$$

and the lemma is proven.

Now, let's prove the proposition B.1. By the same analogy used in Lemma B.1 to compute probability  $P_{n,m}, 1 \geq k \geq n$ , the probability weight assigned to  $\sum_{i=1}^k \xi_i^+$  when we decompose  $\sum_{i=1}^k Y_i$ , it is equivalent to consider the probability of the random walk ever reach  $(k,0)$  starting from the point  $(i,n-i)$  being  $\binom{n}{i} p^i q^{n-i}$ . Note that the point  $(k,0)$  can only be reached from point  $(i,n-i)$  such that  $k \geq i \geq n-1$ , because the random walk can only go left or down, and stops once it reaches the horizontal axis. Therefore, for  $1 \geq k \geq n-1$ , (B3) leads to

$$\begin{aligned} P_{n,k} &= \sum_{i=k}^{n-1} n-1 P(\text{going from } (i, n-i) \text{ to } (k, 0)). P(\text{starting from } (i, n-i)) \\ &= \sum_{i=k}^{n-1} \binom{i + (n-i) - k - 1}{(n-i) - 1} \binom{n}{i} \left(\frac{\eta_1}{\eta_1 + \eta_2}\right)^{i-k} \left(\frac{\eta_2}{\eta_1 + \eta_2}\right)^{n-i} p^i q^{n-i} \\ &= \sum_{i=k}^{n-1} \binom{n-k-1}{n-i-1} \binom{n}{i} \left(\frac{\eta_1}{\eta_1 + \eta_2}\right)^{i-k} \left(\frac{\eta_2}{\eta_1 + \eta_2}\right)^{n-i} p^i q^{n-i} \\ &= \sum_{i=k}^{n-1} \binom{n-k-1}{i-k} \binom{n}{i} \left(\frac{\eta_1}{\eta_1 + \eta_2}\right)^{i-k} \left(\frac{\eta_2}{\eta_1 + \eta_2}\right)^{n-i} p^i q^{n-i} \end{aligned}$$

Of course  $P_{n,n} = p^n$ . Similarly, we can compute  $Q_{n,k}$ :

$$\begin{aligned} Q_{n,k} &= \sum_{i=k}^{n-1} n-1 P(\text{going from } (n-i, i) \text{ to } (0, k)). P(\text{starting from } (n-i, i)) \\ &= \sum_{i=k}^{n-1} \binom{i + (n-i) - k - 1}{(n-i) - 1} \binom{n}{n-i} \left(\frac{\eta_1}{\eta_1 + \eta_2}\right)^{n-i} \left(\frac{\eta_2}{\eta_1 + \eta_2}\right)^{i-k} p^{n-i} q^i \end{aligned}$$

$$= \sum_{i=k}^{n-1} \binom{n-k-1}{i-k} \binom{n}{i} \left(\frac{\eta_1}{\eta_1 + \eta_2}\right)^{n-i} \left(\frac{\eta_2}{\eta_1 + \eta_2}\right)^{i-k} p^{n-i} q^i$$

with  $Q_{n,n} = q^n$ . Incidentally, we have also got  $\sum k = 1n(P_{n,k} + Q_{n,k}) = 1$

B.2. Let's develop now the results on Hh functions. First of all, note that  $Hh_n(x) \rightarrow 0$ , as  $x \rightarrow \infty$ , for  $n \geq -1$ ; and  $Hh_n(x) \rightarrow \infty$ , as  $x \rightarrow -\infty$ , for  $n \geq -1$ ; and  $Hh_0(x) = \sqrt{2\pi}\phi(-x) \rightarrow \sqrt{2\pi}$ , as  $x \rightarrow -\infty$ . Also, for every  $n \geq -1$ , as  $x \rightarrow \infty$ ,

$$\lim Hh_n(x) / \left\{ \frac{1}{x^{n+1}} e^{-\frac{x^2}{2}} \right\} = 1$$

and as  $x \rightarrow \infty$

$$Hh_n(x) = O(|x|^n)$$

Here (B4) is clearly true for  $n = -1$ , while for  $n \geq 0$  note that as  $x \rightarrow \infty$ ,

$$\begin{aligned} Hh_n(x) &= \frac{1}{n!} \int_x^\infty (t-x)^n e^{-\frac{t^2}{2}} dt \\ &\leq \frac{2^n}{n!} \int_{-\infty}^\infty |t|^n e^{-t^2} 2dt + \frac{2^n}{n!} \int_{-\infty}^\infty |x|^n e^{-t^2} 2dt = O(|x|^n) \end{aligned}$$

For option pricing it is important to evaluate the integral  $I_n(c; \alpha; \beta; \delta)$ ,

$$I_n(c; \alpha; \beta; \delta) = \int_c^\infty e^{\alpha x} Hh_n(\beta x - \delta) dx, n \geq 0$$

for arbitrary constants  $\alpha, c$  and  $\beta$ .

## REFERENCES

- Garfinkle, David, Horowitz, Gary T, and Strominger, Andrew. "Charged black holes in string theory." *Physical Review D* 43 (1991).10: 3140.
- Green, Karen L. "A wrinkle in time." *comiXology* (2008).
- L'engle, Madeleine. *A Wrinkle in Time: 50th Anniversary Commemorative Edition*, vol. 1. Macmillan, 2012.
- Strickler, Howard D, Rosenberg, Philip S, Devesa, Susan S, Hertel, Joan, Fraumeni Jr, Joseph F, and Goedert, James J. "Contamination of poliovirus vaccines with simian virus 40 (1955-1963) and subsequent cancer rates." *Jama* 279 (1998).4: 292–295.

## BIOGRAPHICAL SKETCH

This section is where your biographical sketch is typed in the [bio.tex](#) file. It should be in third person, past tense. Do not put personal details such as your birthday in the file. Again, to make a full paragraph you must write at least three sentences.