

# Documentation

*A. Casares M.*


*January 5th, 2017*

**Problem:** Given the distances from an interior point to three vertices of a rectangle, which are the maximum and minimum possible areas?

## Description and analysis:


Originally, this problem asked which was the distance to the fourth vertex. Let's answer this question first. See next a plot of the problem, taken the original data:  $AX = 3$ ,  $BX = 12$ ,  $CX = 11$ .  $X$  is an interior point, such as its distances agree with the data.  $DX$  is the unknown distance.

**Fig.1**



Since we don't know the dimensions of the rectangle, nor any angle, it may seem that some data is missing. But with the help of Pithagoras and placing a rectangular coordinate system in A; calling  $(x,y)$  the coordinates of  $X$  in that system, and tracing parallel lines to the sides of the rectangle through  $X$ , we have:  
 $\overline{XD} = \sqrt{\overline{Xp}^2 + \overline{Xo}^2}$ .

**Fig.2**



Considering that  $\overline{Xp}^2 = \overline{CX}^2 - x^2$ ,  $\overline{Xo}^2 = \overline{BX}^2 - y^2$  and  $x^2 + y^2 = r^2$ , the replacement in the previous equality gives  $\overline{XD} = \sqrt{\overline{CX}^2 + \overline{BX}^2 - (x^2 + y^2)} = \sqrt{\overline{CX}^2 + \overline{BX}^2 - \overline{AX}^2}$

Replacing our data, we get  $\overline{XD} = \sqrt{11^2 + 12^2 - 3^2} = 16$

And this value doesn't depend on the height or base of the rectangle, nor on the position of point X. So, given only the 3 known distances, the fourth is a constant.

However, depending of the position of X (which can be expressed also by its polar coordinates  $(r, \alpha)$ ), the rectangle's dimensions must change to conform with the data and the fourth distance. Let's find which are the base and length of the rectangle for a given  $\alpha$ ,  $0 \leq \alpha \leq 90$ .

From Fig.2 is not difficult to deduce that the base  $b = \overline{Xo} + x = \sqrt{\overline{BX}^2 - y^2} + x$  and the height  $h = \overline{Xp} + y = \sqrt{\overline{CX}^2 - x^2} + y$ .

Since  $x$  and  $y$  can be expressed as functions of  $\alpha$ :  $x = r \times \cos(\alpha)$ ,  $y = r \times \sin(\alpha)$ , we can finally get  $b$  and  $h$  as functions of  $\alpha$  and the data.

And so is the area  $A$ , which is taking us to the core of the problem:

$$A = b \times h = (\sqrt{\overline{BX}^2 - y^2} + x) \times (\sqrt{\overline{CX}^2 - x^2} + y) \Leftrightarrow$$

$$A = (\sqrt{\overline{BX}^2 - (r \times \sin(\alpha))^2} + r \times \cos(\alpha)) \times (\sqrt{\overline{CX}^2 - (r \times \cos(\alpha))^2} + r \times \sin(\alpha))$$

And as a function only of  $\sin(\alpha)$ :

$$A = (\sqrt{\overline{BX}^2 - (r \times \sin(\alpha))^2} + r \times \sqrt{1 - \sin^2(\alpha)}) \times (\sqrt{\overline{CX}^2 - (r \times \sqrt{1 - \sin^2(\alpha)})^2} + r \times \sin(\alpha))$$

Even replacing the data of the particular given problem, we have a long formula:

$$A = (\sqrt{144 - (3 \times \sin(\alpha))^2} + 3 \times \sqrt{1 - \sin^2(\alpha)}) \times (\sqrt{121 - (3 \times \sqrt{1 - \sin^2(\alpha)})^2} + 3 \times \sin(\alpha))$$

We could simplify a little using the new variable  $s = \sin^2(\alpha)$ :

$$A = (\sqrt{BX^2 - r^2 \times s} + r \times \sqrt{1 - s}) \times (\sqrt{CX^2 - (r \times \sqrt{1 - s})^2} + r \times \sqrt{s})$$

So, we have arrived to an expression giving the rectangle's area as a function of  $\alpha$ , sort of indirectly, via the new  $s$  variable. Now, to find analytically the maximum or minimum of this function, we'd have to find its derivative with respect to  $\alpha$ , equate it to zero, and solve the corresponding equation.

Something like:  $\frac{dA}{d\alpha} = \frac{dA}{ds} \times \frac{ds}{d\alpha} = 0$ .

Now,  $\frac{ds}{d\alpha} = 2 \times \sin(\alpha) \times \cos(\alpha) = \sin(2\alpha)$ , and this is zero at  $\alpha = 0$  and  $\alpha = 90$  degrees. In those cases, the point X is on the base or the height of the rectangle. We shall see later if these are maxima or minima.

But the hard, substantial, part of the derivative is  $\frac{dA}{ds}$ . To equate it to zero gives an embarrassingly hard equation, to be solved only by some numerical method, that I spare the reader. Instead of this half-closed procedure, we may recur from the beginning to a numerical solution.

We can take two routes to find the  $\alpha$  giving the maximum or minimum areas, and the value of these extremes. The first, graphical and empirical; the second, thoroughly numerical. Both must give us compatible answers.

### Graphical procedure:

- In the computer, take a sample of angle's values covering the whole  $[0,90]$  interval, with an adequate increment (say 0.1 degrees), and find the area with the formula or the product  $b \times h$  in each point. Using these results, plot the resulting areas to have an image of the behaviour of the curve, and find the maximum and minimum from the data. We can try to do it by hand, but the computer comparison of values will give us the answer, with a precision of one decimal digit for the angle, and several more for the area.

### Numerical procedure:

- a).- Using the area values collected, find the numerical derivative of  $A$  with respect to  $\alpha$ :  
There are several formulas to do this computation, giving us approximated values of the derivative in the selected points. For instance, we can use the so called "central differences" formula:  
 $\frac{dA}{d\alpha} = \frac{A_{i+1} - A_{i-1}}{2h}$ , where  $h$  is the difference  $\alpha_{i+1} - \alpha_i$ .

This is a second degree's formula, meaning that its error is roughly proportional to  $h^2$ , which is good for small  $h$ . The  $h$  value should be rather small, to avoid errors of mathematical modeling; but not too much, because in that case the cancellation errors, typical of floating point representation of real numbers in the computer, are presented. The selected value  $h = 0.1$  may be a good one for this case.

- b).- Plot the found values of the derivative, so we can appreciate its linearity.
- c).- Apply an adequate regression model to the dataset of obtained values. In our case, the plot showed a high linearity, so a straightforward linear model is the most obvious option. We'll get a formula  $\frac{dA}{d\alpha} = \beta_1 \alpha + \beta_0$ . That is the regression line.
- d).- If the squared correlation coefficient  $R^2$  is near 1, showing that the adjustment is good, solve the linear equation  $\beta_1 \alpha + \beta_0 = 0$ , giving us  $\alpha = -\frac{\beta_0}{\beta_1}$ . This is the angle corresponding to the zero derivative, and, in accordance to Calculus, is the point where the function  $A(\alpha)$  takes a maximum or minimum value. In order to know this, we must look at the sign of the second derivative - the slope of the regression line (see how much work spares us this equation) - in this case simply  $\beta_1$ . If this is negative, as happens in this case, the extreme is a maximum.
- e).- If all goes right, the found root value and respective extreme must strongly agree with the empirical values found in the graphical procedure.

## Implementation:

The data product shows an implementation of these little theory, making use of R and Shiny.

You have the chance of work with several cases, one of which you must select in the beginning. Once selected, you can play with the values of  $\alpha$  angle (called BAX), through a provided slider widget, watching how the variation of  $\alpha$  changes the dimensions  $b$  and  $h$  of the rectangle, in order to adjust to the conditions of the problem. These variations also affect the second and third plots.

The second plot corresponds to the graphical procedure, and there you can see the maximum and minimum areas found in that procedure, altogether with the value of the angle and the rectangle's dimensions. Using the slider widget you can try to get the same answer printed on the plot.

In the third plot you find the derivative's values, making a remarkably linear path (in fact, only because of the approximation inaccuracies the path is not exactly linear), and the regression line altogether, showing their close vicinity. The difference between the root of the regression equation and the intersection of the derivative data points with the x axis is practically indistinguishable. These facts are confirmed numerically by the high value of the adjusted R-squared coefficient, and the concordance to several decimal digits between the greatest areas found in the empirical and numerical proceeding.

The minimal area, as was suggested before, happens to be on the  $\alpha = 0$  extreme. It is a global minimum.

Finally, it should be noticed the easiness provided by the R's linear modeling function (lm) to the c) and d) points in the numerical method: everything gets done there with only two propositions.