

PARTIAL PROOF TERMS IN THE STUDY OF PROOF SEARCH

José Espírito Santo and Ana Catarina Sousa
CMAT - University of Minho

Women in Logic Workshop
University of Tallinn, Estonia

9th July 2024

Finished Proofs

$$\frac{\frac{\frac{}{A, B \vdash A} Ax}{A \vdash B \supset A} I \supset}{\vdash A \supset B \supset A} I \supset$$



C.H. Isomorphism

Proof Terms

$$\lambda x^A. \lambda y^B. x$$

Finished Proofs

$$\frac{\frac{\frac{}{A, B \vdash A} \text{Ax}}{A \vdash B \supset A} I \supset}{\vdash A \supset B \supset A} I \supset$$

$\longrightarrow$

C.H. Isomorphism

Proof Terms

$$\lambda x^A. \lambda y^B. x$$

Unfinished Proofs

$$\frac{\frac{?}{A \vdash B \supset A}}{\vdash A \supset B \supset A} I \supset$$

$\longrightarrow$

?

Finished Proofs

$$\frac{\frac{\frac{}{A, B \vdash A} Ax}{A \vdash B \supset A} I \supset}{\vdash A \supset B \supset A} I \supset$$



C.H. Isomorphism

Proof Terms

$$\lambda x^A. \lambda y^B. x$$

Unfinished Proofs

$$\frac{\frac{?}{A \vdash B \supset A}}{\vdash A \supset B \supset A} I \supset$$



In the literature...

Finished Proofs

$$\frac{\frac{\frac{}{A, B \vdash A} Ax}{A \vdash B \supset A} I \supset}{\vdash A \supset B \supset A} I \supset$$



C.H. Isomorphism

Proof Terms

$$\lambda x^A. \lambda y^B. x$$

Unfinished Proofs

$$\frac{?}{\frac{A \vdash B \supset A}{\vdash A \supset B \supset A} I \supset}$$



In the literature...

Meta-variables

e.g. [Muñoz, 2001]

- Model, with the help of explicit substitutions, the reuse of proofs/ delay of choices in unification and proof search;
- Used in the meta-theory of proof assistants based on type theory.

Finished Proofs

$$\frac{\frac{\frac{}{A, B \vdash A} Ax}{A \vdash B \supset A} I \supset}{\vdash A \supset B \supset A} I \supset$$



C.H. Isomorphism

Proof Terms

$$\lambda x^A. \lambda y^B. x$$

Unfinished Proofs

$$\frac{?}{\frac{A \vdash B \supset A}{\vdash A \supset B \supset A} I \supset}$$



In the literature...

Open Proofs and Open Proof Terms

[Geuvers, Jojgov, 2002]

- Developed directly for HOL; deal with proofs in ND, possibly containing a generic "claim" inference;
- Does not have an explicit formalization of a search process.

Finished Proofs

$$\frac{\frac{\frac{}{A, B \vdash A} Ax}{A \vdash B \supset A} I \supset}{\vdash A \supset B \supset A} I \supset$$



C.H. Isomorphism

Proof Terms

$$\lambda x^A. \lambda y^B. x$$

Unfinished Proofs

$$\frac{?}{\frac{A \vdash B \supset A}{\vdash A \supset B \supset A} I \supset}$$



In the literature...

Coinductive Proof Search

e.g. [JES, Matthes, Pinto, 2021]

- Employs proof terms with meta-variables X^σ , allowing the representation of an entire search space;
- Does not yet provide means for expressing proof search procedures.

Finished Proofs

$$\frac{\frac{\frac{}{A, B \vdash A} Ax}{A \vdash B \supset A} I \supset}{\vdash A \supset B \supset A} I \supset$$



C.H. Isomorphism

Proof Terms

$$\lambda x^A. \lambda y^B. x$$

Unfinished Proofs

$$\frac{?}{\frac{A \vdash B \supset A}{\vdash A \supset B \supset A} I \supset}$$



In the literature...

L-Framework

[Olarte, Pimentel, Rocha, 2023]

- Term rewriting as a meta-logical framework for the specification of sequent calculi and the automation of meta-theory;
- More focused in automation and bottom-up proof search.

Finished Proofs

$$\frac{\frac{\frac{}{A, B \vdash A} Ax}{A \vdash B \supset A} I \supset}{\vdash A \supset B \supset A} I \supset$$



C.H. Isomorphism

Proof Terms

$$\lambda x^A. \lambda y^B. x$$

Unfinished Proofs

$$\frac{\frac{?}{A \vdash B \supset A}}{\vdash A \supset B \supset A} I \supset$$



Our approach...

Finished Proofs

$$\frac{\frac{\frac{}{A, B \vdash A} Ax}{A \vdash B \supset A} I \supset}{\vdash A \supset B \supset A} I \supset$$



C.H. Isomorphism

Proof Terms

$$\lambda x^A. \lambda y^B. x$$

Unfinished Proofs

$$\frac{?}{A \vdash B \supset A} I \supset$$



Our approach...

Extension of the
C.H. Isomorphism

Partial Proof Terms

$$\lambda x^A. (A \vdash B \supset A)$$

- *Formal sequents*, i.e., sequents occurring as components of proof terms;
- *Idealized proof search* - theoretical study of proof search without considerations of control, effectiveness or implementation.

A first case study: LJ1

Formulas

$$A, B ::= p \mid A \supset B$$

A first case study: LJT

Formulas

$$A, B ::= p \mid A \supset B$$

Sequents

$$\Gamma \vdash A \qquad \Gamma; A \vdash p$$

A first case study: LJT

Formulas

$$A, B ::= p \mid A \supset B$$

Sequents

$$\Gamma \vdash A \qquad \Gamma; A \vdash p$$

Inference Rules

$$\frac{}{\Gamma; p \vdash p} \text{ AX}$$

$$\frac{\Gamma, x : A; A \vdash p}{\Gamma, x : A \vdash p} \text{ CTR}$$

$$\frac{\Gamma, x : A \vdash B}{\Gamma \vdash A \supset B} \text{ RI}$$

$$\frac{\Gamma \vdash A \quad \Gamma; B \vdash p}{\Gamma; A \supset B \vdash p} \text{ LI}$$

Proof Terms and Proof Lists

$$t, u ::= \lambda x^A. t \mid x \mid l \quad l ::= \star \mid u :: l$$

Proof Terms and Proof Lists

$$t, u ::= \lambda x^A. t \mid x \mid l \quad l ::= \star \mid u :: l$$

Total Sequents

$$\Gamma \vdash t : A \quad \Gamma ; l : A \vdash p$$

Proof Terms and Proof Lists

$$t, u ::= \lambda x^A. t \mid x \mid l \quad l ::= \star \mid u :: l$$

Total Sequents

$$\Gamma \vdash t : A \quad \Gamma ; l : A \vdash p$$

Inference Rules

$$\frac{\Gamma, x : A \vdash t : B}{\Gamma \vdash \lambda x^A. t : A \supset B} \text{ RI}$$

$$\frac{\Gamma, x : A ; l : A \vdash p}{\Gamma, x : A \vdash x \mid l : p} \text{ CTR}$$

$$\frac{}{\Gamma ; \star : p \vdash p} \text{ AX}$$

$$\frac{\Gamma \vdash u : A \quad \Gamma ; l : B \vdash p}{\Gamma, u :: l : A \supset B \vdash p} \text{ LI}$$

Proof Terms and Proof Lists

$$t, u ::= \lambda x^A. t \mid x \mid l \quad l ::= \star \mid u :: l$$

Total Sequents

$$\Gamma \vdash t : A \quad \Gamma; l : A \vdash p$$

Example

$$\frac{\frac{\frac{\overline{\Gamma; \star : p \vdash p} \text{ AX}}{\Gamma \vdash x^{\star} : p} \text{ CTR} \quad \frac{\overline{\Gamma; \star : p \vdash p} \text{ AX}}{\Gamma \vdash x^{\star} : p} \text{ LI}}{\Gamma; ((x^{\star}) :: \star) : p \supset p \vdash p} \text{ CTR}}{\Gamma \vdash f^{\wedge}((x^{\star}) :: \star) : p} \text{ CTR} \\ \frac{}{\vdash \lambda f^{(p \supset p)}. \lambda x^p. f^{\wedge}((x^{\star}) :: \star) : (p \supset p) \supset p \supset p} \text{ RI}$$

where $\Gamma := \{f : p \supset p, x : p\}$

Proof Search (Focusing)

1. Eagerly invert RI to decompose implications in the RHS.

$$\frac{\Gamma, x : A \vdash B}{\Gamma \vdash A \supset B} \text{ RI}$$

Proof Search (Focusing)

1. Eagerly invert RI to decompose implications in the RHS.
2. When the RHS becomes atomic, apply CTR to decide a formula in the LHS to focus on.

$$\frac{\Gamma, x : A \vdash B}{\Gamma \vdash A \supset B} \text{ } RI \qquad \frac{\Gamma, x : A; A \vdash p}{\Gamma, x : A \vdash p} \text{ } CTR$$

Proof Search (Focusing)

1. Eagerly invert RI to decompose implications in the RHS.
2. When the RHS becomes atomic, apply CTR to decide a formula in the LHS to focus on.
3. As long as the focus is an implication, apply LI, keeping the focus on the succedent formula, and launching sub-problems.

$$\frac{\Gamma, x : A \vdash B}{\Gamma \vdash A \supset B} \text{ RI}$$

$$\frac{\Gamma, x : A; A \vdash p}{\Gamma, x : A \vdash p} \text{ CTR}$$

$$\frac{\Gamma \vdash A \quad \Gamma; B \vdash p}{\Gamma; A \supset B \vdash p} \text{ LI}$$

Proof Search (Focusing)

1. Eagerly invert RI to decompose implications in the RHS.
2. When the RHS becomes atomic, apply CTR to decide a formula in the LHS to focus on.
3. As long as the focus is an implication, apply LI, keeping the focus on the succedent formula, and launching sub-problems.
4. When the focus becomes atomic, compare with the atom in the RHS; if equal, finish with AX, otherwise backtrack to 2.

$$\frac{\Gamma, x : A \vdash B}{\Gamma \vdash A \supset B} \text{ RI}$$

$$\frac{\Gamma, x : A; A \vdash p}{\Gamma, x : A \vdash p} \text{ CTR}$$

$$\frac{}{\Gamma; p \vdash p} \text{ AX}$$

$$\frac{\Gamma \vdash A \quad \Gamma; B \vdash p}{\Gamma; A \supset B \vdash p} \text{ LI}$$

- Sequents of LJ_T: $\sigma = (\Gamma \vdash A)$ $\tau = (\Gamma; A \vdash p)$

From LJT to LJT_∂

- Sequents of LJT: $\sigma = (\Gamma \vdash A)$ $\tau = (\Gamma; A \vdash p)$
- Formal Sequents:** $\underline{\sigma} = (\underline{\Gamma \vdash A})$ $\underline{\tau} = (\underline{\Gamma; A \vdash p})$

From LJ_T to LJ_T_∂

- Sequents of LJ_T: $\sigma = (\Gamma \vdash A)$ $\tau = (\Gamma; A \vdash p)$
- **Formal Sequents:** $\underline{\sigma} = (\underline{\Gamma \vdash A})$ $\underline{\tau} = (\underline{\Gamma; A \vdash p})$
- Partial Proof Terms: proof terms with occurrences of formal sequents;

From LJT to LJT_{∂}

- Sequents of LJT: $\sigma = (\Gamma \vdash A)$ $\tau = (\Gamma; A \vdash p)$
- **Formal Sequents:** $\underline{\sigma} = (\underline{\Gamma \vdash A})$ $\underline{\tau} = (\underline{\Gamma; A \vdash p})$
- **Partial Proof Terms:** proof terms with occurrences of formal sequents;
- **Partial Derivations:** derivations in LJT_{∂} which formalize incomplete proofs found in the proof search process;

From LJT to LJT_{∂}

- Sequents of LJT: $\sigma = (\Gamma \vdash A)$ $\tau = (\Gamma; A \vdash p)$
- **Formal Sequents:** $\underline{\sigma} = (\underline{\Gamma \vdash A})$ $\underline{\tau} = (\underline{\Gamma; A \vdash p})$
- **Partial Proof Terms:** proof terms with occurrences of formal sequents;
- **Partial Derivations:** derivations in LJT_{∂} which formalize incomplete proofs found in the proof search process;
- **Rewriting System:** reduction relation on partial proof terms expressing proof search, such that **total terms are normal forms**.

From LJT to LJT_{∂}

- Sequents of LJT: $\sigma = (\Gamma \vdash A)$ $\tau = (\Gamma; A \vdash p)$
- **Formal Sequents:** $\underline{\sigma} = (\underline{\Gamma \vdash A})$ $\underline{\tau} = (\underline{\Gamma; A \vdash p})$
- **Partial Proof Terms:** proof terms with occurrences of formal sequents;
- **Partial Derivations:** derivations in LJT_{∂} which formalize incomplete proofs found in the proof search process;
- **Rewriting System:** reduction relation on partial proof terms expressing proof search, such that **total terms are normal forms**.

Aim: Proof Search as Normalization

LJT derives $\Gamma \vdash t : A$ iff $\underline{\Gamma \vdash A} \twoheadrightarrow t$

From LJT to LJT_{∂}

- Sequents of LJT: $\sigma = (\Gamma \vdash A)$ $\tau = (\Gamma; A \vdash p)$
- **Formal Sequents:** $\underline{\sigma} = (\underline{\Gamma \vdash A})$ $\underline{\tau} = (\underline{\Gamma; A \vdash p})$
- **Partial Proof Terms:** proof terms with occurrences of formal sequents;
- **Partial Derivations:** derivations in LJT_{∂} which formalize incomplete proofs found in the proof search process;
- **Rewriting System:** reduction relation on partial proof terms expressing proof search, such that **total terms are normal forms**.

Aim: Proof Search as Normalization

LJT derives $\Gamma \vdash t : A$ iff $\underline{\Gamma \vdash A} \rightarrow t$
proof search succeeds

From LJT to LJT_{∂}

- Sequents of LJT: $\sigma = (\Gamma \vdash A)$ $\tau = (\Gamma; A \vdash p)$
- **Formal Sequents:** $\underline{\sigma} = (\underline{\Gamma \vdash A})$ $\underline{\tau} = (\underline{\Gamma; A \vdash p})$
- **Partial Proof Terms:** proof terms with occurrences of formal sequents;
- **Partial Derivations:** derivations in LJT_{∂} which formalize incomplete proofs found in the proof search process;
- **Rewriting System:** reduction relation on partial proof terms expressing proof search, such that **total terms are normal forms**.

Aim: Proof Search as Normalization

LJT derives $\Gamma \vdash t : A$
proof search succeeds

iff

$\underline{\Gamma \vdash A} \rightarrow t$
the formal sequent normalizes

Proof Terms and

Proof Lists

Proof Terms

Proof Lists

$t, u ::= \lambda x^A. t \mid x^{\wedge} l \mid$

$l ::= \star \mid u :: l \mid$

Partial Proof Terms and Partial Proof Lists

Partial Proof Terms

Partial Proof Lists

$t, u ::= \lambda x^A. t \mid x^{\wedge} l \mid \underline{\sigma}$

$l ::= \star \mid u :: l \mid \underline{\tau}$

Partial Proof Terms and Partial Proof Lists

Partial Proof Terms

Partial Proof Lists

$t, u ::= \lambda x^A. t \mid x^{\wedge} l \mid \underline{\sigma}$

$l ::= \star \mid u :: l \mid \underline{\tau}$

Formal Term Sequents

Formal List Sequents

$\underline{\sigma} ::= \frac{}{\Gamma \vdash A}$

$\underline{\tau} ::= \frac{}{\Gamma; A \vdash p}$

LJT_∂ : partial proof terms

Partial Proof Terms and Partial Proof Lists

Partial Proof Terms

Partial Proof Lists

$$t, u ::= \lambda x^A. t \mid x^{\wedge} l \mid \underline{\sigma}$$

$$l ::= \star \mid u :: l \mid \underline{\tau}$$

Formal Term Sequents

Formal List Sequents

$$\underline{\sigma} ::= \frac{}{\Gamma \vdash A}$$

$$\underline{\tau} ::= \frac{}{\Gamma; A \vdash p}$$

Examples

Partial Proof Term

Partial Proof List

$$\lambda f^{(p \supset p)}. \lambda x^p. (\Gamma \vdash p)$$

$$(x^{\wedge} \star) :: (\underline{\Gamma; p \vdash p})$$

Partial Sequents

$$\Xi \Vdash t : \sigma \qquad \Xi \Vdash l : \tau$$

Partial Sequents

Proof State

$\Xi \Vdash t : \sigma$

$\Xi \Vdash l : \tau$

Partial Sequents

Proof State

proof obligations
record of the search
goal sequent

$\Xi \Vdash t : \sigma$

$\Xi \Vdash l : \tau$

list of formal sequents
partial proof term/list
 σ/τ

Ξ
 t/l

Partial Sequents

Proof State

proof obligations
record of the search
goal sequent

$\Xi \Vdash t : \sigma$

list of formal sequents
partial proof term/list
 σ/τ

$\Xi \Vdash l : \tau$

Ξ
 t/l

Example

Proof State

proof obligations
record of the search
goal sequent

$[\underline{\Gamma \vdash p}] \Vdash \lambda f^{(p \supset p)}. \lambda x^p. (\underline{\Gamma \vdash p}) : (\vdash (p \supset p) \supset p \supset p)$
where $\Gamma := \{f : p \supset p, x : p\}$

list of formal sequents
partial proof term/list
 $\vdash (p \supset p) \supset p \supset p$

$\frac{[\underline{\Gamma \vdash p}]}{\lambda f^{(p \supset p)}. \lambda x^p. (\underline{\Gamma \vdash p})}$

Inference Rules

$$\overline{[\sigma] \Vdash \underline{\sigma} : \sigma} \quad \partial$$

$$\frac{}{\epsilon \Vdash \star : (\Gamma; p \vdash p)} \quad AX$$

$$\frac{\Xi \Vdash t : (\Gamma, x : A \vdash B)}{\Xi \Vdash \lambda x^A. t : (\Gamma \vdash A \supset B)} \quad RI$$

$$\overline{[\tau] \Vdash \underline{\tau} : \tau} \quad \partial$$

$$\frac{\Xi \Vdash l : (\Gamma, x : A; A \vdash p)}{\Xi \Vdash x^l : (\Gamma, x : A \vdash p)} \quad CTR$$

$$\frac{\Xi_1 \Vdash u : (\Gamma \vdash A) \quad \Xi_2 \Vdash l : (\Gamma; B \vdash p)}{\Xi_1 @ \Xi_2 \Vdash u :: l : (\Gamma; A \supset B \vdash p)} \quad LI$$

$$\begin{array}{llll} (IR) & \frac{\Gamma \vdash A \supset B}{\Gamma \vdash A \supset B} & \rightarrow & \lambda x^A. (\Gamma, x : A \vdash B) \\ (SFL) & \frac{\Gamma, x : A \vdash p}{\Gamma, x : A \vdash p} & \rightarrow & x^{\wedge}(\Gamma, x : A; A \vdash p) \\ (KFL) & \frac{\Gamma; A \supset B \vdash p}{\Gamma; A \supset B \vdash p} & \rightarrow & (\Gamma \vdash A) :: (\Gamma; B \vdash p) \\ (FFL) & \frac{\Gamma; p \vdash p}{\Gamma; p \vdash p} & \rightarrow & \star \end{array}$$

$$\begin{array}{llll} (IR) & \frac{\Gamma \vdash A \supset B}{\Gamma \vdash A \supset B} & \rightarrow & \lambda x^A. (\Gamma, x : A \vdash B) \\ (SFL) & \frac{\Gamma, x : A \vdash p}{\Gamma, x : A \vdash p} & \rightarrow & x^{\wedge}(\Gamma, x : A; A \vdash p) \\ (KFL) & \frac{\Gamma; A \supset B \vdash p}{\Gamma; A \supset B \vdash p} & \rightarrow & (\frac{\Gamma \vdash A}{\Gamma \vdash A}) :: (\frac{\Gamma; B \vdash p}{\Gamma; B \vdash p}) \\ (FFL) & \frac{\Gamma; p \vdash p}{\Gamma; p \vdash p} & \rightarrow & \star \end{array}$$

Comments

$$\begin{array}{lll} (IR) & \frac{\Gamma \vdash A \supset B}{\Gamma \vdash A \supset B} & \rightarrow \lambda x^A. (\Gamma, x : A \vdash B) \\ (SFL) & \frac{\Gamma, x : A \vdash p}{\Gamma, x : A \vdash p} & \rightarrow x^{\wedge}(\Gamma, x : A; A \vdash p) \\ (KFL) & \frac{\Gamma; A \supset B \vdash p}{\Gamma; A \supset B \vdash p} & \rightarrow (\frac{\Gamma \vdash A}{\Gamma \vdash A}) :: (\frac{\Gamma; B \vdash p}{\Gamma; B \vdash p}) \\ (FFL) & \frac{\Gamma; p \vdash p}{\Gamma; p \vdash p} & \rightarrow \star \end{array}$$

Comments

- One reduction rule for each inference rule;

$$\begin{array}{lll}
 (IR) & \frac{\Gamma \vdash A \supset B}{\Gamma \vdash A \supset B} & \rightarrow \lambda x^A. (\Gamma, x : A \vdash B) \\
 (SFL) & \frac{\Gamma, x : A \vdash p}{\Gamma, x : A \vdash p} & \rightarrow x^{\wedge}(\Gamma, x : A; A \vdash p) \\
 (KFL) & \frac{\Gamma; A \supset B \vdash p}{\Gamma; A \supset B \vdash p} & \rightarrow (\frac{\Gamma \vdash A}{\Gamma \vdash A}) :: (\frac{\Gamma; B \vdash p}{\Gamma; B \vdash p}) \\
 (FFL) & \frac{\Gamma; p \vdash p}{\Gamma; p \vdash p} & \rightarrow \star
 \end{array}$$

Comments

- One reduction rule for each inference rule;
- The reduction rule tells how to use the inference rule during proof search;

$$\begin{array}{lll} (IR) & \frac{\Gamma \vdash A \supset B}{\Gamma \vdash A \supset B} & \rightarrow \lambda x^A. (\Gamma, x : A \vdash B) \\ (SFL) & \frac{\Gamma, x : A \vdash p}{\Gamma, x : A \vdash p} & \rightarrow x^{\wedge}(\Gamma, x : A; A \vdash p) \\ (KFL) & \frac{\Gamma; A \supset B \vdash p}{\Gamma; A \supset B \vdash p} & \rightarrow (\frac{\Gamma \vdash A}{\Gamma \vdash A}) :: (\frac{\Gamma; B \vdash p}{\Gamma; B \vdash p}) \\ (FFL) & \frac{\Gamma; p \vdash p}{\Gamma; p \vdash p} & \rightarrow \star \end{array}$$

Comments

- One reduction rule for each inference rule;
- The reduction rule tells how to use the inference rule during proof search;
- Redexes are formal sequents;

$$\begin{array}{lll} (IR) & \frac{\Gamma \vdash A \supset B}{\Gamma \vdash A \supset B} & \rightarrow \lambda x^A. (\Gamma, x : A \vdash B) \\ (SFL) & \frac{\Gamma, x : A \vdash p}{\Gamma, x : A \vdash p} & \rightarrow x^{\wedge}(\Gamma, x : A; A \vdash p) \\ (KFL) & \frac{\Gamma; A \supset B \vdash p}{\Gamma; A \supset B \vdash p} & \rightarrow (\frac{\Gamma \vdash A}{\Gamma \vdash A}) :: (\frac{\Gamma; B \vdash p}{\Gamma; B \vdash p}) \\ (FFL) & \frac{\Gamma; p \vdash p}{\Gamma; p \vdash p} & \rightarrow \star \end{array}$$

Comments

- One reduction rule for each inference rule;
- The reduction rule tells how to use the inference rule during proof search;
- Redexes are formal sequents;
- Reduction builds the record of the search.

LJT_∂ : example

Informal LJ_T Derivation

$$\frac{}{\vdash (p \supset p) \supset p \supset p} \partial$$

Reduction Sequence

Proof State

LJT_∂ : example

Informal LJT Derivation

$$\frac{}{\vdash (p \supset p) \supset p \supset p} \partial$$

Reduction Sequence

$$\frac{}{\vdash (p \supset p) \supset p \supset p}$$

Proof State

LJT_∂ : example

Informal LJT Derivation

$$\frac{}{\vdash (p \supset p) \supset p \supset p} \partial$$

Reduction Sequence

$$\frac{}{\vdash (p \supset p) \supset p \supset p}$$

Proof State

$$[\frac{}{\vdash (p \supset p) \supset p \supset p}] \Vdash \frac{}{\vdash (p \supset p) \supset p \supset p} : (\vdash (p \supset p) \supset p \supset p)$$

LJT_∂ : example

Informal LJTDerivation

$$\frac{\frac{\overline{\Gamma \vdash p} \quad \partial}{\vdash (p \supset p) \supset p \supset p} \quad RI}$$

$$\Gamma := \{f : p \supset p, x : p\}$$

Reduction Sequence

$$\underline{\vdash (p \supset p) \supset p \supset p}$$

Proof State

Informal LJTDerivation

$$\frac{\frac{\overline{\Gamma \vdash p} \quad \partial}{\vdash (p \supset p) \supset p \supset p} \quad RI}{\vdash (p \supset p) \supset p \supset p} \quad RI$$

$$\Gamma := \{f : p \supset p, x : p\}$$

Reduction Sequence

$$\xrightarrow{\rightarrow_{IR}^2} \frac{\vdash (p \supset p) \supset p \supset p}{\lambda f (p \supset p) . \lambda x p . (\Gamma \vdash p)}$$

Proof State

Informal LJTDerivation

$$\frac{\frac{\overline{\Gamma \vdash p} \quad \partial}{\vdash (p \supset p) \supset p \supset p} \quad RI}{\vdash (p \supset p) \supset p \supset p}$$

$$\Gamma := \{f : p \supset p, x : p\}$$

Reduction Sequence

$$\xrightarrow{2}_{IR} \frac{\vdash (p \supset p) \supset p \supset p}{\lambda f (p \supset p) . \lambda x p . (\underline{\Gamma \vdash p})}$$

Proof State

$$[(\underline{\Gamma \vdash p})] \Vdash \lambda f (p \supset p) . \lambda x p . (\underline{\Gamma \vdash p}) : (\vdash (p \supset p) \supset p \supset p)$$

LJT_∂ : example

Informal LJTDerivation

$$\frac{\frac{\frac{\overline{\Gamma \vdash p} \quad \partial \quad \overline{\Gamma; p \vdash p} \quad \partial}{\Gamma; p \supset p \vdash p} LI}{\Gamma \vdash p} CTR}{\vdash (p \supset p) \supset p \supset p} RI$$

$$\Gamma := \{f : p \supset p, x : p\}$$

Reduction Sequence

$$\xrightarrow{2}_{IR} \frac{\vdash (p \supset p) \supset p \supset p}{\lambda f (p \supset p) . \lambda x p . (\Gamma \vdash p)}$$

Proof State

LJT_∂ : example

Informal LJTDerivation

$$\frac{\frac{\frac{\frac{}{\Gamma \vdash p} \partial \quad \frac{}{\Gamma; p \vdash p} \partial}{\Gamma; p \supset p \vdash p} LI}{\Gamma \vdash p} CTR}{\vdash (p \supset p) \supset p \supset p} RI$$

$$\Gamma := \{f : p \supset p, x : p\}$$

Reduction Sequence

$$\begin{aligned} & \vdash (p \supset p) \supset p \supset p \\ \rightarrow_{IR}^2 & \lambda f (p \supset p) . \lambda x P . (\Gamma \vdash p) \\ \rightarrow_{SFL} & \lambda f (p \supset p) . \lambda x P . f^*(\Gamma; p \supset p \vdash p) \\ \rightarrow_{KFL} & \lambda f (p \supset p) . \lambda x P . f^*((\Gamma \vdash p) :: (\Gamma; p \vdash p)) \end{aligned}$$

Proof State

Informal LJTDerivation

$$\begin{array}{c}
 \frac{\frac{\frac{}{\Gamma \vdash p} \partial \quad \frac{}{\Gamma; p \vdash p} \partial}{\Gamma; p \supset p \vdash p} LI}{\Gamma \vdash p} CTR \\
 \hline
 \vdash (p \supset p) \supset p \supset p \quad RI
 \end{array}$$

$$\Gamma := \{f : p \supset p, x : p\}$$

Reduction Sequence

$$\begin{array}{l}
 \vdash (p \supset p) \supset p \supset p \\
 \rightarrow_{IR}^2 \quad \lambda f^{(p \supset p)}. \lambda x^p. (\underline{\Gamma \vdash p}) \\
 \rightarrow_{SFL} \quad \lambda f^{(p \supset p)}. \lambda x^p. f^*(\underline{\Gamma; p \supset p \vdash p}) \\
 \rightarrow_{KFL} \quad \lambda f^{(p \supset p)}. \lambda x^p. f^*(\underline{(\Gamma \vdash p)} :: \underline{(\Gamma; p \vdash p)})
 \end{array}$$

Proof State

$$[(\underline{\Gamma \vdash p}), (\underline{\Gamma; p \vdash p})] \Vdash \lambda f^{(p \supset p)}. \lambda x^p. f^*(\underline{(\Gamma \vdash p)} :: \underline{(\Gamma; p \vdash p)}) : (\vdash (p \supset p) \supset p \supset p)$$

LJT_∂ : example

Informal LJTDerivation

$$\frac{\frac{\frac{}{\Gamma; p \vdash p} \text{AX}}{\Gamma \vdash p} \text{CTR} \quad \frac{}{\Gamma; p \vdash p} \text{AX}}{\frac{}{\Gamma; p \supset p \vdash p} \text{CTR}} \text{LI} \quad \frac{}{\Gamma \vdash p} \text{CTR} \quad \frac{}{\vdash (p \supset p) \supset p \supset p} \text{RI}$$

$$\Gamma := \{f : p \supset p, x : p\}$$

Reduction Sequence

$$\begin{aligned} & \vdash (p \supset p) \supset p \supset p \\ \rightarrow_{IR}^2 & \lambda f (p \supset p) . \lambda x P . (\Gamma \vdash p) \\ \rightarrow_{SFL} & \lambda f (p \supset p) . \lambda x P . f^*(\Gamma; p \supset p \vdash p) \\ \rightarrow_{KFL} & \lambda f (p \supset p) . \lambda x P . f^*((\Gamma \vdash p) :: (\Gamma; p \vdash p)) \end{aligned}$$

Proof State

Informal LJTDerivation

$$\begin{array}{c}
 \frac{}{\Gamma; p \vdash p} \text{AX} \\
 \frac{}{\Gamma \vdash p} \text{CTR} \quad \frac{}{\Gamma; p \vdash p} \text{AX} \\
 \frac{}{\Gamma; p \supset p \vdash p} \text{LI} \\
 \frac{}{\Gamma \vdash p} \text{CTR} \\
 \frac{}{\vdash (p \supset p) \supset p \supset p} \text{RI}
 \end{array}$$

$$\Gamma := \{f : p \supset p, x : p\}$$

Reduction Sequence

$$\begin{array}{l}
 \vdash (p \supset p) \supset p \supset p \\
 \xrightarrow{2} \text{IR} \quad \lambda f (p \supset p) . \lambda x P . (\Gamma \vdash p) \\
 \xrightarrow{\text{SFL}} \quad \lambda f (p \supset p) . \lambda x P . f^*(\Gamma; p \supset p \vdash p) \\
 \xrightarrow{\text{KFL}} \quad \lambda f (p \supset p) . \lambda x P . f^*((\Gamma \vdash p) :: (\Gamma; p \vdash p)) \\
 \xrightarrow{\text{FFL}} \quad \lambda f (p \supset p) . \lambda x P . f^*((\Gamma \vdash p) :: *) \\
 \xrightarrow{\text{SFL}} \quad \lambda f (p \supset p) . \lambda x P . f^*((x^*(\Gamma; p \vdash p)) :: *) \\
 \xrightarrow{\text{FFL}} \quad \lambda f (p \supset p) . \lambda x P . f^*((x^* *) :: *)
 \end{array}$$

Proof State

LJT_∂ : example

Informal LJTDerivation

$$\begin{array}{c}
 \frac{}{\Gamma; p \vdash p} \text{AX} \\
 \frac{}{\Gamma \vdash p} \text{CTR} \quad \frac{}{\Gamma; p \vdash p} \text{AX} \\
 \frac{}{\Gamma; p \supset p \vdash p} \text{LI} \\
 \frac{}{\Gamma \vdash p} \text{CTR} \\
 \frac{}{\vdash (p \supset p) \supset p \supset p} \text{RI}
 \end{array}$$

$$\Gamma := \{f : p \supset p, x : p\}$$

Reduction Sequence

$$\begin{array}{l}
 \vdash (p \supset p) \supset p \supset p \\
 \xrightarrow{2} \text{IR} \quad \lambda f (p \supset p) . \lambda x P . (\Gamma \vdash p) \\
 \rightarrow \text{SFL} \quad \lambda f (p \supset p) . \lambda x P . f^*(\Gamma; p \supset p \vdash p) \\
 \rightarrow \text{KFL} \quad \lambda f (p \supset p) . \lambda x P . f^*((\Gamma \vdash p) :: (\Gamma; p \vdash p)) \\
 \rightarrow \text{FFL} \quad \lambda f (p \supset p) . \lambda x P . f^*((\Gamma \vdash p) :: *) \\
 \rightarrow \text{SFL} \quad \lambda f (p \supset p) . \lambda x P . f^*((x^*(\Gamma; p \vdash p)) :: *) \\
 \rightarrow \text{FFL} \quad \lambda f (p \supset p) . \lambda x P . f^*((x^* *) :: *)
 \end{array}$$

Proof State

$$\epsilon \Vdash \lambda f (p \supset p) . \lambda x P . f^*((x^* *) :: *) : (\vdash (p \supset p) \supset p \supset p)$$

Conservativity

- LJT derives $\Gamma \vdash t : A$ iff LJT_{∂} derives $\epsilon \Vdash t : (\Gamma \vdash A)$.
- LJT derives $\Gamma ; I : A \vdash p$ iff LJT_{∂} derives $\epsilon \Vdash I : (\Gamma ; A \vdash p)$.

Conservativity

- LJT derives $\Gamma \vdash t : A$ iff LJT_{∂} derives $\epsilon \Vdash t : (\Gamma \vdash A)$.
- LJT derives $\Gamma; I : A \vdash p$ iff LJT_{∂} derives $\epsilon \Vdash I : (\Gamma; A \vdash p)$.

Record of Search

- If $\Xi \Vdash t : \sigma$ then $\underline{\sigma} \twoheadrightarrow t$.
- If $\Xi \Vdash I : \tau$ then $\underline{\tau} \twoheadrightarrow I$.

Conservativity

- LJT derives $\Gamma \vdash t : A$ iff LJT_{∂} derives $\epsilon \Vdash t : (\Gamma \vdash A)$.
- LJT derives $\Gamma; I : A \vdash p$ iff LJT_{∂} derives $\epsilon \Vdash I : (\Gamma; A \vdash p)$.

Record of Search

- If $\Xi \Vdash t : \sigma$ then $\underline{\sigma} \rightarrow t$.
- If $\Xi \Vdash I : \tau$ then $\underline{\tau} \rightarrow I$.

Subject Reduction

- If $\Xi \Vdash t : \sigma$ and $t \rightarrow t'$, then there is Ξ' such that $\Xi \rightarrow \Xi'$ and $\Xi' \Vdash t' : \sigma$.
- If $\Xi \Vdash I : \tau$ and $I \rightarrow I'$, then there is Ξ' such that $\Xi \rightarrow \Xi'$ and $\Xi' \Vdash I' : \tau$.

Proof Search as Normalization

- For all $t \in LJT$: LJT derives $\Gamma \vdash t : A$ iff $\underline{\sigma} \rightarrow t$, where $\underline{\sigma} = \underline{\Gamma \vdash A}$.
- For all $l \in LJT$: LJT derives $\Gamma; l : A \vdash p$ iff $\underline{\tau} \rightarrow l$, where $\underline{\tau} = \underline{\Gamma; A \vdash p}$.

Second case study: NJT

- Bidirectional Natural Deduction system;

Second case study: NJT

- Bidirectional Natural Deduction system;
- Proof Search combines bottom-up and top-down reasoning, following the ideas of the intercalation calculus [W. Sieg];

Second case study: NJT

- Bidirectional Natural Deduction system;
- Proof Search combines bottom-up and top-down reasoning, following the ideas of the intercalation calculus [W. Sieg];
- ∂ -rule applications represent not only totally undefined derivations, but also gaps between formulas already obtained top-down and an atom previously obtained bottom-up;

Second case study: NJT

- Bidirectional Natural Deduction system;
- Proof Search combines bottom-up and top-down reasoning, following the ideas of the intercalation calculus [W. Sieg];
- ∂ -rule applications represent not only totally undefined derivations, but also gaps between formulas already obtained top-down and an atom previously obtained bottom-up;

Consequences:

Proof Terms - one kind of formal sequent does not constitute, by itself, a partial proof term;

Partial Deductive System - the second ∂ -rule becomes more complex;

Rewriting System - some redexes are not formal sequents.

Second case study: NJT

- Bidirectional Natural Deduction system;
- Proof Search combines bottom-up and top-down reasoning, following the ideas of the intercalation calculus [W. Sieg];
- ∂ -rule applications represent not only totally undefined derivations, but also gaps between formulas already obtained top-down and an atom previously obtained bottom-up;

Consequences:

Proof Terms - one kind of formal sequent does not constitute, by itself, a partial proof term;

Partial Deductive System - the second ∂ -rule becomes more complex;

Rewriting System - some redexes are not formal sequents.

Theorem of Idealized Proof Search

Proof search in LJ_T (following *focusing*)
is **isomorphic** to
proof search in NJ_T (following *intercalation*).

Second case study: NJT

- Bidirectional Natural Deduction system;
- Proof Search combines bottom-up and top-down reasoning, following the ideas of the intercalation calculus [W. Sieg];
- ∂ -rule applications represent not only totally undefined derivations, but also gaps between formulas already obtained top-down and an atom previously obtained bottom-up;

Consequences:

Proof Terms - one kind of formal sequent does not constitute, by itself, a partial proof term;

Partial Deductive System - the second ∂ -rule becomes more complex;

Rewriting System - some redexes are not formal sequents.

Theorem of Idealized Proof Search

Proof search in LJ_T (following *focusing*)
is **isomorphic** to
proof search in NJ_T (following *intercalation*).

Espírito Santo, J., Sousa, A.C. (2024). Partial proof terms in the study of idealized proof search. In: Intelligent Computer Mathematics. CICM 2024. (*to appear*)