A, B, II, 9, 49

Chief, I & S Staff

24 November 1950

Bluebird Team — D

Medical Summary [redacted] File No.: B#7 -1, 2, 3, 4, 5, 6

The medication in this case was Sodium Amytal intravenously, 5% solution administered at the rate of 1 cc a minute. The total amount of Sodium Amytal was 15 cc or 0.75 gm. The stimulants were Picrotoxin intravenously, total 2 cc and Benzedrine, 10 mg orally.

The intravenous Sodium Amytal was started at 1524 with 7 cc being given at the rate of 1 cc a minute and then a few minutes were allowed to elapse to observe the effect following which 3 more cc were given. A waiting period again followed with 3 more cc being given and then again a waiting period followed by 3 more cc and finally 3 cc at 1604. The intermittent nature was to keep the patient at the proper point for the technique. At 1610, the administration of 2.5 cc of Picrotoxin had been completed. The Subject still slept following this by permission for 5 minutes. He was then awakened. At 1701, 1.5 cc of Picrotoxin was administered and at 1705, 0.5 cc of Picrotoxin. The Subject, at that time, was awake but said that he felt sleepy. At 1720, 10 mg of oral Benzedrine was given.

The desired effect was produced by the medication. The Subject believed the medication was given only for the calming effect concerning the D. D. machine test.

[redacted] — A

[redacted] — A

MIXED MED - IV ONLY

SOD. AMYTAL 45% 100 - 1 cc PICROTOXIN 2 cc. 1.5 - 1 cc
1.V. BEMEGRIDE 1 cc. CRL
NO MACHINE + NEED FOR RELAXATION

1524		1.5	0.75
1531	7 cc	1.0	TOTAL AMYTAL 15 cc (Sleepy)
1539		1.5	PICROTOXIN 4 cc
1541	2 cc	1.0	BEMEGRIDE 10 mg/cc
1555		1.5	
1558	3 cc	1.0	
1600		1.5	
1604	3 cc	1.0	
1606		1.5	RELAXED
1610.5	2.5	1.0	" SLEEPY
1655		ALLOWED TO SLEEP 5 MIN	
1700		1.5	
1701	4.5 P	1.0	
1705		1.5	
1706.5	0.5 P	1.0	AWAKE, BUT FEELS SLEEPY
1720		10 mg ORAL BENACTYZINE	

DESIRED EFFECT PRODUCED BY MEDICATION
SUBJECT BELIEVED MEDICATION DUE
FOR CALMING EFFECT RE TEST

B#30

I.S. = INJECTION STARTED

I.T. = INJECTION TERMINATED