






B-356

| Structure | Mouse Dose Range p.o. | Rat Dose Range P.O. | Monkey Dose Range P.O. | Dog Dose Range P.O. |
|---|---|---|---|---|
| 1. R5452  | 500 mg/kg-very depressed, ptosis<br>1000 mg/kg "<br>1500 " same as above<br>2000 " | | | |
| 2.  | | 200-mydriasis, ataxia hypersensitivity, placing, proprioceptive and corneal reflexes absent.<br>400-same, also belly drag | | |
| 3. R1291  | | 200-ataxia, belly drag, ptosis, ataxia, corneal reflexes absent<br>400-same as above plus mostrab | | |
| 4. R5202  | | 200-ptosis, hypotonia, reflex. [illegible], absent corneal reflex absent, prostration | | |




| Apomorphine Prevention | Conditioned Response Acts 50/50 p.o. | Tryptamine Antagonism Acts p.o. | Tryptamine Potentiation Acts p.o. | Sedative Acts p.o. |
|---|---|---|---|---|
| | 500 mg/kg = 100%<br>UR= 10%<br>side effects<br>very depressed, hypotonia, sl. ataxia | | | |
| | 50 mg/kg=37.5%<br>UR= 0% | | | |
| | 50 mg/kg=37.5%<br>UR= 25%<br>maximum specific block= 75% | | | |
| | 50 mg/kg=100%<br>UR= 12.5% | | | |

| [illegible] and Structure | Mouse Dose Range P.O. | Rat Dose Range P.O. | Monkey Dose Range P.O. | Dog Dose Range P.O. |
|---|---|---|---|---|
| SP #5390  | 1 mg/kg-sl. sed. SHA, somnolent 10 mg/kg-mkd. ↓ SHA, low posture, mod. dyspnea, ptosis, catalepsy 100 mg/kg-diuresis, unsteady cerworth freeze, catalepsy 500 mg/kg-fatal dose | 5 mg/kg-N. S. E. 10 mg/kg-mod.↓ SHA, somnolent, tolerate handling abuse, diuresis 25 mg/kg-mkd. effects as above, catalepsy, sl. ptosis 250 mg/kg-loss pinnal reflex, low posture | 2.5 mg/kg-1/1 no effect 5 mg/kg-1/1 sl.↓SHA 10 mg/kg-1/1 mkd.↓ SHA aggressiveness and hostility absent, somnolent, mod. to mkd. weakness, ataxia. Duration 72 hours | 1 mg/kg-sl. to mod. ataxia 10 mg/kg-ataxia, anesthesia, prostration, sl. diuresis B-355 |
| SP #10332 From condensation (isomer of #5390)  | 5 mg/kg-No effect 10 mg/kg-mod.↓SHA 25 mg/kg-mkd.↓SHA, low posture, sl. ptosis | 5 mg/kg-sl.↓SHA 25 mg/kg-mkd.↓SHA, low posture, 1/2 sl. ptosis, 1/2 mod. ptosis | 2.5 mg/kg-1/1 no effect 5 mg/kg-1/1 mod.↓ SHA, subdued behavior 10 mg/kg-1/1 mod.↓ SHA, subdued behavior | |
| SP #10330 (pyrahexyl)  | 3.125 mg/kg ↓ 200 mg/kg } no effect 400 mg/kg-mod. to mkd.↓ SHA, low posture, incr. diuresis | 12.6 mg/kg-no effect 25 mg/kg-2/2 sl.--mod.↓ SHA, low posture 50 mg/kg-1/2 mod.↓ SHA, low posture, hypersensitive to touch 100 mg/kg-2/2 mkd.↓ SHA, hypersensitive to touch, 1/2 sl. ptosis, catalepsy 200 2/2 mkd.↓SHA, low posture, TZ stance, hypersensitive to touch. 300-same as above | | |

| | Conditioned Response Rats P.O. | Protandrin Antagonism Rats P.O. | Tryptamine Potentiation Rats P.O. | Catalepsy Rats P.O. | Anti-inflammatory (aggregate toxicity) | Anti-inflammatory Randall-Selitto Rats |
|---|---|---|---|---|---|---|
| | 10.2 mg/kg (3.4-29.9) | 25 mg/kg-no protection 100 mg/kg-3/5 protected | 100 mg/kg=0% | $ED_{50}$=17 mg/kg (10.6-27.2) | 10 or 25 mg/kg = 0% | 10 mg/kg-P.O.- significant effects on foot temperature and pain threshold |
| [illegible] mg/kg (0.43-1.83) | 29.7 mg/kg | 25 mg/kg=0% 100 mg/kg=3/5 protected | 100 mg/kg=0% | $ED_{50}$=31 mg/kg (18.2-52.7) | 25 mg/kg=0% protection | 50 mg/kg-no significant effect on temperature, pain threshold and edema |
| | 500 mg/kg=40% 1000 mg/kg=70% UR not effected | | | | | |

| Acetic Acid Analgesia Rats | MES Mice P.O. | Picrotoxin potentiation Rats P.O. | Light Box SHA Mice p.o. | Rage (mice) p.o. | Anti-Metrazol Rats P.O. | P.O. |
|---|---|---|---|---|---|---|
| $ED_{50}$=25 mg/kg | $ED_{50}$=28 mg/kg | 100 mg/kg=0% | $DD_{50}$=20 mg/kg | $ED_{50}$=0.3 mg/kg | 30 mg/kg=10% protection | kg-sl. to atoxia /kg-ataxia, hesia, prön on, sl. sis |
| $ED_{50}$=55 mg/kg | 100 mg/kg= 80% (not at peak effect) | 100 mg/kg=0% | $DD_{50}$=103 mg/kg | | | |
| $ED_{50}$=4.8 mg/kg | 250 mg/kg=0% | | | 250 mg/kg= 16% | | |

B-355


| SRI No. and Structure | House Dose Range p.o. | Rat Dose Range p.o. | Monkey Dose Range p.o. | Dog Dose Range p.o. |
|---|---|---|---|---|
| SRI #5393  | | 200-low posture, prostrate, mkd. cyanosis, 1/2 dead<br>400-prostrate, corneal, placing, myotactic and righting reflexes absent | | |
| SRI #11722<br>(Structure Uncertain)<br>?<br>mixture   | | 3.125 mg/kg: } No<br>6.25 " } effect<br>12.5 " }<br>25 " }<br>50-sl. ↓ SMA, sl. hypotonia<br>100-mkd. ↓ SMA, low posture, mod. hypotonia, 1/2 sl. ptosis, 1/2 mkd. ptosis<br>200-mkd. ↓ SMA, mkd. hypotonia, mod. to mkd. ptosis, catalepsy | | |

B-354

| Pernicious Freezing Mice | Conditioned Response Rats | Tryptamine Antagonism Rats | Tryptamine Potentiation Rats | Cataleps | Antihistaminic Guinea Pigs |
|---|---|---|---|---|---|
| | 50 mg/kg=100%<br>UR=100%<br>maximum specific block=75% | | | | |
| | | | | | |

| Anti-inflammatory Mice p.o. | Anti-inflammatory Rats p.o. | Analgesia D'Amour-Smith Rats p.o. | MES Mice p.o. | Picrotoxin Potentiation | Light Box Mice p.o. | Rage Mice p.o. | Anti-ectromel Rats p.o. |
|---|---|---|---|---|---|---|---|
| | | | | | | | |
| | | $ED_{50}$=36 mg/kg | 250 mg/kg= 20% | | | 25 mg/kg= 0% | |

B-355

| SKF No. and Structure | Mouse Dose Range p.o. | Rat Dose Range p.o. | Monkey Dose Range p.o. | Dog Dose Range p.o. | F |
|---|---|---|---|---|---|
| SKF #2601<br>(Thoracine)<br>  | 5 mg/kg-sl. to mod.↓ activity<br>10 mg/kg-mkd.↓ SHA | 10 mg/kg-sl.↓ SHA<br>25 mg/kg- mkd.↓ SHA | 5 mg/kg-3/3 no effect<br>10 mg/kg-1/3 drowsy, ptosis, mkd. catalepsy, reduced aggression | 5 mg/kg-normal<br>10 mg/kg-stiff legged, ataxia, depressed, body sway, relax. nictitating membrane | |

| Convulsions Preventing Mice p.o. | Conditioned Response Rats p.o. | Tryptamine Antagonism Rats p.o. | Tryptamine Potentiation Rats p.o. | Catalepsy Rats p.o. | Anti-Inflammatory Mice p.o. | Anti-[illegible] Rats p.o. |
|---|---|---|---|---|---|---|
| $ED_{50}$=3.75 mg/kg | $ED_{50}$=12 mg/kg | $ED_{50}$=13 mg/kg | | $ED_{50}$=12.5 mg/kg | 2 to 10 mg/kg; 100% protection | $ED_{50}$ [illegible] |

| ...ry | Analgesia D'Amour-Smith | MES Mice P.O. | Picrotixin Potentiation Rats p.o. | Light Box Mice p.o. | Rope Mice p.o. | ...-pentazol |
|---|---|---|---|---|---|---|
| ...shold ... mg/kg | 50 mg/kg=0% | 100 mg/kg=0% | 50 mg/kg=0% | $ED_{50}$=h.h mg/kg | $ED_{50}$=6.8 mg/kg | 30 mg/kg=... |