



MINISTRY OF ENVIRONMENT, CLIMATE CHANGE AND FORESTRY
KENYA METEOROLOGICAL
DEPARTMENT



Climate
Hazards
Center
UC SANTA BARBARA

Indian Ocean SST and Dipole Outlook

Forecast issued 19 September 2026, covering 19 September to 18 October 2026

ACCORD / CHC — draft for review

21 September 2026

Abstract

A positive Indian Ocean Dipole is forecast at every horizon over the coming month. The index is $+1.03\text{ }^{\circ}\text{C}$ in week 1 and $+0.86\text{ }^{\circ}\text{C}$ over the next 30 days. Six days earlier the same two figures were $+0.94$ and $+1.02\text{ }^{\circ}\text{C}$. The western box is 0.85 to $1.03\text{ }^{\circ}\text{C}$ above its 2006–2025 average and the eastern box is within $0.15\text{ }^{\circ}\text{C}$ of normal.

Forecasts come from the ECMWF sub-seasonal ensemble, corrected against NOAA OI SST by linear regression on the twenty past years of the same calendar date. At two of the ten box and window combinations the forecast is more accurate than 30-day persistence and the 80% interval on that difference stays above zero, both in the eastern box. At one of the two that interval reaches down to $+0.0002$ and so sits on zero.

The forecast has been too positive in all four windows that have verified, by 0.20 to $0.38\text{ }^{\circ}\text{C}$. In each of them the eastern correction cooled the forecast by 0.15 to $0.29\text{ }^{\circ}\text{C}$ and made it worse by 0.14 to $0.29\text{ }^{\circ}\text{C}$, the uncorrected value having been within $0.06\text{ }^{\circ}\text{C}$ of what was observed. A dipole computed as west minus east is inflated by an eastern value that is too cool, so the numbers above should be read as an upper estimate.

Two alternative methods were also tested. Neither is more accurate overall, but both give a weaker dipole for this run, and an ARIMA model fitted to the observed record alone is more accurate than the corrected forecast in week 1.

Method

We take the ECMWF sub-seasonal forecast issued on 19 September 2026, which runs with 100 members on a 1.5° grid, together with the matching reforecast for the same calendar date in each year from 2006 to 2025 at 10 members. Observations are NOAA OI SST v2.1 at 0.25° daily resolution. Both are averaged over the two dipole boxes of Saji et al. (1999), the west at $50\text{--}70^{\circ}\text{E}$ and $10^{\circ}\text{S}\text{--}10^{\circ}\text{N}$ and the east at $90\text{--}110^{\circ}\text{E}$ and $10^{\circ}\text{S}\text{--}0$, weighting each cell by the cosine of its latitude.

The lead windows are days 1 to 7, 8 to 14, 15 to 21, 22 to 28 and 1 to 30. Temperature is filed as a 24-hour mean, so day 1 is the day of issue and week 1 covers 19 to 25 September. An independent calculation labels day 1 as the following day.

For each box and window we fit a straight line of observed on forecast temperature across the twenty reforecast years and apply it to this year's forecast. Regression is used rather than quantile matching, which would pull values back inside the historical range. The 80% intervals are prediction intervals from that regression and describe how wrong this method has been in past years rather than the spread of the members. Accuracy figures are leave-one-year-out. Reforecasts exist only for odd calendar days of the month. The calculation is reproduced by `iod_pipeline.run("2026-09-19")`.




Figure 1: **(a)** Forecast anomaly in each box with 80% prediction intervals. **(b)** The dipole from the corrected and uncorrected forecasts, the gap being the change in amplitude made by the regression rather than the removal of a bias. **(c,d)** Leave-one-year-out correlation against two persistence baselines, shaded by the gain over the 30-day one.

Results

Horizon	Valid	West box	East box	Dipole
Week 1	19 – 25 Sep	28.92 °C (+1.03)	28.17 °C (–0.00)	+1.03 ± 0.56
Week 2	26 Sep – 2 Oct	28.94 °C (+0.89)	28.23 °C (+0.08)	+0.81 ± 0.52
Week 3	3 – 9 Oct	29.19 °C (+0.93)	28.30 °C (+0.14)	+0.79 ± 0.43
Week 4	10 – 16 Oct	29.23 °C (+0.85)	28.41 °C (+0.12)	+0.74 ± 0.42
Days 1–30	19 Sep – 18 Oct	29.10 °C (+0.94)	28.29 °C (+0.09)	+0.86 ± 0.35

Absolute temperatures, with the departure from the 2006–2025 average in brackets and the dipole with its 80% interval. That baseline is the reforecast period rather than the 1991–2020 climatology used by BoM, CPC and NOAA, so these anomalies are not comparable with the dipole values those centres publish. The uncorrected model dipole is +0.78 to +0.93. The regression changes amplitude rather than removing a bias, with western slopes of 0.83 to 1.12 and eastern slopes of 0.72 to 0.75.

Accuracy

Leave-one-year-out correlation across the twenty reforecast years against persistence, meaning the observed mean of the 31 days ending at issue, carried forward through the same leave-one-year-out regression. The gain carries an 80% bootstrap interval.

	Week 1	Week 2	Week 3	Week 4	Days 1–30
West forecast / persistence	0.83 / 0.84	0.84 / 0.84	0.80 / 0.70	0.76 / 0.72	0.87 / 0.85

	Week 1	Week 2	Week 3	Week 4	Days 1–30
gain	–0.00 (–0.07, +0.10)	–0.00 (–0.09, +0.10)	+0.10 (–0.04, +0.23)	+0.03 (–0.07, +0.18)	+0.02 (–0.06, +0.12)
East forecast / persistence	0.83 / 0.72	0.84 / 0.71	0.88 / 0.78	0.79 / 0.79	0.90 / 0.82
gain	+0.11 (–0.02, +0.23)	+0.13 (–0.01, +0.24)	+0.09 (+0.01, +0.16)	–0.00 (–0.11, +0.08)	+0.08 (+0.00, +0.15)

The gain is the forecast’s correlation minus persistence’s. Two of the ten box and window combinations have an interval above zero, both in the east, and the other eight include zero. The days 1 to 30 lower bound is shown as +0.00 above because its actual value is +0.0002, so that interval sits on zero rather than above it. The previous issuance had four above zero. Across four issuances the eastern gain at weeks 2 to 4 has been positive in eleven of twelve cases, with an interval above zero in six, and the western gain in one of twenty. Against 7-day persistence the forecast is less accurate in both boxes at week 1.




Figure 2: Corrected SST anomaly at every horizon, in °C against the 2006–2025 observed average. The western box is outlined in orange and the eastern in teal. Grey ocean marks cells where leave-one-year-out correlation falls below 0.4 and the forecast should not be relied on. Each cell is corrected by its own regression. Fields are interpolated for contouring, so the zero contour is placed by the interpolation rather than measured.

Alternative methods

Two alternatives were tested alongside the standing correction. Both use the observed ocean temperature at the time of issue, which the standing correction does not use.

The first fits observed temperature on the observed mean of the 30 days to issue and on the model’s forecast change from day 1, so the starting point comes from observations and the model supplies only the change. Out of sample it is

less accurate at every window in both boxes, and the weight on the model's change is not distinguishable from zero in the west.

The second is an ARIMA(1,0,1) model fitted to the daily observed record up to the day before issue, using no model data. Root mean square error in °C, on the seventeen years both methods cover, 2009 to 2025.

	Week 1	Week 2	Week 3	Week 4	Days 1–30
West forecast / ARIMA	0.196 / 0.117	0.167 / 0.207	0.198 / 0.294	0.240 / 0.337	0.139 / 0.210
East forecast / ARIMA	0.283 / 0.184	0.290 / 0.293	0.266 / 0.342	0.284 / 0.443	0.219 / 0.290

ARIMA is more accurate in week 1 in both boxes and less accurate from week 2 onward, consistent with the persistence comparison above. No interval is attached, and the smaller differences are not resolvable on seventeen years.

The comparison is uneven in both directions. ARIMA sees the observed temperature at issue and the regression does not, but the regression is refitted leaving out one year at a time from all twenty, so its prediction for an early year is informed by later ones. The seasonal cycle removed before the ARIMA fit is rebuilt from prior data only, which removes one bias and introduces another that runs cool. The two effects cannot be separated here. Under the alternative cycle the ordering is unchanged at weeks 3 and 4 and over days 1–30, while the east at week 2 reverses.

Dipole forecasts for this run, on the 2006–2025 average used in Results.

	Week 1	Week 2	Week 3	Week 4	Days 1–30
Standing correction	+1.03	+0.81	+0.79	+0.74	+0.86
Observed start, model change	+0.32	+0.17	+0.02	+0.11	+0.14
ARIMA	+0.54	+0.39	+0.20	+0.21	+0.33

Both are below the standing forecast, which the verification record also shows running too positive. Because ARIMA is a stronger comparison than flat persistence at week 1, the measured gain over persistence at that lead may overstate what the forecast adds. Neither is proposed as a replacement.

Verification record

run	valid	forecast	observed	error	east raw err	east cal err
31 Aug	31 Aug – 6 Sep	+0.43	+0.05	+0.38	–0.06	–0.35
31 Aug	7 – 13 Sep	+0.82	+0.56	+0.26	+0.04	–0.18
7 Sep	7 – 13 Sep	+0.76	+0.56	+0.20	–0.02	–0.23
13 Sep	13 – 19 Sep	+0.94	+0.69	+0.25	–0.01	–0.16

Every error is inside its 80% interval. In all four the dipole forecast was too positive.

Limitations

- **The eastern correction** has made the forecast worse in all four verified windows, by 0.14 to 0.29 °C. Read the eastern box and the dipole as upper estimates.
- **The gain over persistence** depends on which persistence is used, and at week 1 the forecast is less accurate than 7-day persistence in both boxes.
- **The baseline** is 2006–2025 rather than 1991–2020, and on a different SST analysis from BoM's, so these anomalies are not comparable with published dipole values.
- **Twenty years** is a short training record. The western forecast lies 2.6 to 2.9 standard deviations beyond the training mean, so the regression is extrapolating.
- **Separate correction** of the two boxes before differencing changes the dipole's amplitude in a direction that varies by window. This run it is amplified at weeks 1 and 2 and damped at weeks 3 and 4.
- **The intervals** come from past errors, not the ensemble spread, and none is attached to the alternative methods.
- **Box and map numbers** differ by up to 0.19 °C in this run. Quote the table.
- **One model.** All figures come from the ECMWF ensemble.