

The Glaciolacustrine Sediment Record of Cariboo Lake, BC: Implications for Holocene Fluvial and Glacial Watershed Dynamics

Alex Cebulski & Dr. Joseph Desloges




UNIVERSITY OF
TORONTO

Background


- Alpine environments & climate change
- Changes Include:
 - Volume & area of glacier extent
 - The production, connectivity, and delivery of sediment
- Environmental proxies in glaciated watersheds
 - Ice cores
 - Dating of material in glacier forefields
 - Glaciolacustrine sediment cores

Background

- Regional climate comparison of various glaciolacustrine sediment study sites



(Doughty et al., 1997)



(Hodder et al., 2006)



Purpose: To determine if the distal glacier-fed sediment record from Cariboo Lake provides a proxy of past geomorphic and hydroclimatic change

Objectives:

1. Determine the mechanisms that control the production, connection, and transport of sediment
2. Analyze the sediment record of Cariboo Lake and determine if it reflects changes in past watershed activity
3. Compare the Cariboo Lake sediment record to other regional climate proxies



Study Site

Cariboo Lake, British Columbia





Study Site

Cariboo Lake, British Columbia



Study Site

Cariboo Lake, British Columbia




Methods

- Sub-Bottom Acoustic Survey



Methods

- Sub-Bottom Acoustic Survey
- CTD water column characteristics



Methods

- Sub-Bottom Acoustic Survey
- CTD water column characteristics
- 20 Ekman surficial cores (~10 cm length)
 - Laminae thickness measurements
 - Organic content
 - Sediment grain diameter
 - Provide a spatial record



Methods

- Sub-Bottom Acoustic Survey
- CTD water column characteristics
- 20 Ekman dredge cores (~ 10 cm length)
 - Laminae thickness measurements
 - Organic content
 - Sediment grain diameter
 - Provide a spatial record
- 4 Vibrocores (~ 4 m length)
 - Laminae thickness measurements
 - Organic content
 - Sediment grain diameter
 - AMS radiocarbon analysis
 - Provide a temporal record





V2 – (96 – 166 cm)


Results: Sub-Bottom Acoustics Transects




Sub-Bottom Acoustics: Transect C




Sediment Thickness Map




Ekman Surficial Sediment Record




Ekman Cores




Ekman Grain Size Diameter



Vibro Core Location Map



Long Core Bottoms



V1



AD 450 - 525

V2



AD 150 - 200

V3



N. D.

V4



N. D.

Distance From Cariboo River Delta (km)


3.75

5.50


7.55

10.75


Regional Comparison




Regional Comparison




Regional Comparison



Regional Comparison



Sediment Yield Comparison



Cariboo Lake: ◆

6.35-7.59 Mg·km⁻²·a⁻¹

2.4% Perm. Snow Cover

Thank you for your time!



Acknowledgements

- Funding provided by the University of Toronto.
- Lab assistance by Anna Soleski, Selina Amaral, Mike Gorton, George Kretschmann, and Yanan Liu at the University of Toronto.



UNIVERSITY OF
TORONTO

- Field support provided by Michael Allchin, Laszlo Enyedy, and Caitlin Langford at University of Northern British Columbia via the Quesnel River Research Centre.

