

DRP: Fourier Analysis on Groups

Adithya Ganesh

October 12, 2019

1 Introduction to Finite Markov Chains

A finite Markov chain is a process which moves among the elements of a finite set Ω in the following manner: when at $x \in \Omega$, the next position is chosen according to a fixed probability distribution $P(x, \cdot)$.

(More formal definition.)

Theorem.

Definition. For $x \in \Omega$, define the **hitting time** for x to be

$$\tau_x := \min \{t \geq 0, X_t = x\},$$

that is, the first time at which the chain visits state x .

2 Meeting 1

s_0, \dots, s_n family of random variables. \mathcal{F}_i is the filtration at time i , which is equal to $\sigma(s_0, s_1, \dots, s_i)$, the signal algebra generated by s_0, \dots, s_i .

A probability space is given of three things:

$$(\Omega, \mathcal{F}, p),$$

where Ω is the state space, \mathcal{F} is a sigma-algebra, and p is a prob measures

When \mathcal{F} is the power set in the discrete case. The σ algebra is a collection C of subsets such that

- $\emptyset \in C$.
- $A \in C \implies A^C \in C$.
- $A_1, A_2, \dots, A_n \in C \implies \bigcup A_i \in C$.

A probability measure is a function $p : \mathcal{F} \rightarrow [0, 1]$ such that

- $p(\emptyset) = 0$

- $p(\bigcup_{i=1}^{\infty} A_i) = \sum_{i=1}^{\infty} P(A_i)$ if $A_i \cap A_j = \emptyset$.

Why sigma algebras? Consider a sequence of n coin tosses. We can write

$$(\Omega, \mathcal{F}, p) = \left(\{0, 1\}^n, 2^{\{0, 1\}^n}, p = \text{uniform} \right).$$

In the continuous case, (Ω, \mathcal{F}, p) , $\Omega = [0, 1]$, where $p = \text{uniform measure}$, $p((a, b)) = b - a$.

Definition. Martingales.

Definition. If \mathcal{F} is a σ -algebra, X is an RV, then

$$\mathbb{E}[X | \mathcal{F}]$$

is the unique rv s.t.

$$\mathbb{E}[X 1_A] = \mathbb{E}[\mathbb{E}[X | \mathcal{F}] 1_A],$$

for all $A \in \mathcal{F}$.

Let $T = \text{stopping time}$, which is the property that

$$\{T \leq n\} \in \mathcal{F}_n = \sigma(S_0, \dots, S_n).$$

The optimal stopping theorem states that

$$\mathbb{E}[S_T] = 0,$$

if $S_{\min(N, T)}$ is bounded.

The Gambler's problem is a martingale.

$$\mathbb{E}[S_T] = 0,$$

so

$$P(S_T = -b)(-b) + P(S_T = a)a = 0,$$

so $p = \frac{b}{a+b}$.

Exercise. Gambler ruin with bias. Suppose $S_n = X_1 + X_2 + \dots + X_n$, $\mathbb{E}[X_i] = p < 0$.