

SGD with Variance Reduction beyond Empirical Risk Minimization

MCM 2017

Massil Achab

S. Gaïffas, E. Bacry, A. Guilloux

4th July 2017

CMAP, Ecole Polytechnique

1. Introduction
2. SGD and Variance Reduction
3. Beyond Empirical Risk Minimization
4. HSVRG: Hybrid SVRG
5. Theoretical Guarantees
6. Numerical Experiments

Introduction

- Most machine learning problems can be expressed as a convex optimization problem

$$\hat{\theta} = \arg \min_{\theta \in \mathbb{R}^d} \left(\frac{1}{n} \sum_{i=1}^n f_i(\theta) + g(\theta) \right),$$

- Usually, $f = \frac{1}{n} \sum_{i=1}^n f_i$ is a convex data fitting term (usually smooth), and g is a convex penalty on the predictor (smooth or not).
- Example (Lasso): $f_i(\theta) = (y_i - \theta^\top x_i)^2$ and $g(\theta) = \|\theta\|_1$.

Usual supervised machine learning framework

- **Data:** n observations $(x_i, y_i) \in \mathcal{X} \times \mathcal{Y}, i = 1, \dots, n$
- **Prediction function:** $h(x, \theta) \in \mathbb{R}$ parametrized by $\theta \in \mathbb{R}^d$
- **Empirical Risk Minimization:** find $\hat{\theta}$ solution of

$$\min_{\theta \in \mathbb{R}^d} \frac{1}{n} \sum_{i=1}^n f_i(\theta) + g(\theta), \quad \text{with} \quad f_i(\theta) = \ell(y_i, h(x_i, \theta))$$

- **Examples:** linear regression, logistic regression, support vector machines, neural networks, ...

Cox partial likelihood

- Goal: relate covariates of a patient to its survival time
- Cox regression model: *regression* that can extract information from patients whose failure time is not observed
- Semi-parametric model on the hazard function of a patient

$$\lim_{h \rightarrow 0} \frac{\mathbb{P}(t \leq T \leq t + h | t \leq T)}{h} = \lambda_0(t) \exp(\theta^\top x)$$

- Estimation of θ through maximization of the partial log-likelihood

$$\ell(\theta) = -\frac{1}{|D|} \sum_{i \in D} \left[-\theta^\top x_i + \log \left(\sum_{j \in R_i} \exp(\theta^\top x_j) \right) \right]$$

Vanilla algorithm to find $\hat{\theta}$

Proximal operator

The proximal operator of h is defined by

$$\text{prox}_h(y) = \arg \min_{x \in \mathbb{R}^d} \{h(x) + 1/2 \|y - x\|_2^2\},$$

where $\|\cdot\|_2$ is the usual Euclidean norm.

Proximal Gradient Descent

- Given a starting point θ_0 and η small enough
- Until convergence, do

$$\theta^{t+1} \leftarrow \text{prox}_{\eta g} [\theta^t - \eta \nabla f(\theta^t)]$$

SGD and Variance Reduction

- Large-scale and high-dimensional machine learning: both \mathbf{d} , dimension of each observation, and \mathbf{n} , number of observations, are large
- Consequence: computation of $\nabla f(\theta^t) = \frac{1}{n} \sum_{i=1}^n \nabla f_i(\theta^t)$ is time-consuming.
- Idea behind **Stochastic Gradient Descent**: replace $\nabla f(\theta^t)$ with a *descent direction* d^t , faster to compute

$$d^t = \nabla f(\theta^t) + \epsilon^t \quad \text{with} \quad \mathbb{E}[\epsilon^t] = 0.$$

- The usual version of SGD from Robbins and Monro (1951) writes

$$\begin{aligned}i_t &\sim \mathcal{U}[n] \\d^t &= \nabla f(\theta^t) + \left(\nabla f_{i_t}(\theta^t) - \frac{1}{n} \sum_{j=1}^n \nabla f_j(\theta^t) \right) \\&= \nabla f_{i_t}(\theta^t)\end{aligned}$$

- **ERM** framework with linear prediction $h(x, \theta) = \theta^\top \Phi(x)$,

$$\nabla f_i(\theta) = \partial_2 \ell(y_i, \theta^\top \Phi(x_i)) \Phi(x_i),$$

then computing d^t is **n times faster** than computing $\nabla f(\theta)$.

SGD's high variance




Figure 1: Picture borrowed from Francis Bach's presentations.

SGD's high variance

Assumptions

We assume f is L -smooth i.e.

$$\forall x, y : \|\nabla f(x) - \nabla f(y)\|_2 \leq L\|x - y\|_2,$$

and f μ -strongly convex i.e.

$$\forall x, y : f(y) \geq f(x) + \nabla f(x)^\top (y - x) + \frac{\mu}{2}\|y - x\|_2^2$$

Convergence rates

$$\begin{aligned}\mathbb{E}(f(\theta^t) - f(\theta^*)) &= O(1/t) \text{ for Stochastic Gradient Descent} \\ &= O(\rho^t) \text{ with } \rho < 1 \text{ for Gradient Descent}\end{aligned}$$

The latter rate is called *linear convergence rate*.

Variance Reduction approach

Variance Reduction approach

- We want to compute $\mathbb{E}[X]$, and we can easily compute $\mathbb{E}[Y]$, where Y is highly correlated to X .

- We design the estimator $Z_\alpha = \alpha(X - Y) + \mathbb{E}[Y]$. Then,

$$\text{Var}(Z_\alpha) = \alpha^2[\text{Var}(X) + \text{Var}(Y) - 2\text{Cov}(X, Y)].$$

- When $\text{Cov}(X, Y)$ is high enough, $\text{Var}(Z_\alpha) \leq \text{Var}(X)$, giving the method its name.
- The standard approach uses $\alpha = 1$, leading to an unbiased estimate $\mathbb{E}[Z_\alpha] = \mathbb{E}[X]$.

SGD with Variance Reduction

- Surprisingly enough, recent findings (M. Schmidt, N. Le Roux & F. Bach, 2012), (R. Johnson & T. Zhang, 2013), (A. Defazio, F. Bach & S. Lacoste-Julien, 2014) proved that reducing the variance in SGD enables reaching a **linear convergence rate**.
- Descent directions of these algorithms

$$\text{(SAG)} \quad \theta \leftarrow \theta - \eta \left(\frac{\nabla f_i(\theta) - y_i}{n} + \frac{1}{n} \sum_{j=1}^n y_j \right),$$

$$\text{(SAGA)} \quad \theta \leftarrow \theta - \eta \left(\nabla f_i(\theta) - y_i + \frac{1}{n} \sum_{j=1}^n y_j \right),$$

$$\text{(SVRG)} \quad \theta \leftarrow \theta - \eta \left(\nabla f_i(\theta) - \nabla f_i(\tilde{\theta}) + \frac{1}{n} \sum_{j=1}^n \nabla f_j(\tilde{\theta}) \right).$$

- SAG's descent direction is biased ($\alpha = 1/n$), while SAGA's and SVRG's are unbiased ($\alpha = 1$)

Beyond Empirical Risk Minimization

Remarks

- These methods work well for problems where computing $\nabla f_i(\theta)$ is **n times faster** than computing $\nabla f(\theta)$.
- True for Generalized Linear Models since $\nabla f_i(\theta)$ is colinear to x_i .
- In more complex problems, computing $\nabla f_i(\theta)$ can be long as computing $\nabla f(\theta)$.

How to adapt the previous algorithms to this new case ?

- The negative Cox partial log-likelihood takes the form

$$-\ell(\theta) = \frac{1}{|D|} \sum_{i \in D} \left[-\theta^\top x_i + \log \left(\sum_{j \in R_i} \exp(\theta^\top x_j) \right) \right]$$

- Likelihood and gradient

$$f_i(\theta) = -\theta^\top x_i + \log \left(\sum_{j \in R_i} \exp(\theta^\top x_j) \right)$$

$$\nabla f_i(\theta) = -x_i + \sum_{j \in R_i} \pi_\theta^i(j) x_j, \quad \text{with} \quad \pi_\theta^i(j) = \frac{\exp(\theta^\top x_j)}{\sum_{k \in R_i} \exp(\theta^\top x_k)}$$

Gradient of a subfunction as expectation

- Each subfunction's gradient ∇f_i can be expressed as the expectation of a random variable.
- Computing the exact expectation is expensive due to the summation over all possible configurations $k \in R_i$.
- **Our approach:** consider $\nabla f_i(\theta)$ the expectation of a random variable, and approximate it using MCMC:

$$\text{replace } \nabla f_i(\theta) = \mathbb{E}[G_i(\theta)] \quad \text{with} \quad \hat{\nabla} f_i(\theta) = \hat{G}_i(\theta)$$

HSVRG: Hybrid SVRG

Algorithm 1 Hybrid SVRG

```
1: for  $k = 1$  to  $K$  do
2:   for  $t = 0$  to  $m - 1$  do
3:     Pick  $i \sim \mathcal{U}[n]$ 
4:      $\hat{\nabla} f_i(\theta^t) \leftarrow \text{APPROXMCMC}(\theta^t, i, N_k)$ .
5:      $d^t = \hat{\nabla} f_i(\theta^t) - \nabla f_i(\tilde{\theta}) + \frac{1}{n} \sum_{j=1}^n \nabla f_j(\tilde{\theta})$ 
6:      $\omega^{t+1} \leftarrow \theta^t - \gamma d^t$ 
7:      $\theta^{t+1} \leftarrow \text{prox}_{\gamma g}(\omega^{t+1})$ 
8:   end for
9:   Update  $\tilde{\theta} \leftarrow \frac{1}{m} \sum_{t=1}^m \theta^t$ ,  $\theta^0 \leftarrow \tilde{\theta}$ 
10:  Compute  $\nabla f_i(\tilde{\theta})$  for  $i = 1, \dots, n$ 
11: end for
```

$\text{APPROXMCMC}(\theta^t, i, N_k)$ outputs an approximation of $\nabla f_i(\theta^t)$ using N_k iterations of a MCMC. We focused on two implementations:

- Independent Metropolis-Hastings¹ (IMH)
- Adaptive Importance Sampling (AIS)

¹with uniform proposal

Theoretical Guarantees

Assumption

We assume that the bias and the expected squared error of the Monte Carlo error $\eta = \widehat{G}_i(\theta) - \mathbb{E}[G_i(\theta)]$ can be bounded this way

$$\|\mathbb{E}_t[\eta]\| \leq \frac{C_1}{N_k} \text{ and } \mathbb{E}_t[\|\eta\|^2] \leq \frac{C_2}{N_k},$$

where N_k is the length of the Markov chain.

Proposition

Suppose that there exists $M > 0$ such that the proposal Q and the stationary distribution π satisfy $\pi(x) \leq MQ(x)$, for all x in the support of π . Then, the error η^t obtained by Algorithm IMH satisfies the previous assumption.

Remark: We can compute C_1 and C_2 from special cases (for Cox model, for instance).

Theorem

Suppose that $F = f + g$ is μ -strongly convex. Consider Algorithm **HSVRG**, with a phase length m and a step-size $\gamma \in (0, \frac{1}{16L})$ satisfying

$$\rho = \frac{1}{m\gamma\mu(1-8L\gamma)} + \frac{8L\gamma(1+1/m)}{1-8L\gamma} < 1. \quad (1)$$

Assuming there exists $B > 0$ such that $\sup_{t \geq 0} \|\theta^t - \theta^*\|_2 \leq B$, we have:

$$\mathbb{E}[F(\tilde{\theta}^K)] - F(\theta^*) \leq C\rho^K + D \sum_{k=1}^K \rho^{K-k} \frac{1}{N_k}, \quad (2)$$

where $C = F(\theta^0) - F(\theta^*)$, and $D = \frac{3\gamma C_2 + BC_1}{1-8L\gamma}$.

Corollary

In the previous theorem, the choice $N_k = k^\alpha \rho^{-k}$ with $\alpha > 1$ gives

$$\mathbb{E}[F(\tilde{\theta}^K)] - F(\theta^*) \leq D' \rho^K,$$


where $D' = F(\theta^0) - F(\theta^*) + D \sum_{k \geq 1} k^{-\alpha}$.

This entails that **HSVRG** achieves a **linear rate** under strong convexity.

Numerical Experiments

Experiments

We ran experiments on the Cox model with IMH with (uniform and adaptative proposal) and AIS (adaptative proposal).



- CRFs model the conditional probability of a structured output $y \in \mathcal{Y}$ (such as a sequence of labels) given an input $x \in \mathcal{X}$ (such as a sequence of words) based on features $F(x, y)$ and parameter θ using

$$\mathbb{P}(y|x, \theta) = \frac{\exp(\theta^\top F(x, y))}{\sum_{y'} \exp(\theta^\top F(x, y'))}.$$

- Likelihood and gradient,

$$\begin{aligned} f_i(\theta) &= -\log \mathbb{P}(y_i | x_i, \theta) \\ \nabla f_i(\theta) &= -F(x_i, y_i) + \sum_{y' \in \mathcal{Y}} \mathbb{P}(y' | x_i, \theta) F(x_i, y') \end{aligned}$$

Questions?

Adaptative Importance Sampling

- IMH with uniform proposal outputs an estimate with high variance.
- Use Normalized Importance Sampling in APPROXMCMC.

$$I = \mathbb{E}_p[f(X)] = \mathbb{E}_q \left[f(X) \frac{p(X)}{q(X)} \right]$$

$$\hat{I}_n = \frac{1}{n} \sum_{k=1}^n f(X^{(k)}) \frac{p(X^{(k)})}{q(X^{(k)})}, \text{ with } X^{(k)} \sim q$$

$$\hat{J}_n = \sum_{k=1}^n f(X^{(k)}) \frac{p(X^{(k)})}{q(X^{(k)})} / \sum_{k=1}^n \frac{p(X^{(k)})}{q(X^{(k)})}, \text{ with } X^{(k)} \sim q$$

- Use $\pi_{\tilde{\theta}}$ as adaptative proposal, where $\tilde{\theta}$ is updated every phase.

- Apply this new APPROXMCMC(θ, i, N) to CRF outputs

$$\hat{J}_n = -F(x_i, y_i) + \sum_{k=1}^N \frac{\exp((\theta - \tilde{\theta})^\top F(x_i, y^{(k)}))}{\sum_{j=1}^N \exp((\theta - \tilde{\theta})^\top F(x_i, y^{(j)}))} F(x_i, y^{(k)})$$

- The sequence $(y^{(k)})$ is sampled from $\mathbb{P}(\bullet | x_i, \tilde{\theta})$.
- We remind the true subgradient is

$$\nabla f_i(\theta) = -F(x_i, y_i) + \sum_{y \in \mathcal{Y}} \frac{\exp((\theta - \tilde{\theta})^\top F(x_i, y))}{\sum_{y'} \exp((\theta - \tilde{\theta})^\top F(x_i, y'))} F(x_i, y)$$