| Variable | Confirmed case N = 15 | Not confirmed N = 19 | p-valu |
|--|--------------------------|-------------------------|--------|
| HIV status, n (%) | | | 0.2 |
| Person with HIV | 0 (0%) | 1 (5.3%) | |
| Person without HIV | 13 (87%) | 18 (95%) | |
| Unknown | 2 (13%) | 0 (0%) | |
| ever hospitalized, n (%) | | | 0.4 |
| Yes | 14 (93%) | 15 (79%) | |
| No | 1 (6.7%) | 4 (21%) | |
| Unknown | 0 (0%) | 0 (0%) | |
| Length of hospitalization, Median (Q1, Q3) | 6 (5, 9) | 6 (4, 10) | 0.3 |
| Missing/Unknown | 1 | 4 | |
| Escalation of care, n (%) | | | >0.9 |
| Yes | 0 (0%) | 1 (6.7%) | |
| No | 14 (100%) | 14 (93%) | |
| Unknown | 0 (0%) | 0 (0%) | |
| Missing/Unknown | 1 | 4 | |
| supplemental oxygen, n (%) | | | >0.9 |
| Yes | 0 (0%) | 1 (6.7%) | |
| No | 14 (100%) | 14 (93%) | |
| Unknown | 0 (0%) | 0 (0%) | |
| Missing/Unknown | 1 | 4 | |
| receive nasogastric fluids, n (%) | | | >0.9 |
| Yes | 0 (0%) | 0 (0%) | |
| No | 15 (100%) | 19 (100%) | |
| Unknown | 0 (0%) | 0 (0%) | |
| receive IV fluids during course monkeypox infection, n (%) | | | >0.9 |
| Yes | 0 (0%) | 1 (5.3%) | |
| No | 15 (100%) | 18 (95%) | |
| Unknown | 0 (0%) | 0 (0%) | |
| receive analgesia for pain management, n (%) | | | >0.9 |
| Yes | 4 (27%) | 5 (26%) | |
| No | 11 (73%) | 14 (74%) | |
| Unknown | 0 (0%) | 0 (0%) | |
| receive antibiotics during course monkeypox infection, n (%) | | | >0.9 |
| Yes | 13 (87%) | 17 (89%) | |
| No | 2 (13%) | 2 (11%) | |
| Unknown | 0 (0%) | 0 (0%) | |
| receive antifungals during course monkeypox infection, n (%) | | | >0.9 |

| /ariable | Confirmed case N = 15 | Not confirmed N = 19 | p-value |
|--|--------------------------|-------------------------|---------|
| Yes | 0 (0%) | 0 (0%) | |
| No | 15 (100%) | 19 (100%) | |
| Unknown | 0 (0%) | 0 (0%) | |
| experimental antiviral during course monkeypox infection, n (% | b) | | >0.9 |
| Yes | 0 (0%) | 0 (0%) | |
| No | 15 (100%) | 19 (100%) | |
| Unknown | 0 (0%) | 0 (0%) | |