| Variable | N = 16 |
|--|-------------|
| Sex at birth, n/N (%) | |
| Male | 5/15 (33%) |
| Female | 10/15 (67%) |
| Missing/Unknown | 1 |
| Age group, n/N (%) | |
| 10-14 years | 1/15 (6.7%) |
| 20-24 years | 4/15 (27%) |
| 25-29 years | 2/15 (13%) |
| 30-34 years | 1/15 (6.7%) |
| 35-39 years | 3/15 (20%) |
| 40-44 years | 2/15 (13%) |
| 50-54 years | 1/15 (6.7%) |
| 65-69 years | 1/15 (6.7%) |
| Missing/Unknown | 1 |
| Race, n/N (%) | |
| African/Black | 15/15 (100% |
| Missing/Unknown | 1 |
| health care worker, n/N (%) | |
| No | 15/15 (100% |
| Missing/Unknown | 1 |
| Laboratory worker, n/N (%) | |
| No | 15/15 (100% |
| Missing/Unknown | 1 |
| Sex worker, n/N (%) | 7/15 (47%) |
| Missing/Unknown | 1 |
| Nature of contact, n/N (%) | |
| Direct contact with sexual fluids | 1/1 (100%) |
| Missing/Unknown | 15 |
| Most recent CD4, Median (Min, Max) | 98 (1, 98) |
| Missing/Unknown | 1 |
| Antiretroviral (ARV) treatments, n/N (%) | |
| Yes | 14/14 (100% |
| Missing/Unknown | 2 |
| Viral load, n/N (%) | |
| Undetectable | 3/3 (100%) |
| Missing/Unknown | 13 |
| Most recent viral load, n/N (%) | 0/0 (NA%) |
| Missing/Unknown | 16 |
| AIDS - defining illness, n/N (%) | |

| Variable | N = 16 |
|---|--------------|
| No | 15/15 (100%) |
| Missing/Unknown | 1 |
| New diagnosis of HIV, n/N (%) | |
| No | 13/13 (100%) |
| Missing/Unknown | 3 |
| Current new STI (Sexually Transmitted Infection) diagnosis, n/N (%) | |
| No | 3/3 (100%) |
| Missing/Unknown | 13 |
| Syphilis, n/N (%) | |
| Yes | 4/4 (100%) |
| Missing/Unknown | 12 |
| N. gonorrhoeae, n/N (%) | |
| No | 3/3 (100%) |
| Missing/Unknown | 13 |
| Chlamydia trachomatis, n/N (%) | |
| No | 3/3 (100%) |
| Missing/Unknown | 13 |
| Immunosuppressive condition, n/N (%) | 1/16 (6.3%) |
| Length of hospitalization, Median (Min, Max) | 12 (5, 26) |
| Missing/Unknown | 5 |