Lab₀

Indiana University | M303 Linear Algebra | Fall 2025

Anthony Chao

format rat
format compact

1.) Store the matrix $\begin{bmatrix} 2 & 5 & 7 & 1 \\ 3 & 1 & 1 & 0 \\ -1 & -3 & 7 & 2 \end{bmatrix}$ in MATLAB and call it M. Do not suppress the output.

M = [2, 5, 7, 1; 3, 1, 1, 0; -1, -3, 7, 2]

2.) Store the matrix $\begin{bmatrix} 1 & 3 \\ -1 & 1 \\ 0 & 0 \\ 2 & 4 \end{bmatrix}$ and call it *N*. Suppress the output.

N = [1, 3; -1, 1; 0, 0; 2, 4];

3.) Store the matrix $\begin{bmatrix} 3 & \frac{1}{2} \\ -\frac{1}{2} & \frac{5}{3} \end{bmatrix}$ and call it F. Do not suppress the output.

F = [3, 1/2; -1/2, 5/3]

F = 3 1/2 5/3

4.) Store the matrix $\begin{bmatrix} 2 & 3 & 4 & 5 & 6 & 1 \\ 3 & 4 & 5 & 6 & 1 & 2 \\ 4 & 5 & 6 & 1 & 2 & 3 \\ 5 & 6 & 1 & 2 & 3 & 4 \\ 6 & 1 & 2 & 3 & 4 & 5 \end{bmatrix}$ using **the multiline method** and call it *G*. Do not suppress the output.

1

G = [1, 2, 3, 4, 5, 6; 2, 3, 4, 5, 6, 1; 3, 4, 5, 6, 1, 2; 4, 5, 6, 1, 2, 3; 5, 6, 1, 2, 3, 4;

6, 1, 2, 3, 4, 5]

G =	
-----	--

1	2	3	4	5	6
2	3	4	5		
3	4	5	6	1	2
4	5	6	1	2	3
5	6	1	2	3	4
6	1	2	3	4	5