

The sign test

Packages

```
library(tidyverse)
library(smmr)
```

`smmr` is new. See later how to install it.

Duality between confidence intervals and hypothesis tests

- ▶ Tests and CIs really do the same thing, if you look at them the right way. They are both telling you something about a parameter, and they use same things about data.
- ▶ To illustrate, some data (two groups):

```
my_url <- "http://ritsokiguess.site/datafiles/duality.txt"  
twogroups <- read_delim(my_url," ")
```

The data

```
twogroups
```

```
# A tibble: 15 x 2
```

| | y | group |
|----|-------|-------|
| | <dbl> | <dbl> |
| 1 | 10 | 1 |
| 2 | 11 | 1 |
| 3 | 11 | 1 |
| 4 | 13 | 1 |
| 5 | 13 | 1 |
| 6 | 14 | 1 |
| 7 | 14 | 1 |
| 8 | 15 | 1 |
| 9 | 16 | 1 |
| 10 | 13 | 2 |
| 11 | 13 | 2 |
| 12 | 14 | 2 |
| 13 | 17 | 2 |
| 14 | 18 | 2 |

95% CI (default)

for difference in means, group 1 minus group 2:

```
t.test(y ~ group, data = twogroups)
```

Welch Two Sample t-test

data: y by group

t = -2.0937, df = 8.7104, p-value = 0.0668

alternative hypothesis: true difference in means between gr

95 percent confidence interval:

-5.5625675 0.2292342

sample estimates:

mean in group 1 mean in group 2

13.00000

15.66667

90% CI

```
t.test(y ~ group, data = twogroups, conf.level = 0.90)
```

Welch Two Sample t-test

data: y by group

t = -2.0937, df = 8.7104, p-value = 0.0668

alternative hypothesis: true difference in means between groups

90 percent confidence interval:

-5.010308 -0.323025

sample estimates:

mean in group 1 mean in group 2

13.00000

15.66667

Too high Hypothesis test

Null is that difference in means is zero:

```
t.test(y ~ group, mu=0, data = twogroups)
```

Welch Two Sample t-test

data: y by group

t = -2.0937, df = 8.7104, p-value = 0.0668

alternative hypothesis: true difference in means between gr

95 percent confidence interval:

-5.5625675 0.2292342

sample estimates:

mean in group 1 mean in group 2

13.00000

15.66667

Comparing results

Recall null here is $H_0 : \mu_1 - \mu_2 = 0$. P-value 0.0668.

- ▶ 95% CI from -5.6 to 0.2 , contains 0 .
- ▶ 90% CI from -5.0 to -0.3 , does not contain 0 .
- ▶ At $\alpha = 0.05$, would not reject H_0 since P-value > 0.05 .
- ▶ At $\alpha = 0.10$, *would* reject H_0 since P-value < 0.10 .

Test and CI

Not just coincidence. Let $C = 100(1 - \alpha)$, so $C\%$ gives corresponding CI to level- α test. Then following always true. (Symbol \iff means “if and only if”.)

| Test decision | | Confidence interval |
|---------------------------------------|--------|---------------------------------------|
| Reject H_0 at level α | \iff | $C\%$ CI does not contain H_0 value |
| Do not reject H_0 at level α | \iff | $C\%$ CI contains H_0 value |

Idea: “Plausible” parameter value inside CI, not rejected;
“Implausible” parameter value outside CI, rejected.

The value of this

- ▶ If you have a test procedure but no corresponding CI:
- ▶ you make a CI by including all the parameter values that would not be rejected by your test.
- ▶ Use:
 - ▶ $\alpha = 0.01$ for a 99% CI,
 - ▶ $\alpha = 0.05$ for a 95% CI,
 - ▶ $\alpha = 0.10$ for a 90% CI, and so on.

Testing for non-normal data

- ▶ The IRS (“Internal Revenue Service”) is the US authority that deals with taxes (like Revenue Canada).
- ▶ One of their forms is supposed to take no more than 160 minutes to complete. A citizen’s organization claims that it takes people longer than that on average.
- ▶ Sample of 30 people; time to complete form recorded.
- ▶ Read in data, and do t -test of $H_0 : \mu = 160$ vs. $H_a : \mu > 160$.
- ▶ For reading in, there is only one column, so can pretend it is delimited by anything.

Read in data

```
my_url <- "http://ritsokiguess.site/datafiles/irs.txt"
irs <- read_csv(my_url)
irs
```

```
# A tibble: 30 x 1
```

```
  Time
```

```
<dbl>
```

| | |
|----|-----|
| 1 | 91 |
| 2 | 64 |
| 3 | 243 |
| 4 | 167 |
| 5 | 123 |
| 6 | 65 |
| 7 | 71 |
| 8 | 204 |
| 9 | 110 |
| 10 | 178 |

```
# i 20 more rows
```

Test whether mean is 160 or greater

```
with(irs, t.test(Time, mu = 160,  
                  alternative = "greater"))
```


One Sample t-test

```
data: Time  
t = 1.8244, df = 29, p-value = 0.03921  
alternative hypothesis: true mean is greater than 160  
95 percent confidence interval:  
 162.8305      Inf  
sample estimates:  
mean of x  
 201.2333
```

Reject null; mean (for all people to complete form) greater than 160.

But, look at a graph

```
ggplot(irs, aes(x = Time)) + geom_histogram(bins = 6)
```



Comments

- ▶ Skewed to right.
- ▶ Should look at *median*, not mean.

The sign test

- ▶ But how to test whether the median is greater than 160?
- ▶ Idea: if the median really is 160 (H_0 true), the sampled values from the population are equally likely to be above or below 160.
- ▶ If the population median is greater than 160, there will be a lot of sample values greater than 160, not so many less. Idea: test statistic is number of sample values greater than hypothesized median.

Getting a P-value for sign test 1/3

- ▶ How to decide whether “unusually many” sample values are greater than 160? Need a sampling distribution.
- ▶ If H_0 true, pop. median is 160, then each sample value independently equally likely to be above or below 160.
- ▶ So number of observed values above 160 has binomial distribution with $n = 30$ (number of data values) and $p = 0.5$ (160 is hypothesized to be *median*).

Getting P-value for sign test 2/3

- ▶ Count values above/below 160:

```
irs %>% count(Time > 160)
```

```
# A tibble: 2 x 2
  `Time > 160`      n
  <lgl>          <int>
1 FALSE           13
2 TRUE            17
```

- ▶ 17 above, 13 below. How unusual is that? Need a *binomial table*.

Getting P-value for sign test 3/3

- ▶ R function `dbinom` gives the probability of eg. exactly 17 successes in a binomial with $n = 30$ and $p = 0.5$:

```
dbinom(17, 30, 0.5)
```

```
[1] 0.1115351
```

- ▶ but we want probability of 17 *or more*, so get all of those, find probability of each, and add them up:

```
tibble(x=17:30) %>%  
  mutate(prob=dbinom(x, 30, 0.5)) %>%  
  summarize(total=sum(prob))
```

```
# A tibble: 1 x 1  
  total  
  <dbl>  
1 0.292
```

Using my package smmr

- ▶ I wrote a package `smmr` to do the sign test (and some other things). Installation is a bit fiddly:
 - ▶ Install `devtools` (once) with

```
install.packages("devtools")
```

- ▶ then install `smmr` using `devtools` (once):

```
library(devtools)  
install_github("nxskok/smmr")
```

- ▶ Then load it:

```
library(smmr)
```

smmr for sign test

- ▶ smmr's function `sign_test` needs three inputs: a data frame, a column and a null median:

```
sign_test(irs, Time, 160)
```

```
$above_below
```

```
below above
```

```
13      17
```

```
$p_values
```

```
alternative    p_value
```

```
1         lower 0.8192027
```

```
2         upper 0.2923324
```

```
3    two-sided 0.5846647
```

Comments (1/3)

- ▶ Testing whether population median *greater than* 160, so want *upper-tail* P-value 0.2923. Same as before.
- ▶ Also get table of values above and below; this too as we got.

Comments (2/3)

- P-values are:

| Test | P-value |
|------|---------|
| t | 0.0392 |
| Sign | 0.2923 |

- These are very different: we reject a mean of 160 (in favour of the mean being bigger), but clearly *fail* to reject a median of 160 in favour of a bigger one.
- Why is that? Obtain mean and median:

```
irs %>% summarize(mean_time = mean(Time),  
                  median_time = median(Time))
```

```
# A tibble: 1 x 2  
  mean_time median_time  
    <dbl>      <dbl>  
1    201.        172.
```

Comments (3/3)

- ▶ The mean is pulled a long way up by the right skew, and is a fair bit bigger than 160.
- ▶ The median is quite close to 160.
- ▶ We ought to be trusting the sign test and not the t-test here (median and not mean), and therefore there is no evidence that the “typical” time to complete the form is longer than 160 minutes.
- ▶ Having said that, there are clearly some people who take a lot longer than 160 minutes to complete the form, and the IRS could focus on simplifying its form for these people.
- ▶ In this example, looking at any kind of average is not really helpful; a better question might be “do an unacceptably large fraction of people take longer than (say) 300 minutes to complete the form?”: that is, thinking about worst-case rather than average-case.

Confidence interval for the median

- ▶ The sign test does not naturally come with a confidence interval for the median.
- ▶ So we use the “duality” between test and confidence interval to say: the (95%) confidence interval for the median contains exactly those values of the null median that would not be rejected by the two-sided sign test (at $\alpha = 0.05$).

For our data

- ▶ The procedure is to try some values for the null median and see which ones are inside and which outside our CI.
- ▶ `smmr` has `pval_sign` that gets just the 2-sided P-value:

```
pval_sign(160, irs, Time)
```

```
[1] 0.5846647
```

- ▶ Try a couple of null medians:

```
pval_sign(200, irs, Time)
```

```
[1] 0.3615946
```

```
pval_sign(300, irs, Time)
```

```
[1] 0.001430906
```

- ▶ So 200 inside the 95% CI and 300 outside.

Doing a whole bunch

- Choose our null medians first:

```
(d <- tibble(null_median=seq(100,300,20)))
```

```
# A tibble: 11 x 1
```

```
  null_median
```

```
    <dbl>
```

| | |
|----|-----|
| 1 | 100 |
| 2 | 120 |
| 3 | 140 |
| 4 | 160 |
| 5 | 180 |
| 6 | 200 |
| 7 | 220 |
| 8 | 240 |
| 9 | 260 |
| 10 | 280 |
| 11 | 300 |

... and then

“for each null median, run the function `pval_sign` for that null median and get the P-value”:

```
d %>% rowwise() %>%  
  mutate(p_value = pval_sign(null_median, irs, Time))
```

A tibble: 11 x 2

Rowwise:

| | null_median | p_value |
|---|-------------|----------|
| | <dbl> | <dbl> |
| 1 | 100 | 0.000325 |
| 2 | 120 | 0.0987 |
| 3 | 140 | 0.200 |
| 4 | 160 | 0.585 |
| 5 | 180 | 0.856 |
| 6 | 200 | 0.362 |
| 7 | 220 | 0.0428 |
| 8 | 240 | 0.0161 |
| 9 | 260 | 0.00522 |

Make it easier for ourselves

```
d %>% rowwise() %>%  
  mutate(p_value = pval_sign(null_median, irs, Time)) %>%  
  mutate(in_out = ifelse(p_value > 0.05, "inside", "outside"))
```

A tibble: 11 x 3

Rowwise:

| | null_median | p_value | in_out |
|----|-------------|----------|---------|
| | <dbl> | <dbl> | <chr> |
| 1 | 100 | 0.000325 | outside |
| 2 | 120 | 0.0987 | inside |
| 3 | 140 | 0.200 | inside |
| 4 | 160 | 0.585 | inside |
| 5 | 180 | 0.856 | inside |
| 6 | 200 | 0.362 | inside |
| 7 | 220 | 0.0428 | outside |
| 8 | 240 | 0.0161 | outside |
| 9 | 260 | 0.00522 | outside |
| 10 | 280 | 0.00143 | outside |
| 11 | 300 | 0.00143 | outside |

confidence interval for median?

- ▶ 95% CI to this accuracy from 120 to 200.
- ▶ Can get it more accurately by looking more closely in intervals from 100 to 120, and from 200 to 220.

A more efficient way: bisection

- ▶ Know that top end of CI between 200 and 220:

```
lo <- 200  
hi <- 220
```

- ▶ Try the value halfway between: is it inside or outside?

```
try <- (lo + hi) / 2  
try
```

```
[1] 210
```

```
pval_sign(try,irs,Time)
```

```
[1] 0.09873715
```

- ▶ Inside, so upper end is between 210 and 220. Repeat (over):

... bisection continued

```
lo <- try  
try <- (lo + hi) / 2  
try
```

```
[1] 215
```

```
pval_sign(try, irs, Time)
```

```
[1] 0.06142835
```

- ▶ 215 is inside too, so upper end between 215 and 220.
- ▶ Continue until have as accurate a result as you want.

Bisection automatically

- A loop, but not a for since we don't know how many times we're going around. Keep going while a condition is true:

```
lo = 200
hi = 220
while (hi - lo > 1) {
    try = (hi + lo) / 2
    ptry = pval_sign(try, irs, Time)
    print(c(try, ptry))
    if (ptry <= 0.05)
        hi = try
    else
        lo = try
}
```

The output from this loop

```
[1] 210.00000000    0.09873715
[1] 215.00000000    0.06142835
[1] 217.50000000    0.04277395
[1] 216.25000000    0.04277395
[1] 215.62500000    0.04277395
```

- 215 inside, 215.625 outside. Upper end of interval to this accuracy is 215.

Using smmr

- ▶ smmr has function `ci_median` that does this (by default 95% CI):

```
ci_median(irs, Time)
```

```
[1] 119.0065 214.9955
```

- ▶ Uses a more accurate bisection than we did.
- ▶ Or get, say, 90% CI for median:

```
ci_median(irs, Time, conf.level=0.90)
```

```
[1] 123.0031 208.9960
```

- ▶ 90% CI is shorter, as it should be.


Bootstrap

- ▶ but, was the sample size (30) big enough to overcome the skewness?
- ▶ Bootstrap, again:

```
tibble(sim = 1:1000) %>%  
  rowwise() %>%  
  mutate(my_sample = list(sample(irs$Time, replace = TRUE)))  
  mutate(my_mean = mean(my_sample)) %>%  
  ggplot(aes(x=my_mean)) + geom_histogram(bins=10) -> g
```

The sampling distribution

g



Comments

- ▶ A little skewed to right, but not nearly as much as I was expecting.
- ▶ The t -test for the mean might actually be OK for these data, *if the mean is what you want*.
- ▶ In actual data, mean and median very different; we chose to make inference about the median.
- ▶ Thus for us it was right to use the sign test.