

## Numerical Summaries

# Summarizing data in R 1/2

- ▶ Have seen `summary` (5-number summary of each column). But what if we want:
  - ▶ a summary or two of just one column
  - ▶ a count of observations in each category of a categorical variable
  - ▶ summaries by group
  - ▶ a different summary of all columns (eg. SD)
- ▶ To do this, meet pipe operator `%>%`. This takes input data frame, does something to it, and outputs result. (Learn: `Ctrl-Shift-M`.)

## Summarizing data in R 2/2

- ▶ Output from a pipe can be used as input to something else, so can have a sequence of pipes.
- ▶ Summaries include: `mean`, `median`, `min`, `max`, `sd`, `IQR`, `quantile` (for obtaining quartiles or any percentile), `n` (for counting observations).
- ▶ Use our Australian athletes data again.

## Packages for this section

```
library(tidyverse)
```

## Summarizing one column

► Mean height:

```
athletes %>% summarize(m=mean(Ht))
```

```
# A tibble: 1 x 1
```

```
      m
```

```
  <dbl>
```

```
1  180.
```

or to get mean and SD of BMI:

```
athletes %>% summarize(m=mean(BMI),s=sd(BMI))
```

```
# A tibble: 1 x 2
```

```
      m
```

```
      s
```

```
  <dbl> <dbl>
```

```
1  23.0  2.86
```

# Quartiles

- ▶ quantile calculates percentiles (“fractiles”), so we want the 25th and 75th percentiles:

```
athletes %>% summarize( Q1=quantile(Wt, 0.25),  
                        Q3=quantile(Wt, 0.75))
```

```
# A tibble: 1 x 2
```

```
      Q1      Q3
```

```
  <dbl> <dbl>
```

```
1  66.5  84.1
```

## Creating new columns

- ▶ These weights are in kilograms. Maybe we want to summarize the weights in pounds.
- ▶ Convert kg to lb by multiplying by 2.2.
- ▶ Create new column and summarize that:

```
athletes %>% mutate(wt_lb=Wt*2.2) %>%  
  summarize(Q1_lb=quantile(wt_lb, 0.25),  
            Q3_lb=quantile(wt_lb, 0.75))
```

```
# A tibble: 1 x 2  
  Q1_lb Q3_lb  
  <dbl> <dbl>  
1  146.  185.
```

## Counting how many

for example, number of athletes in each sport:

```
athletes %>% count(Sport)
```

```
# A tibble: 10 x 2
```

	Sport	n
	<chr>	<int>
1	BBall	25
2	Field	19
3	Gym	4
4	Netball	23
5	Row	37
6	Swim	22
7	T400m	29
8	TSprnt	15
9	Tennis	11
10	WPolo	17



## Counting how many, variation 2:

Another way (which will make sense in a moment):

```
athletes %>% group_by(Sport) %>%  
  summarize(count=n())
```

```
# A tibble: 10 x 2
```

	Sport	count
	<chr>	<int>
1	BBall	25
2	Field	19
3	Gym	4
4	Netball	23
5	Row	37
6	Swim	22
7	T400m	29
8	TSprnt	15
9	Tennis	11
10	WPolo	17

## Summaries by group

- ▶ Might want separate summaries for each “group”, eg. mean and SD of height for males and females. Strategy is `group_by` (to define the groups) and then `summarize`:

```
athletes %>% group_by(Sex) %>%  
  summarize(mean_Ht = mean(Ht), sd_Ht = sd(Ht))
```

```
# A tibble: 2 x 3  
  Sex      mean_Ht sd_Ht  
  <chr>    <dbl> <dbl>  
1 female    175.  8.24  
2 male     186.  7.90
```

## Count plus stats

- ▶ If you want number of observations per group plus some stats, you need to go the `n()` way:

```
athletes %>% group_by(Sex) %>%  
summarize(n = n(), mean_Ht = mean(Ht), sd_Ht = sd(Ht))
```

```
# A tibble: 2 x 4  
  Sex      n mean_Ht sd_Ht  
  <chr> <int>   <dbl> <dbl>  
1 female   100    175.   8.24  
2 male    102    186.   7.90
```

- ▶ This explains second variation on counting within group:  
“within each sport/Sex, how many athletes were there?”

## Summarizing several columns

- ▶ Standard deviation of each (numeric) column:

```
athletes %>% summarize(across(where(is.numeric), \ (x) sd(x))
```

```
# A tibble: 1 x 11
```

	RCC	WCC	Hc	Hg	Ferr	BMI	SSF	`%Bfat`	LBM	
	<dbl>	<dbl>	<dbl>	<dbl>	<dbl>	<dbl>	<dbl>	<dbl>	<dbl>	<dbl>
1	0.458	1.80	3.66	1.36	47.5	2.86	32.6	6.19	13.1	

- ▶ Median and IQR of all columns whose name starts with H:

```
athletes %>% summarize(across(starts_with("H"),  
                               list(med = \ (x) median(x),  
                                    iqr = \ (x) IQR(x))))
```

```
# A tibble: 1 x 6
```

	Hc_med	Hc_iqr	Hg_med	Hg_iqr	Ht_med	Ht_iqr
	<dbl>	<dbl>	<dbl>	<dbl>	<dbl>	<dbl>
1	43.5	4.98	14.7	2.07	180.	12.2

## Same thing by group

```
athletes %>%  
  group_by(Sex) %>%  
  summarize(across(starts_with("H"),  
                    list(
                      med = \ (h) median(h),  
                      iqr = \ (h) IQR(h))))
```

```
# A tibble: 2 x 7
```

	Sex	Hc_med	Hc_iqr	Hg_med	Hg_iqr	Ht_med	Ht_iqr
	<chr>	<dbl>	<dbl>	<dbl>	<dbl>	<dbl>	<dbl>
1	female	40.6	4.03	13.5	1.60	175	8.68
2	male	45.5	2.57	15.5	0.975	186.	11.3