



**Universidade do Estado do Rio de Janeiro**

centro

unidade patrono

Nome Sobrenome

**Título do trabalho acadêmico**

Cidade

aaaa

Nome Sobrenome

**Título do trabalho acadêmico**

Dissertação apresentada, como requisito parcial para obtenção do título de Mestre, ao Programa de Pós-Graduação em Curso, da Universidade do Estado do Rio de Janeiro.



Orientador: Cargo Titulação Nome Sobrenome  
Coorientador: Cargo Titulação Nome Sobrenome

Cidade

aaaa

CATALOGAÇÃO NA FONTE  
UERJ / REDE SIRIUS / BIBLIOTECA CTC/D

---

D979

Sobrenome, Nome

Título do trabalho acadêmico / Nome Sobrenome. – Cidade, aaaa-  
33 f.

Orientador: Cargo Titulação Nome Sobrenome

Dissertação (Mestrado) – Universidade do Estado do Rio de Janeiro,  
unidade, Programa de Pós-Graduação em Curso, aaaa.

1. primeira palavra chave.. 2. segunda palavra chave.. 3. terceira  
palavra chave.. I. Cargo Titulação Nome Sobrenome. II. Universidade do  
Estado do Rio de Janeiro. III. unidade. IV. Título

CDU 02:141:005.7

---

Autorizo, apenas para fins acadêmicos e científicos, a reprodução total ou parcial desta  
dissertação, desde que citada a fonte.

---

Assinatura

---

Data

Nome Sobrenome

**Título do trabalho acadêmico**

Dissertação apresentada, como requisito parcial para obtenção do título de Mestre, ao Programa de Pós-Graduação em Curso, da Universidade do Estado do Rio de Janeiro.

Aprovada em dd de Mês de aaaa.

Banca Examinadora:

---

Cargo Titulação Nome Sobrenome (Orientador)  
Unidade – Instituição

---

Cargo Titulação Nome Sobrenome (Coorientador)  
Unidade – Instituição

---

primeiro membro titular da banca  
instituição

---

segundo membro titular da banca  
instituição

---

terceiro membro titular da banca  
instituição

---

primeiro membro suplente da banca  
instituição

---

segundo membro suplente da banca  
instituição

---

terceiro membro suplente da banca  
instituição instituição instituição instituição

Cidade

aaaa

## DEDICATÓRIA

Texto da dedicatória

## AGRADECIMENTOS

Texto de agradecimento



## RESUMO

SOBRENOME, Iniciais. Do. Nome. *Título do trabalho acadêmico*. aaaa. 33 f.  
Dissertação (Mestrado em Curso) – unidade, Universidade do Estado do Rio de Janeiro,  
Cidade, aaaa.

Texto do resumo em português.

Palavras-chave: primeira palavra chave. segunda palavra chave. terceira palavra chave.



## ABSTRACT

SOBRENOME, Iniciais. Do. Nome. *Title of dissertation.* aaaa. 33 f. Dissertação (Mestrado em Curso) – unidade, Universidade do Estado do Rio de Janeiro, Cidade, aaaa.

Abstract in English.

Keywords: first keyword. second keyword. third keyword.

## LIST OF FIGURES

Figure 1 - Título da figura. . . . .	23
Figure 2 - Título da figura. . . . .	24

## LIST OF TABLES

Table	1 - Título da tabela. . . . .	23
-------	-------------------------------	----

## LISTA DE ALGORITMOS

Algoritmo 1 - Título do algoritmo. . . . .	25
--	----

## LISTA DE ABREVIATURAS E SIGLAS

sigla1	por extenso
sigla2	por extenso
sigla3	por extenso

## LISTA DE SÍMBOLOS

<i>simbolo1</i>	significado e/ou valor
<i>simbolo2</i>	significado e/ou valor
<i>simbolo3</i>	significado e/ou valor

## CONTENTS

	<b>INTRODUÇÃO</b>	14
1	<b>BOLTZMANN MACHINES</b>	15
1.1	<b>Boltzmann Machines</b>	15
2	<b>TÍTULO DO CAPÍTULO 1</b>	22
3	<b>TÍTULO DO CAPÍTULO 2</b>	25
	<b>CONCLUSÃO</b>	27
	<b>BIBLIOGRAPHY</b>	28
	<b>GLOSSÁRIO</b>	29
	<b>APPENDIX A – Primeiro apêndice</b>	30
	<b>APPENDIX B – Segundo apêndice</b>	31
	<b>ANNEX A – Primeiro anexo</b>	32
	<b>ANNEX B – Segundo anexo</b>	33

## INTRODUÇÃO

Texto da introdução. Texto, texto texto (AMADO, 1969), texto Amado (1969).  
Texto Andrade em 1997, texto ANDRADE, texto.



## 1 BOLTZMANN MACHINES

Here we begin by explaining the theory behind Boltzmann Machines

EXPLICAR OS ELEMENTOS DE PROBABILIDADE COM OS QUAIS ESTAMOS LIDANDO: QUEM SÃO AS VARIÁVEIS ALEATÓRIAS E COMO VAMOS DOMINÁ-LAS NO TEXTO, COMENTAR QUE ESTAMOS TRATANDO COM VARIÁVEIS DISCRETAS, COMO VAMOS IDENTIFICAR AS PROBABILIDADES,...

EXPLICAR O QUE OS ÍNDICES REPRESENTAM, e as variáveis.

EXPLICAR OS TERMOS  $\omega$  e  $\phi$ .  $\omega$  É para os pesos, e  $\phi$  para o bias de cada unidade.

EXPLICAR MINHA NOTAÇÃO sobre o sobre-escrito entre parênteses!!!

### 1.1 Boltzmann Machines

Boltzmann Machines (BM) are a type of stochastic neural networks (SNN) where the connections between units, which are described by  $\omega$ , are symmetrical, i.e.,  $\omega_{ij} = \omega_{ji}$  [HERTZ]. This kind of stochastic neural networks are capable of learning internal representation and to model an input distribution. Boltzmann Machines were named after the Boltzmann distribution. Due to its stochastic behaviour, the probability of the state of the system to be found in a certain configuration is given by previous mentioned distribution [HERTZ]. According to [MONTUFAR, 2018], BM can be seen as an extension of Hopfield networks to include hidden units.

Boltzmann Machines have visible and hidden units. The visible units are linked to the external world and they correspond to the components of an observation. On the other hand, the hidden units do not have any connection outside of the network and model the dependencies between the components of the observations [FISCHER, 2012]. In BM, there is no connection restriction, this means that every unit, visible or hidden, can be connected to every other unit as in a complete graph, this pattern is not mandatory as some of the connections may not exist depending on the network layout.

Training Boltzmann Machines means finding the right connection between the units.

Boltzmann Machines (BM) are stochastic neural networks with symmetric connections, i.e.,  $\omega_{ij} = \omega_{ji}$ . Boltzmann Machines use the Boltzmann distribution to determine the probability of the state of the system of the network. BM resembles the Hopfield networks with the inclusion of hidden units. Finding the right connections between the hidden units without knowing it from the training patterns what the hidden units represent is part of solving the Boltzmann Machine problem.

Units  $x_i$  in BM are split into two kinds: visible and hidden units. The visible units have connection to the outside world and are the units that receive the data input. On the other hand, the hidden units do not have any connection to the outside of the network and they are responsible to find the data relation from the input. In a BM, the connections between units can be complete or not. Regardless of how the connections are, every connection in a BM is symmetric.

BM are made of stochastic units  $x_i$  which each of them can assume a binary value with a certain probability as follows:

$$x_i = \begin{cases} 1 & \text{with probability } g(h_i) \\ 0 & \text{with probability } 1 - g(h_i) \end{cases}, \quad (1)$$

where

$$h_i = \sum_j \omega_{ij} x_j, \quad (2)$$

and

$$g(h_i) = \frac{1}{1 + e^{-2\beta h_i}}. \quad (3)$$

Due to the symmetrical connections, there is an energy function given by

$$H(\mathbf{x}) = - \sum_i \sum_j \omega_{ij} x_i x_j - \sum_i \phi_i x_i, \quad (4)$$

where  $\mathbf{x} = (x_1, x_2, \dots, x_n)$ , and  $n$  is equal to the number of units in the network, and the above energy function has minimum when there is a stable state characterised by

$$x_i = \text{sgn}(h_i). \quad (5)$$

The probability  $P$  of finding the system in a given state  $\mathbf{x}$  after the equilibrium is reached can be computed as follows:

$$P(\mathbf{x}) = \frac{1}{Z} e^{-\beta H(\mathbf{x})}, \quad (6)$$

where

$$Z = \sum_{\mathbf{x}'} e^{-\beta H(\mathbf{x}')} \quad (7)$$

is the partition function.

The learning process of a Boltzmann Machine consists in adjusting the connections

$\omega_{ij}$  in such a way that the state of the visible units have a particular desired probability distribution.

Let us identify the state of the visible units by an index  $v$  and the state of the hidden units by an index  $h$ . Considering a system which has  $N$  visible units and  $K$  hidden units, the whole system have  $2^{N+K}$  possibilities of states in which it can be found.

The joint probability  $P_{vh}$  is the probability of finding the visible and hidden units in the states  $v$  and  $h$ , respectively. This probability measurement is given by the Boltzmann distribution:

$$P_{vh} = \frac{e^{-\beta H_{vh}}}{Z}, \quad (8)$$

where

$$Z = \sum_u \sum_k e^{-\beta H_{uk}}, \quad (9)$$

and

$$H_{vh} = - \sum_i \sum_j \omega_{ij} x_i^{(vh)} x_j^{(vh)} - \sum_i \phi_i x_i^{(vh)}. \quad (10)$$

As mentioned above, the problem a Boltzmann Machine is trying to solve is determining the connections  $\omega_{ij}$  between units such that the visible units have a certain probability distribution. In order to do that, we need to find the marginal probability of the state  $v$  in which the visible units are found regardless of the state  $h$  of the hidden units. The marginal probability  $P_v$  is given by

$$P_v = \sum_h P_{vh} = \sum_h \frac{e^{-\beta H_{vh}}}{Z}. \quad (11)$$

Although we know that  $P_v$  is a function of the connections  $\omega_{ij}$ , and that this is the probability of finding the visible units in the state  $v$ . We want the states to have a certain probability  $Q_v$ , i.e., a desired probability. This means that ideally we would like to match the empirical distribution of the data, even though we do not have access to the correct distribution, only to what the observed data has given us as an input to training the model.

One way to evaluate the difference between two probability distribution, for example,  $P_v$  and  $Q_v$ , is using the Kullback-Leibler divergence, which can also be referred to relative entropy,  $E$ , which will be our cost function. (EXPLICAR FUNÇÃO DE CUSTO

e  $D_{KL}!!!$ ).

$$E = \sum_v Q_v \ln \left( \frac{Q_v}{P_v} \right). \quad (12)$$

The relative entropy  $E$  has the property of always being equal or greater than zero. It reaches zero only if  $P_v = Q_v$ , which means that we are able to retrieve the exactly probability distribution of the input data at the visible units.

$$\begin{aligned} E &= \sum_v Q_v \ln \left( \frac{Q_v}{P_v} \right) \\ &\geq \sum_v Q_v \left( 1 - \frac{P_v}{Q_v} \right) \\ &= \sum_v (Q_v - P_v) \\ &= \sum_v Q_v - \sum_v P_v = 1 - 1 \\ &\Rightarrow E \geq 0. \end{aligned} \quad (13)$$

From the gradient descent equation

$$\Delta\omega_{ij} = -\eta \frac{\partial E}{\partial \omega_{ij}}, \quad (14)$$

where

$$\begin{aligned} E &= \sum_v Q_v \ln \left( \frac{Q_v}{P_v} \right) \\ &= \sum_v [\ln(Q_v) - \ln(P_v)]. \end{aligned} \quad (15)$$

In the following steps, we present the gradient descent derivation

$$\begin{aligned} \Delta\omega_{ij} &= -\eta \frac{\partial E}{\partial \omega_{ij}} \\ &= -\eta \frac{\partial}{\partial \omega_{ij}} \left[ \sum_v Q_v (\ln(Q_v) - \ln(P_v)) \right] \\ &= \eta \frac{\partial}{\partial \omega_{ij}} \left[ \sum_v Q_v \ln(P_v) \right] \\ &= \eta \sum_v Q_v \frac{\partial}{\partial \omega_{ij}} [\ln(P_v)] \\ &\Rightarrow \Delta\omega_{ij} = \eta \sum_v \frac{Q_v}{P_v} \frac{\partial P_v}{\partial \omega_{ij}}. \end{aligned} \quad (16)$$

To continue with the computation of  $\Delta\omega_{ij}$ , we have to find the derivative of  $\partial P_v/\partial\omega_{ij}$ , from the marginal probability, equation 11,

$$P_v = \frac{\sum_h e^{-\beta H_{vh}}}{\sum_u \sum_k e^{-\beta H_{uk}}}, \quad (17)$$

thus the derivative of  $P_v$  follows

$$\begin{aligned} \frac{\partial P_v}{\partial\omega_{ij}} &= \frac{\partial}{\partial\omega_{ij}} \left[ \frac{\sum_h e^{-\beta H_{vh}}}{\sum_u \sum_k e^{-\beta H_{uk}}} \right] \\ &= \frac{1}{\sum_u \sum_k e^{-\beta H_{uk}}} \sum_h (-\beta) e^{-\beta H_{vh}} \frac{\partial H_{vh}}{\partial\omega_{ij}} \\ &\quad - \sum_h e^{-\beta H_{vh}} \frac{1}{(\sum_u \sum_k e^{-\beta H_{uk}})^2} \sum_u \sum_k e^{-\beta H_{uk}} (-\beta) \frac{\partial H_{uk}}{\partial\omega_{ij}}. \end{aligned} \quad (18)$$

Following the equation 18, we need to compute the term  $\partial H_{vh}/\partial\omega_{ij}$ ,

$$\begin{aligned} \frac{\partial H_{vh}}{\partial\omega_{ij}} &= \frac{\partial}{\partial\omega_{ij}} \left[ -\frac{1}{2} \sum_m \sum_n \omega_{mn} x_m^{(vh)} x_n^{(vh)} - \sum_m \phi_{mm} x_m^{(vh)} \right] \\ &= \frac{\partial}{\partial\omega_{ij}} \left[ -\frac{1}{2} \sum_{m \neq i,j} \sum_{n \neq i,j} \omega_{mn} x_m^{(vh)} x_n^{(vh)} - \frac{1}{2} \omega_{ij} x_i^{(vh)} x_j^{(vh)} - \frac{1}{2} \omega_{ji} x_j^{(vh)} x_i^{(vh)} - \sum_m \phi_m x_m^{(vh)} \right], \end{aligned} \quad (19)$$

as the connections between units are symmetric, i.e.,  $\omega_{ij} = \omega_{ji}$ , then we can simplify equation 19,

$$\begin{aligned} \frac{\partial H_{vh}}{\partial\omega_{ij}} &= \frac{\partial}{\partial\omega_{ij}} \left[ -\frac{1}{2} \sum_{m \neq i,j} \sum_{n \neq i,j} \omega_{mn} x_m^{(vh)} x_n^{(vh)} - \omega_{ij} x_i^{(vh)} x_j^{(vh)} - \sum_m \phi_m x_m^{(vh)} \right] \\ &= \frac{\partial}{\partial\omega_{ij}} \left[ -\frac{1}{2} \sum_{m \neq i,j} \sum_{n \neq i,j} \omega_{mn} x_m^{(vh)} x_n^{(vh)} \right] + \frac{\partial}{\partial\omega_{ij}} \left[ -\omega_{ij} x_i^{(vh)} x_j^{(vh)} \right] + \frac{\partial}{\partial\omega_{ij}} \left[ -\sum_m \phi_m x_m^{(vh)} \right] \\ &\Rightarrow \frac{\partial H_{vh}}{\partial\omega_{ij}} = -x_i^{(vh)} x_j^{(vh)}. \end{aligned} \quad (20)$$

Analogous to  $\partial H_{vh}/\partial\omega_{ij}$ , we have  $H_{uk}$  derivative, which is

$$\frac{\partial H_{uk}}{\partial\omega_{ij}} = -x_i^{(uk)} x_j^{(uk)}. \quad (21)$$

Going back to equation 18, we can replace the derivatives of  $H$ , and solve the

derivative of the marginal probability  $P_v$ ,

$$\begin{aligned}
\frac{\partial P_v}{\partial \omega_{ij}} &= \frac{1}{Z} \sum_h e^{-\beta H_{vh}} (-\beta) (-x_i^{(vh)} x_j^{(vh)}) - \frac{1}{Z^2} \sum_h e^{-\beta H_{vh}} \sum_u \sum_k e^{-\beta H_{uk}} (-\beta) (-x_i^{(uk)} x_j^{(uk)}) \\
&= \beta \left[ \sum_h \frac{e^{-\beta H_{vh}}}{Z} x_i^{(vh)} x_j^{(vh)} - \sum_h \frac{e^{-\beta H_{vh}}}{Z} \sum_u \sum_k \frac{e^{-\beta H_{uk}}}{Z} x_i^{(uk)} x_j^{(uk)} \right] \\
&= \beta \left[ \sum_h P_{vh} x_i^{(vh)} x_j^{(vh)} - P_v \sum_u \sum_k P_{uk} x_i^{(uk)} x_j^{(uk)} \right] \\
&\Rightarrow \frac{\partial P_v}{\partial \omega_{ij}} = \beta \left[ \sum_h P_{vh} x_i^{(vh)} x_j^{(vh)} - P_v \langle x_i x_j \rangle \right].
\end{aligned} \tag{22}$$

Given the derivative of  $P_v$  in relation to  $\omega_{ij}$ , we can compute the learning term  $\Delta \omega_{ij}$ , from equation 16,

$$\begin{aligned}
\Delta \omega_{ij} &= \eta \sum_v \frac{Q_v}{P_v} \beta \left[ \sum_h P_{vh} x_i^{(vh)} x_j^{(vh)} - P_v \langle x_i x_j \rangle \right] \\
&= \eta \beta \left[ \sum_v \sum_h \frac{Q_v}{P_v} P_{vh} x_i^{(vh)} x_j^{(vh)} - \sum_v \frac{Q_v}{P_v} P_v \langle x_i x_j \rangle \right] \\
&= \eta \beta \left[ \sum_v \sum_h Q_v \frac{P_{vh}}{P_v} x_i^{(vh)} x_j^{(vh)} - \sum_v Q_v \langle x_i x_j \rangle \right] \\
&= \eta \beta \left[ \sum_v \sum_h Q_v P_{h|v} x_i^{(vh)} x_j^{(vh)} - \langle x_i x_j \rangle \right].
\end{aligned} \tag{23}$$

In the above derivation, we have used the following relations,

$$P_{h|v} = \frac{P_{vh}}{P_v}, \tag{24}$$

which is the conditional probability equation. In our scenario this equation means that the probability distribution of the hidden units in state  $h$  given the state  $v$  of the visible units is the joint probability distribution of both states,  $v$  and  $h$ , divided by the marginal probability distribution of the visible units in state  $v$ .

The second term in equation 23, is the average of units  $i$  and  $j$  over all combinations of states  $v$  and  $h$  of the system. In other words, we would have to compute all possible combination of states of visible and hidden units,  $v$  and  $h$ , and then average over the specific units  $i$  and  $j$ .

The first term can be simplified by

$$\sum_v Q_v \sum_h P_{h|v} x_i^{(vh)} x_j^{(vh)} = \sum_v Q_v \langle x_i x_j \rangle^{(v)} = \langle \langle x_i x_j \rangle^{(v)} \rangle. \quad (25)$$

Then equation 23 becomes

$$\Delta\omega_{ij} = \eta\beta \left[ \langle \langle x_i x_j \rangle^{(v)} \rangle_{clamped} - \langle x_i x_j \rangle_{free} \right], \quad (26)$$

it is important to notice that the subscripts *clamped* means that we have to fix a certain  $v$  state on the visible units otherwise the second term in the equation does not have a reference. On the other hand, the subscript *free* identify the ...

## 2 TÍTULO DO CAPÍTULO 1

Texto do capítulo. Texto, texto, Figura 1. Texto Figura 2(a).



Figure 1 - Título da figura.



Legend: Texto da legenda.

Source: Citação da fonte ou 'O autor.'.

Table 1 - Título da tabela.

X	Y
1,20	15,7
1,23	15,6
1,19	15,3
1,26	15,1
1,22	15,5
1,16	15,3
1,37	15,7

Legend: Texto da legenda.

Source: Citação da fonte ou 'O autor.'.

Figure 2 - Título da figura.



(a)



(b)



(c)

Legend: Texto da legenda. (a) Texto da imagem. (b) Texto da imagem. (c) Texto da imagem.

Source: Citação da fonte ou 'O autor'.

### 3 TÍTULO DO CAPÍTULO 2

Texto do capítulo. Texto, texto Algoritmo 1. Texto.

Algoritmo 1 - Título do algoritmo.

#### DOCUMENTAÇÃO

##### TÍTULO

**Nome do algoritmo**

##### PROPÓSITO

Propósito do algoritmo.

##### MÉTODO

Método utilizado no algoritmo.

##### ENTRADAS

a, m: multiplicador e módulo

n0: semente

i: contador auxiliar

##### SAÍDAS

n: número aleatório

##### OBSERVAÇÕES, RESTRIÇÕES, REQUISITOS

Observações, restrições e requisitos.

##### ALGORITMO IDENTIFICAÇÃO

**declarar**  $a, m, i$  numéricos

**declarar**  $n0, n$  numéricos

```

1.   $m \leftarrow 13$ 
2.   $n0 \leftarrow 1$ 
3.  para  $a$  de 2 até  $m - 1$  , fazer      {para cada possível valor de 'a'}
4.  |   escrever "a = ",  $a$ , ":",  $n = \{$ 
5.  |    $n \leftarrow n0$       {reinicia a geração com a semente  $n0$ }
6.  |   para  $i$  de 0 até  $m - 1$  , fazer
7.  |   |    $n \leftarrow \text{resto}(a * n, m)$       {gerador de números aleatórios}
8.  |   |   se  $(n == n0)$ , então      {se fim da sequencia...}
9.  |   |   |   escrever  $n, \{$ 
10. |   |   |   parar
11. |   |   senão
12. |   |   |   escrever  $n$ 
13. |   |   fim se
14. |   fim para
15. fim para
— continua —

```

Algoritmo 1 - Título do algoritmo. (continuação)

```
— continuação —  
a ← 1  
enquanto (a < 10), fazer      {comentário}  
|   escrever a  
|   a ← a + 1  
fim enquanto  
a ← 1  
repetir      {comentário}  
|   escrever a  
|   a ← a + 1  
até que (a ≥ 10)  
  
a ← 1  
fazer      {comentário}  
|   escrever a  
|   a ← a + 1  
enquanto (a < 10)  
FIM ALGORITMO  
FIM DOCUMENTAÇÃO
```

## CONCLUSÃO

Texto da conclusão.

**BIBLIOGRAPHY**

AMADO, J. *Dona Flor e seus dois maridos*: historia moral e de amor. 9. ed. Rio de Janeiro: Record, 1969. 535 p.

ANDRADE, M. M. d. *Introdução à metodologia científica*. São Paulo: Atlas, 1997. 151 p.

## GLOSSÁRIO

termo	significado
termo	significado
termo	significado

## APPENDIX A – Primeiro apêndice

### A.1 Primeira seção

Texto da primeira seção.

#### A.1.1 Primeira subseção

Texto da primeira subseção.

##### A.1.1.1 Primeira subsubseção

Texto da primeira subsubseção.



## APPENDIX B – Segundo apêndice

### B.1 Primeira seção

Texto da primeira seção.

#### B.1.1 Primeira subseção

Texto da primeira subseção.

##### B.1.1.1 Primeira subsubseção

Texto da primeira subsubseção.

## ANNEX A – Primeiro anexo

### A.1 Primeira seção

Texto da primeira seção.

#### A.1.1 Primeira subseção

Texto da primeira subseção.

##### A.1.1.1 Primeira subsubseção

Texto da primeira subsubseção.

## ANNEX B – Segundo anexo

### B.1 Primeira seção

Texto da primeira seção.

#### B.1.1 Primeira subseção

Texto da primeira subseção.

##### B.1.1.1 Primeira subsubseção

Texto da primeira subsubseção.