

Wouse Liver - All Regions



- Promoter (<=1kb) (36.43%)
 Promoter (1-2kb) (4.39%)
 Promoter (2-3kb) (3.49%)
 5' UTR (0.26%)
 3' UTR (1.9%)

- 1st Exon (1.64%)
 Other Exon (3.01%)
 1st Intron (10.09%)
 Other Intron (16.55%)
 Downstream (<=300) (0.16%)
 Distal Intergenic (22.08%)


Widuse Liver - Conserved in Human Liver



- Promoter (<=1kb) (54.52%)
 Promoter (1-2kb) (3.32%)
 Promoter (2-3kb) (2.46%)
 5' UTR (0.38%)
 3' UTR (1.2%)

- 1st Exon (1.45%)
 Other Exon (2.84%)
 1st Intron (7.61%)
 Other Intron (11.62%)
 Downstream (<=300) (0.09%)
 Distal Intergenic (14.5%)


Mouse Liver - Conserved in numan Pancreas



- □ Promoter (<=1kb) (52.9%)■ Promoter (1-2kb) (3.39%)
- □ Promoter (2–3kb) (3%)
 □ 5' UTR (0.49%)
 □ 3' UTR (1.34%)

- 1st Exon (1.34%)
 Other Exon (2.71%)
 1st Intron (7.83%)
 Other Intron (12.3%)
 Downstream (<=300) (0.13%)
 Distal Intergenic (14.58%)

Mouse Liver - Not Conserved in numan



- □ Promoter (<=1kb) (44.88%)■ Promoter (1-2kb) (4.09%)
- □ Promoter (2–3kb) (2.92%)
 □ 5' UTR (0.29%)
 □ 3' UTR (2.1%)

- 1st Exon (1.22%)
 Other Exon (3.25%)
 1st Intron (8.32%)
 Other Intron (15.28%)
 Downstream (<=300) (0.13%)
 Distal Intergenic (17.54%)