

numan Liver - An Regions



- □ Promoter (<=1kb) (33.64%)
 □ Promoter (1-2kb) (4.06%)
 □ Promoter (2-3kb) (3.24%)
 □ 5' UTR (0.3%)
 □ 3' UTR (2.08%)

- 1st Exon (1.2%)
 Other Exon (2.59%)
 1st Intron (10.75%)
 Other Intron (21.35%)
 Downstream (<=300) (0.09%)
 Distal Intergenic (20.69%)


numan Liver - Conserved in Mouse Liver



- □ Promoter (<=1kb) (48.99%)■ Promoter (1-2kb) (4.17%)
- □ Promoter (2–3kb) (3.1%)
 □ 5' UTR (0.38%)
 □ 3' UTR (1.83%)


- 1st Exon (1.03%)
 Other Exon (1.76%)
 1st Intron (8.22%)
 Other Intron (16.77%)
 Downstream (<=300) (0.04%)
 Distal Intergenic (13.72%)

Human Liver - Conserved in Mouse Pancreas



- □ Promoter (<=1kb) (48.84%)■ Promoter (1-2kb) (3.87%)
- □ Promoter (2–3kb) (2.9%)
 □ 5' UTR (0.81%)
 □ 3' UTR (1.99%)

- 1st Exon (0.91%)
 Other Exon (2.1%)
 1st Intron (7.32%)
 Other Intron (17.7%)
 Downstream (<=300) (0.05%)
 Distal Intergenic (13.5%)



- □ Promoter (<=1kb) (41%)□ Promoter (1-2kb) (4.17%)
- □ Promoter (2–3kb) (3.13%)
 □ 5' UTR (0.33%)
 □ 3' UTR (2.36%)

- 1st Exon (0.81%)
 Other Exon (2.91%)
 1st Intron (9.73%)
 Other Intron (19.44%)
 Downstream (<=300) (0.08%)
 Distal Intergenic (16.05%)