Mountain Pine Beetle Climatic Suitability Maps

Alex M. Chubaty, Barry J. Cooke, Eliot J. B. McIntire October 26, 2016

Contents

Overview	1
Data source	1
Load maps	1
Plot maps	2

Overview

We have 7 MPB climate scenarios (using four different indices: S, L, R, G) ready to use as model drivers. The Logan suitability index (L) is based on summer temperatures. The Regniere suitability index (R) is based on MPB cold tolerance (i.e., winter survival). The Safranyik suitability index (S) is based on aspects of both summer temperatures and winter survival. Finally, the composite SLR index (G) takes the harmonic mean of the S, L, and R models.

description
Safranyik
Logan
Regniere
Composite

For each of the four indices, there are 1981-2010 normals plus the projections from two RCP scenarios (either 4.5 or 8.5), covering a span of 120 years, split into 30-year frames.

All maps are projected using LCC and cover all of Canada.

Data source

Maps were produced in BioSim (need info!).

They were uploaded to and retrieved from the NoFC FTP site on 26 October 2016.


Load maps


```
library(raster)


files <- dir(path = maps.dir, pattern = '[.]tif', full.names = TRUE)


maps <- lapply(files, function(f) {
   raster(f)
})</pre>
```


Plot maps

















```
## NULL
##
## [[7]]
## NULL
## [[8]]
## NULL
##
## [[9]]
## NULL
## [[10]]
## NULL
##
## [[11]]
## NULL
##
## [[12]]
## NULL
##
## [[13]]
## NULL
##
## [[14]]
## NULL
## [[15]]
## NULL
##
## [[16]]
## NULL
##
## [[17]]
## NULL
##
## [[18]]
## NULL
##
## [[19]]
## NULL
##
## [[20]]
## NULL
##
## [[21]]
## NULL
##
## [[22]]
## NULL
##
## [[23]]
## NULL
##
```

[[24]]

NULL

##

[[25]]

NULL

##

[[26]]

NULL

##

[[27]]

NULL

##

[[28]]

NULL