

Date: September 22, 2026  
From: John ZuHone  
To: Chandra Operations Team  
Subject: Chandra Radiation Event and Shutdown in January 2026  
Cc: MSFC Project Science, CXC Director's Office  
File: JAN1926\_memo.md  
Version: 1.0

## 1 Abstract

This memo discusses the thought process that the operations team, especially the ACIS operations team, used during a very high radiation event between January 19-22, 2026. *Chandra* was shut down via an autonomous trigger of SCS-107 via the ACIS txings algorithm. In this case, the txings algorithm was triggered on the “FI Increasing” threshold. The key decision points to resume science operations are reviewed, and the details of the storm are presented.

During this storm, the accumulated attenuated ACE P3 fluence was approximately  $3 \times 10^7$ . Thanks to the autonomous shutdown, ACIS avoided accumulating another  $\sim 4.22 \times 10^9$  of P3 fluence. The agreed annual budget for this quantity is  $2.0 \times 10^{10}$ .

## 2 Introduction

During the days of January 19-22, 2026, *Chandra* experienced a radiation storm which resulted in the autonomous shutdown of the spacecraft via an ACIS txings trigger of SCS-107 to protect the ACIS instrument from the damaging soft solar wind protons that increase CTI. Though we were past solar maximum at the time, solar activity was still high.

The ACIS txings rate onboard monitoring software serves as the only available way to trigger autonomous shutdowns due to high radiation levels. In this storm, the txings rates reached their trigger levels right before the peak of the proton rates in nearly all channels, averting a substantial amount of radiation damage to the ACIS CCDs from soft ( $\sim 100$  keV) proton fluence.

This memo discusses the properties of the storm, the radiation received in terms of the single-orbit and annual budgets, the differences between various radiation measurements, and the response.

All times in this memo are in UTC unless otherwise noted, and may be approximate. The various electron and proton channel fluxes for ACE and GOES are in units of and assumed throughout when numbers for these fluxes are quoted.

## 3 January 18-22 2026 Detailed Timeline

- 2026:018 **Sunday January 18, 2026**
- 2026:018:00:00:00 The JAN1226 load is in progress.
- 2026:018:00:50:01 **Comm** begins (60 min).

- 2026:018:09:00:00 Earth-orbiting satellites detect an X1.9-class solar flare. A full-halo signature CME is associated with it.
- 2026:018:15:50:02 **Comm** begins (60 min).
- 2026:018:20:04:00 GOES proton channels begin a slow rise in flux, the higher energy channels rising later. An hour and a half later, the HRC proxy begins to rise. All increase for the next ~24 hours.
- 2026:019 **Monday January 19, 2026**
- 2026:019:00:52:30 The JAN1926 load begins.
- 2026:019:01:40:01 **Comm** begins (60 min).
- 2026:019:03:35:00 The HRC proxy crosses its nominal trigger threshold.
- 2026:019:05:10:00 The ACE proton channels jump sharply, and continue rising for the next 15 hours. ACE P3 is at ~1000.
- 2026:019:09:52:11 SCS-107 runs due to a txings trigger, out of comm.
- 2026:019:10:20:00 The “strong” (S3) NOAA radiation threshold, with a 10 MeV integrated proton flux of 1000 pfu, is reached.
- 2026:019:11:40:01 **Comm** begins (60 min). The spacecraft comes up with the science loads terminated; ACIS is safe.
- 2026:019:15:45:01 **Comm** begins (60 min).
- 2026:019:16:40:00 ACE and GOES proton channels begin to rise sharply, and will continue to do so for the next ~3 hours.
- 2026:019:19:10:00 The HRC proxy reaches a maximum value of  $\sim 5.8 \times 10^4$ .
- 2026:019:19:24:00 Space weather alerts go out indicating a CME is approaching Earth. Particle rates continue to sharply increase.
- 2026:019:20:05:00 ACE P3 hits a maximum value of  $\sim 5.6 \times 10^5$ .
- 2026:019:21:00:00 ACE and GOES proton channels begin to decline steeply from their peak values.
- 2026:020 **Tuesday January 20, 2026**
- 2026:020:03:30:01 **Comm** begins (60 min).
- 2026:020:14:15:01 **Comm** begins (90 min).
- 2026:020:23:34:38 Time of RADMON disable from the JAN1926 loads.
- 2026:021 **Wednesday January 21, 2026**
- 2026:021:06:50:56 Time of perigee.
- 2026:021:07:10:02 **Comm** begins (60 min).
- 2026:021:11:24:38 Time of RADMON enable from the JAN1926 loads.
- 2026:021:12:10:01 **Comm** begins (60 min).
- 2026:021:14:15:00 First command of the JAN2126 return-to-science loads. ACE P3 has leveled off at a value of ~1000, and will stay at this level for at least the next 24 hours.

## 4 Discussion

In the days leading up to January 18th, 2026, the sunspot region AR 4341 rotated into view. At 09:00, this sunspot produced an X1.9-class solar flare, with which full-halo CME was associated. Early predictions indicated that the CME would arrive by 06:00 UTC on January 20th. Later on in the day, the high-energy GOES proton channels began to rise, as well as the HRC proxy which closely tracks them (see Figures 4 and 5). They would continue to rise for the next ~25 hours. ACIS Ops began to monitor the radiation situation, but no telecon to consider manual actions were planned.

In the early morning of January 19th, the HRC proxy passed its nominal trigger threshold (having no effect on operations as the HRC shield is no longer being continuously operated). At around the same time, the ACE proton and electron channels (see Figures 2 and 3) all sharply rose.

All of the monitored radiation channels continued to rise through the morning of January 19th, and at 2026:019:09:52:11, SCS-107 ran due to a txings trigger, out of comm (see Figure 6), during Obsid 32115. After the dump data was retrieved at the 6:40 am ET comm, it was determined that txings triggered on S2, the lone FI chip turned on for the observation, though given that the BI rates were climbing above their “increasing” triggering threshold, they may have triggered had the FI rates not done so. The storm continued, and shortly after the SCS-107 run the “strong” (S3) NOAA radiation threshold was reached, which [SolarHam.com](http://SolarHam.com) noted had not occurred since October of 2003. At the 6:40 am ET comm, the spacecraft came up with the science loads halted and ACIS safed. Since the CME had not yet arrived and was expected to on January 20th, the tentative plan for a return to science time was the morning of January 21st.

Later on in the afternoon of the 19th, the ACE and GOES channels began to rise very sharply. They continued to rise for the next ~3 hours. Several hours later, alerts went out indicating the early arrival of the CME from the X1.9 flare. Shortly after this, ACE P3 hit a maximum value of  $\sim 5.6 \times 10^5$  (see Figure 1), and all particle rates began to decline. Their decline would flatten early in the morning of January 20th, and either stay relatively flat or slowly declining for a few days afterward. ACE P3 would settle to a value of ~1000.

In the end, the total accumulated attenuated ACE P3 fluence for the orbit was  $3 \times 10^7$ . The total ACE P3 attenuated fluence that would have been accumulated had ACIS not been safed is  $\sim 4.25 \times 10^9$ . The txings trigger prevented the accumulation of this significant amount of fluence.

## 5 Data plots for the January 2026 storm

### 5.1 ACE P3 Flux

In Figure 1, we have plotted the 5-minute averaged ACE P3 flux rate, in the usual units, which are protons  $\text{s}^{-1} \text{cm}^{-2} \text{sr}^{-1} \text{MeV}^{-1}$ , throughout the January 2026 storm. Also marked are radiation belt passages, the time of autonomous SCS-107 execution, and comm times (the same times are marked in the rest of the radiation vs. time plots).

### 5.2 ACE Proton Fluxes

Figure 2 shows the flux from four ACE proton channels (P1, P3, P5, and P7) during the storm. Though only P3 is our proxy for damage to the ACIS CCDs, the other channels can serve as informative diagnostics.




Fig. 1: Figure 1: ACE P3 flux during the January 2026 storm. Purple shaded regions indicate radiation belt passages; blue shaded regions mark scheduled DSN communications.




Fig. 2: Figure 2: Flux from the ACE P1, P3, P5, and P7 proton channels during the January 2026 storm. Shaded regions and vertical lines have the same meaning as in Figure 1.

### 5.3 ACE Electron Fluxes

Figure 3 shows the flux from two ACE electron channels (DE1 and DE4) during the storm. These electron channels more often than not tend to rise and fall rapidly when the higher-energy GOES protons are doing the same.




Fig. 3: Figure 3: Flux from the ACE DE1 and DE4 electron channels during the January 2026 storm. Shaded regions and vertical lines have the same meaning as in Figure 1.

### 5.4 GOES Proton Flux

Figure 4 shows the flux from four GOES proton channels (P1, P3, P5, and P7) during the storm. These higher-energy protons are more representative of the radiation that triggers ACIS txings. GOES P5 and P7 have a steep increase in flux at the time of the txings trigger.

### 5.5 HRC Proxy

Figure 5 shows the HRC Shield Proxy during the storm. The HRC Anti-Coincidence Shield rates are no longer available for radiation monitoring, but had they been, they would have triggered SCS-107 at around the same time as the txings trigger.




Fig. 4: Figure 4: Flux from the GOES P1, P3, P5, and P7 proton channels during the January 2026 storm. Shaded regions and vertical lines have the same meaning as in Figure 1.




Fig. 5: Figure 5: HRC Shield Proxy during the January 2026 storm. Shaded regions and vertical lines have the same meaning as in Figure 1.

## 5.6 txings Rates




Fig. 6: Figure 6: ACIS threshold-crossing (txings) rates during the January 2026 storm. Blue is for FI chips, orange is for BI. The horizontal lines are the increasing-rate trip thresholds for each chip type. Shaded regions and vertical lines have the same meaning as in Figure 1.

## 6 Lessons learned

## 7 Resources

The research required for this memo, as well as the real-time response, would not have been possible without the valiant efforts of <https://www.solarham.com> to provide timely and accurate space weather information.

The archive of ACE data stored in ASCII tables at <https://sohoftp.nascom.nasa.gov/sdb/goes/ace/daily/> has gaps that are not back-filled; the full dataset can however be found in the “ACE Browse” archive:

<ftp://mussel.srl.caltech.edu/pub/ace/browse/>

The data are in HDF4 format, which can be converted to HDF5 data by use of a program `h4toh5` which I downloaded from <https://www.hdfeos.org/software/h4toh5.php>. A Python script, `get_ace.py`, which downloads the data and uses `h4toh5` to convert it is available on the HEAD LAN in `/data/acis/ace`.

Instructions for downloading the data using this script and extracting the ACE proton channels are given in `/data/acis/ace/README_browse.md`.

The HRC Shield Proxy and GOES proton data are stored in HDF5 format here:

`/proj/sot/ska/data/arc/hrc_shield.h5`

Thanks to Peter Ford for providing the ACIS txings data.