

# GOOD COVERS AND ALGEBRAS ON CONICALLY SMOOTH SPACES

ALEKSANDAR IVANOV AND ÖDÜL TETİK

ABSTRACT. We construct good covers for conically smooth spaces. By a result of Karlsson–Scheimbauer–Walde, this implies that, for every such space  $X$ , constructible factorisation algebras on  $X$  and disk algebras over  $X$  coincide. We also give a simplified proof of that result.

## CONTENTS

1.	Introduction	1
2.	Localities of the full unzip	1
3.	Good covers	4
4.	A proof of...	6
	References	6

## 1. INTRODUCTION

*Acknowledgments.* We thank Tashi Walde for very useful exchanges.

## 2. LOCALITIES OF THE FULL UNZIP

**Definition 2.1.**  $S_X \subset X$ .

**Proposition 2.2.**  $S_X \subset X$  proper constructible bundle with  $\text{Unzip}_{S_X}(X) \cong \text{Unzip}(X)$  and  $\text{Link}_{S_X}(X) = \partial \text{Unzip}(X)$ .

**Lemma 2.3.** *Locally finite good cover on  $S_X$ . (?) unnecessary - second-countability implies more than paracompactness - see [Lee, Smooth, Thm 1.15]*

**Lemma 2.4.** *Let  $X$  be a smooth manifold with corners. Then for every boundary collar  $I: \mathbb{R}_{\geq 0} \times \partial X \hookrightarrow X$  there exists a homeomorphism*

$$\alpha_I: X^+ \xrightarrow{\cong} \mathbb{R}_{\geq 0} \times X$$

where  $X^+ = \bigcup_{r \in \mathbb{R}_{\geq 0}} [0, r] \times \partial X \cup_{\{r\} \times \partial X} X$ .

*Proof.* Let  $I: \mathbb{R}_{\geq 0} \times \partial X \hookrightarrow X$  be a collar, which is a homeomorphism onto its image. We have  $X = I(\mathbb{R}_{\geq 0} \times \partial X) \cup_{I((0, \infty) \times \partial X)} X^\circ$ . Similarly, for each  $r$ , we have a diffeomorphism

$$\alpha_r: X \rightarrow X \setminus I([0, r] \times \partial X) = I([r, \infty) \times \partial X) \cup_{I((r, \infty) \times \partial X)} X \setminus I([0, r] \times \partial X)$$

---

The authors were supported by the Austrian Science Fund (FWF) through Project no. P 37046.

given by  $I(t, q) \mapsto I(t + r, q)$  on  $I(\mathbb{R}_{\geq 0} \times \partial X)$  and by the identity on  $X \setminus \text{Im}(I)$ . We suppress the dependence on  $I$  in notation. We obtain a well-defined homeomorphism

$$I \cup \alpha_r: [0, r] \times \partial X \cup_{\{r\} \times \partial X} X \rightarrow X,$$

well-defined since  $\alpha_r(q) = \alpha_r(I(0, q)) = I(r, q)$  for  $q \in \partial X$ , and thereupon, writing  $X^{+r} = [0, r] \times \partial X \cup_{\{r\} \times \partial X} X$ , the bijection

$$\begin{aligned} \alpha_I: X^+ &\rightarrow \mathbb{R}_{\geq 0} \times X, \\ \alpha|_{X^{+r}} &= \{r\} \times I \cup \alpha_r. \end{aligned}$$

We equip  $X^+$  with the induced topology, promoting  $\alpha$  to a homeomorphism.  $\square$

**Lemma 2.5.** *Let  $X$  be a smooth manifold with corners. Then there is a homeomorphism  $\mathbb{R}_{\geq 0} \times \partial(\mathbb{R}_{\geq 0} \times X) \cong \mathbb{R}_{\geq 0} \times X$ . Consequently, there is a homeomorphism*

$$\text{Unzip}_{C(L)}(C(Z)) \cong \mathbb{R}_{\geq 0} \times \text{Link}_{C(L)}(C(Z))$$

for  $L = S_Z$ .

*Proof.* Using Lemma 2.4 it suffices to provide a homeomorphism

$$\phi: \mathbb{R}_{\geq 0} \times \partial(\mathbb{R}_{\geq 0} \times X) \rightarrow \bigcup_{r \in \mathbb{R}_{\geq 0}} [0, r] \times \partial X \cup_{\{r\} \times \partial X} X.$$

Noting

$$\partial(\mathbb{R}_{\geq 0} \times X) = \{0\} \times X \cup_{\{0\} \times \partial X} \mathbb{R}_{\geq 0} \times \partial X,$$

we define  $\phi$  to be the following map:

$$\begin{aligned} \mathbb{R}_{\geq 0} \times \{0\} \times X &\ni (t, 0, x) \mapsto x \in X \subset [0, t] \times \partial X \cup_{\{t\} \times \partial X} X \\ \mathbb{R}_{\geq 0} \times \mathbb{R}_{\geq 0} \times \partial X &\ni (t, s, q) \mapsto (t, q) \in [0, t + s] \times \partial X. \end{aligned}$$

This map and its inverse  $\phi^{-1}$  given by

$$\begin{aligned} [0, r] \times \partial X &\ni (t, q) \mapsto (t, r - t, q) \in \mathbb{R}_{\geq 0} \times \mathbb{R}_{\geq 0} \times \partial X \\ [0, r] \times \partial X \cup_{\{r\} \times \partial X} X &\supset X \ni x \mapsto (r, 0, x) \in \mathbb{R}_{\geq 0} \times \{0\} \times X \end{aligned}$$

are well-defined and continuous. Note that  $\phi$  does not depend on a collar.

The second statement is the special case where  $X = \text{Unzip}_L(Z)$  using Proposition 2.2 and that  $\text{Unzip}_{C(L)} C(Z) = \mathbb{R}_{\geq 0} \times \text{Unzip}_L(Z)$  and  $\text{Link}_{C(L)} C(Z) = \partial \text{Unzip}_{C(L)}(C(Z))$ .  $\square$

*Remark 2.6.* In the situation of Lemma 2.4 we will regard  $X^+$  as a smooth manifold with corners with respect to the smooth structure induced by  $\alpha_I$ . Up to equivalence, this structure does not depend on the choice of collar. Similarly, in the situation of Lemma 2.5 we will regard  $\mathbb{R}_{\geq 0} \times \text{Link}_{C(L)} C(Z)$  as a smooth manifold with corners, tautologically diffeomorphic to  $\text{Unzip}_{C(L)} C(Z)$  with respect to the induced smooth structure.

**Construction 2.7.** Let  $X$  be a smooth manifold with corners of dimension  $n$ , let  $\{(U, \phi_U)\}$  a cover of  $X$  by coordinate neighbourhoods where each  $\phi_U: \mathbb{R}^{n-c_U} \times \mathbb{R}_{\geq 0}^{c_U}$  is a homeomorphism, and let  $\{\rho_U: X \rightarrow [0, 1]\}$  be a partition of unity subordinate to this cover. Recall that a collar  $I = I(\{U, \phi_U, \rho_U\}): \mathbb{R}_{\geq 0} \times \partial X \hookrightarrow X$  is then constructed by defining to be the flow along the nowhere-vanishing inward-pointing vector field  $V = \sum \rho_U V_U$  where, in local coordinates,  $V_U = \sum_{1 \leq i \leq c_U} \partial_i$  where  $\{\partial_i\}$  is the standard basis of  $T_0 \mathbb{R}_{\geq 0}^{c_U} \subset T_0(\mathbb{R}^{n-c_U} \times \mathbb{R}_{\geq 0}^{c_U})$ .

Let  $X$  and  $I = I_{\{U, \phi_U, \rho_U\}}$  be as above. Then there is a canonically induced a collar

$$I^+ = I(\{\mathbb{R}_{\geq 0} \times U, \text{id}_{\mathbb{R}_{\geq 0}} \times \phi_U, \rho_U \circ \text{pr}_X\}): \mathbb{R}_{\geq 0} \times \partial(\mathbb{R}_{\geq 0} \times \partial X) \hookrightarrow \mathbb{R}_{\geq 0} \times X$$

on  $\mathbb{R}_{\geq 0} \times X$ , where  $\text{pr}_X: \mathbb{R}_{\geq 0} \times X \rightarrow X$  is the coordinate projection. It is the flow along the vector field  $V^+ = \sum(\rho_U \circ \text{pr}_X) \cdot V_U^+$  where  $V_U^+ = \partial_s + V_U$  where  $\partial_s$  is the standard basis of  $T_0\mathbb{R}_{\geq 0} \subset T_0(\mathbb{R}_{\geq 0} \times \mathbb{R}^{n-c_U} \times \mathbb{R}_{\geq 0}^{c_U})$ .

**Lemma 2.8.** *Let  $X$  be a smooth manifold with corners and let  $I^+: \mathbb{R}_{\geq 0} \times \partial(\mathbb{R}_{\geq 0} \times \partial X) \hookrightarrow \mathbb{R}_{\geq 0} \times X$  be as in Construction 2.7. Then*

$$I^+ = \alpha_I \circ \phi$$

where  $\alpha_I$  is as in Lemma 2.4 and  $\phi$  is as in the proof of Lemma 2.5.

*Proof.* We observe the restrictions

$$\begin{aligned} \mathbb{R}_{\geq 0} \times \{0\} \times X &\xrightarrow{\alpha_I \circ \phi} \mathbb{R}_{\geq 0} \times X \\ (t, 0, x) &\mapsto \begin{cases} (t, I(t_x + t, q)), & x = I(t_x, q) \\ (t, x), & x \in X \setminus \text{Im}(I) \end{cases} \end{aligned}$$

and

$$\begin{aligned} \mathbb{R}_{\geq 0} \times \mathbb{R}_{\geq 0} \times \partial X &\xrightarrow{\alpha_I \circ \phi} \mathbb{R}_{\geq 0} \times X \\ (t, s, q) &\mapsto (t + s, I(t, q)). \end{aligned}$$

Both maps are the flow (for time  $t$ ) along the vector field  $V^+ = \sum(\rho_U \circ \text{pr}_X) \cdot (\partial_s + V_U)$ .  $\square$

In the following,  $\mathbb{D}^n \subset \mathbb{R}^n$  denotes the unit open  $n$ -disk and  $C^{<1}(Z) = * \amalg_{\{0\} \times Z} [0, 1) \times Z$ .

**Definition 2.9.** We say an embedded basic  $\phi: \mathbb{R}^k \times C(Z) \hookrightarrow X$  in a conically smooth space  $X$  is *extendable* if there exists an embedding  $\hat{\phi}: \mathbb{R}^n \times C(Z) \hookrightarrow X$  such that  $\phi$  factors as  $\phi: \mathbb{R}^k \times C(Z) \rightarrow \mathbb{D}^n \times C^{<1}(Z) \hookrightarrow \mathbb{R}^n \times C^{<1}(Z) \xrightarrow{\hat{\phi}} X$  where the first map is the isomorphism given by the cartesian product of the isomorphisms  $\mathbb{R}^n \rightarrow \mathbb{D}^n$ ,  $x \mapsto \frac{x}{|x|+1}$  and  $C(Z) \rightarrow C^{<1}(Z)$ ,  $(t, z) \mapsto (\frac{t}{t+1}, z)$ .

**Definition 2.10.** Suppose  $\mathcal{U}$  is a cover of a topological space  $X$  which is closed under finite intersections. We say  $\mathcal{U}$  is *generated* by a cover  $\mathcal{V}$  and write  $\mathcal{U} = \langle \mathcal{V} \rangle$  if every member of  $\mathcal{U}$  is a finite intersection of members of  $\mathcal{V}$ .

**Lemma 2.11.** *Every smooth manifold  $M$  has a good cover  $\mathcal{U}$  which is generated by extendable basics.*

*Proof.* Equip  $M$  with a riemannian metric. We can put  $\mathcal{U} = \langle \mathcal{V} \rangle$  for  $\mathcal{V} = \{D_p\}_{p \in M}$  where  $D_p$  is the convex disk which is the interior of the image of a ball in  $T_p M$  under the exponential map, with a radius that is strictly smaller than the radius of injectivity.  $\square$

**Lemma 2.12.** *Suppose  $U \subset C(Z)$  be conically smooth basics with  $S_{C(Z)} \cong U$ , and suppose that  $U \cong C(L)$  is extendable. Let  $\pi: \text{Link}(C(Z))|_U \rightarrow U$  denote the link projection of  $\text{Unzip}(C(Z))$  over  $U$ . Then there is an isomorphism*

$$\overline{\partial\pi^{-1}U} \cong \text{Link}_L(Z)$$

and an induced conically smooth collar

$$(0, 1] \times \partial \overline{\pi^{-1}U} \hookrightarrow \overline{\pi^{-1}U}$$

which is a refinement onto its image.

*Proof.* Up to isomorphism we have that  $U = C^{<1}(L) \subset C^{<1}(Z) \subset C(Z)$  is induced by the inclusion  $L = S_Z \subset Z$ . Abbreviating  $S = S_{C(Z)}$ , the inclusion of  $U$  factors as  $U \subset S = C(L)$ . Now, using Proposition 2.2 we have  $\text{Unzip } C(Z) = \text{Unzip}_S C(Z) = \mathbb{R}_{\geq 0} \times \text{Unzip}_L Z$  and so

$$\begin{aligned} \text{Unzip } C(Z)|_U &= \text{Link}_{C(L)} C(Z)|_{C^{<1}(L)} \\ &= \{0\} \times \text{Unzip}_L Z \cup_{\{0\} \times \text{Link}_L Z} [0, 1] \times \text{Link}_L Z \end{aligned}$$

since the projection  $\pi: \text{Link}_{C(L)} C(Z) = \{0\} \times \text{Unzip}_L Z \cup_{\{0\} \times \text{Link}_L Z} \mathbb{R}_{\geq 0} \times \text{Link}_L Z \rightarrow C(L) = * \amalg_{\{0\} \times L} \mathbb{R}_{\geq 0} \times L$  is given by mapping all of  $\{0\} \times \text{Unzip}_L Z$  to  $*$  and on  $\mathbb{R}_{\geq 0} \times \text{Link}_L Z$  by  $\text{id}_{\mathbb{R}_{\geq 0}} \times \pi'$  where  $\pi': \text{Link}_L Z \rightarrow L$  is the link projection. Thus

$$\pi^{-1}\overline{U} = \pi^{-1}C^{\leq 1}(U) = \{0\} \times \text{Unzip}_L Z \cup [0, 1] \times \text{Link}_L Z$$

and consequently

$$\pi^{-1}\partial\overline{U} = \pi^{-1}\{1\} \times L = \{1\} \times \text{Link}_L Z,$$

proving the first claim. The collar is immediate.  $\square$

*Example 2.13.*

**Corollary 2.14.**

### 3. GOOD COVERS

**Definition 3.1.** Let  $X$  be a CSS. A *good cover* of  $X$  is an open cover  $\mathcal{U}$  of  $X$  consisting of basics whose finite intersections are also in  $\mathcal{U}$ .

**Definition 3.2.** Let  $Y$  be a smooth manifold with or without boundary, equipped with a riemannian metric. We call a cover of  $Y$  *geodesically convex* if it consists of geodesically convex basics.

**Theorem 3.3.** *Let  $X$  be a CSS over a depth-1 poset. Then  $X$  has a good cover.*

*Proof.* Without loss of generality, suppose  $X$  is stratified over  $[1]$  and let  $M = X_0$  and  $N = X_1$ . Recall the blow-up of  $M$ , the smooth manifold

$$\text{Unzip} \cong L \amalg_{L \times (0, \infty)} N$$

with boundary  $L = \partial \text{Unzip}$ . We write  $\pi: L \rightarrow M$  for the accompanying proper fibre bundle. Let us equip  $\text{Unzip}$  with a riemannian metric which splits along the boundary.

Using the paracompactness of  $M$ , let  $\mathcal{U}$  be a locally finite good cover of  $M$  which trivialises  $\pi$ . For each  $U \in \mathcal{U}$ , let  $\epsilon_U > 0$  be the radius of injectivity in the normal direction on the compact closure  $\overline{U}$  so that there is a smooth embedding

$$I_U: \pi^{-1}U \times [0, \epsilon_U) \hookrightarrow \text{Unzip}$$

whose restriction  $I_U|_{\pi^{-1}U \times \{0\}}$  to  $\pi^{-1}U \subset L$  is given by the boundary inclusion, and the path  $I_U(q, -): [0, \epsilon_U) \hookrightarrow \text{Unzip}$ , for every  $q \in \pi^{-1}U$ , is the minimising geodesic

given by the normal exponential map. More precisely, for  $\nu$  the inward-pointing unit normal vector field along the boundary, we set  $I_U(x, t) = \exp_x(t\nu_x)$ .<sup>1</sup> Let

$$C = \bigcup_{U \in \mathcal{U}} \text{Im}(I_U)$$

be the induced ‘collar’ and consider the cover

$$\mathcal{C} = \{I_U(\pi^{-1}U \times [0, \delta]) : U \in \mathcal{U}, \delta \leq \epsilon_U\}$$

of  $C$ . Let us write  $C_{U, \delta} = I_U(\pi^{-1}U \times [0, \delta]) \in \mathcal{C}$ . Note that  $\mathcal{C}$  is closed under finite intersections since so is  $\mathcal{U}$ : we have  $\pi^{-1}U \cap \pi^{-1}V = \pi^{-1}(U \cap V)$  and

$$C_{U, \delta} \cap C_{V, \delta'} = C_{U \cap V, \min(\delta, \delta')} \in \mathcal{C}$$

since  $\min(\delta, \delta') \leq \epsilon_{U \cap V}$  for  $\delta \leq \epsilon_U, \delta' \leq \epsilon_V$ .

Consider now the collection  $\mathcal{V}$  of those convex geodesic disks  $V$  in  $N$  such that  $V \cap C_{U, \delta}$  is either empty or a convex geodesic disk for every  $C_{U, \delta} \in \mathcal{C}$ . Then  $\mathcal{V}$  is a good cover of  $N$ . To prove this, it suffices to show that every  $p \in C$  has a neighbourhood  $V_p \in \mathcal{V}$ , since convex geodesic disks are closed under finite intersections and so  $V \cap V' \cap C_{U, \delta} = V \cap V'' \in \mathcal{V}$ . Let now  $p \in I_U(\pi^{-1}U \times [0, \delta]) \in \mathcal{C}$ , which uniquely determines a point  $l_p = \pi(\text{pr}_1(p)) \in \pi^{-1}U \subset L$ . Let  $W \subset \pi^{-1}U$  be a convex geodesic disk neighbourhood of  $l_p$ . Now, since  $\mathcal{U}$  is locally finite, so is the cover  $\tilde{\mathcal{C}} = \{I_U(\pi^{-1}U \times [0, \epsilon_U])\}$  of  $C$  (which is not necessarily closed under finite intersections), so in particular  $p$  is contained within finitely many members  $I_{U_i}(\pi^{-1}U_i \times [0, \epsilon_i]) \in \tilde{\mathcal{C}}$ . We necessarily have  $p \in I_U(\pi^{-1}U \times [0, \min_i(\epsilon_i)])$  with  $U \in \{U_i\}_i$  an open satisfying  $\epsilon_U = \min_i(\epsilon_i)$ . Then  $W \times [0, \min_i(\epsilon_i)] \ni p$  is a convex geodesic half-disk since the riemannian metric on Unzip splits along the boundary, and so we have  $p \in V_p = W \times (0, \min_i(\epsilon_i)) \in \mathcal{V}$ .

Let us now observe that  $\pi^{-1}U \cong L_p \times U$  for  $p \in U$  implies that each

$$C(L_p) \times U \cong U \amalg_{\pi^{-1}U} C_{U, \delta} \subseteq X$$

is a basic in  $X$ . Thus, writing  $\widehat{C_{U, \delta}} = U \amalg_{\pi^{-1}U} C_{U, \delta}$ , we obtain that

$$\widehat{\mathcal{C}} = \{\widehat{C_{U, \delta}} : C_{U, \delta} \in \mathcal{C}\}$$

is a cover by basics of the ‘tubular neighbourhood’

$$\widehat{C} = \bigcup_{U \in \mathcal{C}} \widehat{C_{U, \epsilon_U}} \subseteq X$$

of  $M$ . It is closed intersections since so is  $\mathcal{C}$ . Finally, since for  $V \in \mathcal{V}$  we have  $\widehat{C_{U, \delta}} \cap V = C_{U, \delta} \cap V \in \mathcal{V}$ , we conclude that

$$\widehat{\mathcal{C}} \cup \mathcal{V}$$

is a good cover of  $X$ . □

---

<sup>1</sup>A global  $\epsilon$  need not exist unless Unzip is of bounded geometry; see e.g. Schick [Sch01].

## 4. A PROOF OF...

## REFERENCES

- [Sch01] T. Schick. ‘Manifolds with Boundary and of Bounded Geometry’. *Mathematische Nachrichten* 223.1 (2001), 103–120.

UNIVERSITY OF VIENNA, FACULTY OF PHYSICS, MATHEMATICAL PHYSICS GROUP, BOLTZMANNGASSE  
5, 1090 VIENNA, AUSTRIA

*Email address:* `aleksandar.ivanov@univie.ac.at`

*Email address:* `oeduel.tetik@univie.ac.at`