

Activity dependent development of maps in the visual system

Activity type

Sensory

- species
 - mammals
 - non-mammalian vertebrates
 - invertebrates
- ~~ignore this Sensory literature for now~~ — too much literature for a 2000 word review limit?
 - *limit review to before vision onset?*
 - just do very brief overview in intro with statement that here we focus on intrinsic activity patterns in visual system and refer to other recent reviews?
- When?
 - Before eye opening and experienced visual patterns
 - Melanopsin
 - After eye opening
 - Hubel and Wiesel

Spontaneous

- What?
 - species
 - mammals
 - can occur before vision— long gestational timecourse
 - non-mammalian vertebrates
 - does not occur before vision— short gestation
 - patterns
 - in vitro
 - early development - before eye opening
 - retinal waves
 - TODO: shatz, Wong, Feller work
 - TODO: J. Zhou work
 - TODO: Subplate references, P. Kanold
 - Calcium waves— Konnerth, Moody, etc
 - after eye opening
 - TODO: Cossart visual cortex?
 - in vivo
 - early development - before eye opening
 - Rat

- Correlated bursting among RGCs [1]
- 'Spindle bursts'
 - spindle shaped field potential oscillations in visual cortex [2]
- 'Slow activity transients'
 - infra-slow LFP wave with nested fast oscillations
 - field oscillations and bursting multiunit activity in visual cortex [3]
 - 87% of L4 MUA occurred during slow activity transients
 - extracellular recordings with glass electrode or multisite silicon probe
 - age P5-P7
 - peak frequency (8 - 31 Hz, mostly 18–30 Hz) and duration form single distribution (400 ms - 5 s), likely all continuous 'spindle burst' oscillations within the slow wave
 - age P9-P13
 - short duration events (<400 ms) first prominent around P9-P10 and increased frequency of these events until becoming the most common type by P13
 - very long duration (>5 s) encountered after P8 ('splitting of events with maturation')
 - two populations of events encountered:
 - long duration, beta band frequency (>5 s, 18–30 Hz)
 - short duration, alpha band frequency (<3 s, 10 Hz)
 - after P12 SATs less common and no longer dominant pattern– continuous cortical activity and slow wave sleep (delta)
- Mouse
 - retinal waves
 - Primary source of patterned activity throughout neonatal visual system [4]
 - Retinal input modulates synchronous calcium signals in cortical neurons [5]
 - they possibly recorded some retinal waves?
 - but most activity was independent of retinal input, and more likely 'spindle bursts'
 - unknown if recordings were strictly from V1 or V2 (no method for identification)
 - calcium recordings were not summed population signals
 - *which might be same as the independent spontaneous V2 activity we saw* [4]
- Human
 - 'Slow activity transients'
 - infra-slow LFP wave with nested fast oscillations
 - EEG field oscillations and bursting multiunit activity in visual cortex

[6][7]

- around eye opening
 - Ferret
 - Correlated bursting activity among LGN and visual cortical neurons [8][9]
 - Rat
 - 'Spindle bursts'
 - field oscillations and bursting multiunit activity in visual cortex [3][7]
 - 'Slow activity transients'
 - infra-slow LFP wave with nested fast oscillations
 - field oscillations and bursting multiunit activity in visual cortex [3][7]
- ~~after eye opening~~
 - *too much literature in adult, limit review to before vision onset?*
 - monkey, cat, ferret, rodent, etc
 - patterned activity
 - Recent Konnerth peri-eye opening calcium imaging paper with direction selectivity?
 - intrinsic signal imaging
 - Stryker work
 - A. Grinvald work
 - multicell recordings
 - any multichannel recordings in newborn monkey (Hubel/Wiesel just did single electrodes?)
 - ~~fast traveling waves~~
 - adult (Y. Dan paper) [Benucci:2007]

- Where?

- Retina
 - Retinal waves propagate among RGCs [4]
- LGN
 - Mouse
 - Spontaneous bursting among dLGN neurons sensitive to retinal input [10]
 - Inferred by matched retinal driven patterns in V1 and SC [4]
 - Ferret
 - Correlated retinal driven activity among LGN neurons [8]
 - multielectrode array recordings
 - neonatal, awake behaving ferret
 - recordings performed just before eye opening in ferret, around P21
 - an age when spontaneous activity likely to be glutamatergic driven waves involving on-off bipolar cells and when retinotopic refinement and ODC formation complete
 - binocular correlations present
 - they state it can't be from retina, cause of statistical independence
 - aspriating cortex (cortico-thalamic projections) abolishes the binocular correlation
 - *but damage due to the ablation?*
 - contralateral inputs drove LGN bursting more strongly than ipsilateral
 - without retinal input, corticothalamic input necessary to sustain LGN neuron

bursting

- Superior colliculus
 - Retinal waves drive collicular neurons [4]
- Visual cortex
 - Mouse
 - Retinal waves propagate to cortical neurons [4]
 - Retinal input modulates synchronous calcium signals in cortical neurons [5]
 - Rat
 - Retinal input can drive spiking multiple unit activity in cortical neurons [2]
 - Retinal input can drive slow activity transients and spiking multiple unit activity in cortical neurons before and around eye opening [3]
 - Ferret
 - Correlated bursting among V1 neurons [9]
 - multielectrode array recordings
 - neonatal, awake behaving ferret
 - recordings performed just before eye opening in ferret, between P24-P29
 - an age when spontaneous activity likely to be glutamatergic driven waves involving on-off bipolar cells and when retinotopic refinement and ODC formation complete
 - Human
 - Slow activity transients in EEG signal in preterm human infants [6][7]
- When?
 - Before eye opening and experienced visual patterns
 - Before birth for some species
 - rabbit [11][12], monkey [13], human [6][7], cat [14]
 - After birth for some species
 - rodent [4][2][3], rabbit [11][12], ferret [14][15][16][8][9]
 - After eye opening
 - experiential pattern replay/dreams
 - analogs to hippocampal - place cell replays (Wilson work) for learning and memory?
 - Y. Dan visual pattern replay paper
 - standing waves and fast travelling waves [#Benucci:2007]
- Why?
 - activity dependent circuit establishment and refinement– lessons from other systems
 - cite R. Wong review
 - *short blurb on other systems*
 - chick neuromuscular junction (lichtmann sanes)
 - spontaneous motor circuit activity V. Hamburger & (Petterssen Nature paper)
 - activity dependent visual map development
 - anatomical - structural
 - axon growth
 - xenopus, zebrafish literature?
 - LGN and SC
 - rodent
 - mouse

- TODO: beta2 nAChR ko mouse?
 - TODO: N. Spitzer reference on activity-dep Ca2+ growth *Cortex
 - Ruthazer and Olavarria paper
- axon refinement
 - xenopus, zebrafish literature?
 - LGN and SC
 - rodent
 - mouse
 - Retinocollicular axon - retinotopic map refinement requires cholinergic retinal waves [\[17\]](#)
 - beta2 nAChR ko mice, focal Dil tracer injections into retina
 - beta2 nAChR ko and transgenic mice show that nAChR mediated spontaneous activity in the retina is essential for retinotopic map refinement, eye specific segregation [\[18\]](#)
 - RGC refinement and deficits in beta2 nAChR ko occurs at the level of single RGCs [\[19\]](#)
 - cortex
 - cortico-collicular axon arborizations [\[20\]](#).
 - Nice Dil reconstructions of cortico-collicular axons in rat
 - Ruthazer and Olavarria paper
- dendrite growth?
 - cortico-collicular recipient cells in SC,
 - Recent constantine-paton paper [\[20\]](#).
 - Cortico-collicular axons needed for 'caliber 3' dendritic filopodia density.
 - Eye opening regulates spine density in 'caliber 3' dendrites
 - Golgi or Dil analysis in ferret, cat, monkey, or rodent cortex?
 - Ruthazer and Olavarria paper
 - Golgi or Dil analysis in LGN or SC?
- dendritic refinement
 - spine dynamics?
 - TODO: xenopus literature?, H. Cline
- cell migration
 - rodent
 - cortex
 - interneurons
 - TODO: recent Fishell paper
 - TODO: recent ZJ Huang papers
 - TODO: Ben-Ari, JB Manent activity dependent interneuron migration in vitro model
 - higher mammals
 - cortex
 - retinal wave evidence in macaque monkey retina in vitro [\[13\]](#)
 - occurs as early as E60, before start of eye specific segregation
 - macaque monkey neocortical neurogenesis (E40, SP/L6), (E56 – E100, L5 – L2/3)

- *unknown, but due to gestational timing, waves likely to have greater influence on cell migration and cortical patterning than in lower mammals*
 - both excitatory and inhibitory cell migration overlaps significantly with the period for retinal waves
- functional - physiological
 - synapse maturation
 - retinocollicular synapse
 - Increased AMPA/NMDA ratios and AMPA quantal amplitudes during first postnatal week [21]
 - burst activation in vitro capable of inducing LTP [21]
 - delayed maturation and greater LTP at beta2-/- nAChR ko synapses [21]
 - retinotopy
 - Retinotopic map refinement requires retinal waves in SC [17]
 - beta2 nAChR ko mice, focal Dil tracer injections into retina
 - preferential anatomical terminal zone elongation along nasal-temporal axis
 - Disrupted retinotopic map in beta2 nAChR ko mice in LGN (first order connections) [22]
 - using tungsten microelectrode extracellular recordings
 - physiological receptive fields preferentially disrupted (elongated) along nasal-temporal (visual field azimuth) axis
 - Abnormal segregation of on- and off-centered cells in LGN that is not seen in wt
 - *precocious glutamate bipolar cell mediated waves?*
 - Altered retinotopic map in beta2 nAChR ko mice in SC (first order connections) [23]
 - using tungsten microelectrode extracellular recordings
 - physiological receptive fields elongated along nasal-temporal axis
 - Altered retinotopic map in beta2 nAChR-/- mice in SC (first order connections) [24]
 - using intrinsic signal imaging
 - retinotopic map preferentially disrupted (elongated) along anterior-posterior (nasal-temporal) axis of SC
 - Altered retinotopic map in beta2 nAChR -/- mice in V1 (second order connections) [25]
 - Intrinsic signal imaging of mouse V1 for visual space map
 - Extracellular microelectrode recordings for single cell receptive fields
 - Preferential disruption (elongation, scatter, response amplitude) along the visual space azimuth (nasal-temporal axis)
 - They speculate that waves regulate ephrinA gradients to explain the nasal-temporal disruption since travelling waves had not been found to have a preferred direction at the time
 - Cortico-collicular alignment of retinotopy (quaternary order connections, L5 -> SC) [26]
 - Transgenic mice, tracer injections, intrinsic signal functional mapping
 - Used ephA3ki/ki (knock in) mice crossed with beta2 nAChR -/- mice for the crucial experiment in Figure 6.
 - These mice have duplicated retinocollicular map, but only a single, non-matched corticocollicular projection when no cholinergic waves are present.
- eye specific segregation
- ocular dominance columns
 - development of ODCs in ferret [27]

- epibatidine injections and tracer injections
 - ocular dominance bias index with extracellular microelectrode recordings
 - spontaneous cholinergic activity in retina required for cortical ODC formation
 - TODO: Crair, Stryker
 - orientation selectivity
 - TODO: Crair, Stryker
 - TODO: Recent Fitzpatrick work
 - TODO: ongoing J. Cang unpublished work? (look at abstr from SFN, our CSHL conf last year)
 - direction selectivity
 - TODO: Recent Konnerth peri-eye opening calcium imaging paper
 - TODO: Recent Fitzpatrick work (the reprogramming of selectivity)
- How?
- Permissive
 - Spatiotemporal pattern does not matter
 - Perhaps just absolute levels of activity needed?
 - homeostasis, cellular growth and survival?
 - Informative
 - Spatiotemporal pattern does matter
 - Temporal activity pattern
 - Time scale
 - eye-specific segregation
 - before eye opening
 - Synchronous activation of RGCs in both eyes with ChR2 disrupts eye-specific segregation in SC and LGN [28]
 - Relevant window for spike timing differences of RGCs in both eyes within 100s of milliseconds [28]
 - chR2 stimulation of RGCs, anatomical segregation analysis in SC
 - Mechanism
 - Coincident pre-post synaptic activity - Hebbian plasticity
 - Dependent on NMDA-R?
 - maybe yes?
 - TODO:
 - HP Xu recent work?
 - maybe not?
 - LTD independent of NMDA-R activation in mouse [29]
 - in vitro explant with extracellular field potentials and high freq stim to mimic retinal waves
 - bidirectional maturation
 - finds LTD early between birth and eye opening
 - finds LTP after eye opening through critical period

- cites [Butts:2007] for bidirectional synaptic strength changes in single LGN cells
 - L-type calcium channel plateau potentials at developing LGN neurons [Lo:2002]
 - retinogeniculate PSC bursting is independent of NMDAR activation (NMDAR1 ko mice, ex vivo, extracellular)[10]
 - HP Xu recent work?
- Independent of NMDA-R?
 - endocannabinoid induced LTD?
 - but this type of activation still requires NMDA activation? [30]
 - this type of coincidence detection reviewed elsewhere [30]
 - mGluR-VSCC-IP3R-eCB coincidence detector [31]
 - this form of LTD independent of postsynaptic NMDA receptors
 - utilizes metabotropic glutamate receptors, voltage sensitive calcium channels, IP3 stores, and endocannabinoid receptors
 - detects firing coincidence at 125 ms time scale (*versus 25 ms time scale for NMDA dependent LTP*)
 - described at L4 to L2/3 synapses in somatosensory cortex
 - bistable switch in spike statistics for postsynaptic neurons?
 - critical level of coincident presynaptic activity needed to cause spike?
 - during early development?
 - biophysical membrane and cable properties different in immature neurons
 - more voltage gated calcium conductance
 - less sodium channels
 - lower fidelity spike transmission initially?
 - Non-coincident – alternate, lagged timing based plasticity rule?
- Spatial activity pattern
 - Unknown: no direct experiment yet
 - experiment needed: to control spatio-temporal activity patterns before start of vision
 - Analogous to the classic Sensory activation experiments
 - owl prism experiments Knudsen
 - cat goggle experiments
 - Instructive role of spatial activation hinted at by Hong-Ping's paper? [18]

- Necessitates the temporal activity mechanisms with an additional spatial dimension via a wavefront
- Spatial patterns setup in periphery (RGCs) and communicated across levels of visual organization [4]
- Spatiotemporal information in retinal waves TODO: make table of these properties?
 - *Perform quick analysis of wavefront size within our in vivo waves dataset and include as a new data in a Figure?*
 - Wave speed from in vitro and in vivo literature consistent (100s of microns/sec)
 - retina in vitro (100 - 500 $\mu\text{m/s}$; mouse, rabbit, ferret, monkey) TODO: zhou, feller, feldheim, chalupa papers
 - SC in vivo (28 - 60 $\mu\text{m/s}$ = 90 - 180 $\mu\text{m/s}$ in retina with 3x scale factor) [4]
 - Wave size (wave area) smaller within in vitro literature [32] and bigger in vivo [4]
 - Wave propagation failure in vitro because of some combination of in vitro conditions such as bath medium, temperature, and cut connections?
 - Wave direction along VT - DN axis in vitro [32] and in vivo [4]
 - This makes the wavefront axis perpendicular to preferred axis for activity mediated refinement

Figure: Schematic of visual cortex primary and secondary areas, pathway illustration, and summary of retinal influence? (incl Olavarria work?)

References

1. Maffei, L. and Galli-Resta, L. (1990). Correlation in the discharges of neighboring rat retinal ganglion cells during prenatal life, *Proc Natl Acad Sci U S A*, 87(7), 2861–2864
2. Hanganu, I. L., Ben-Ari, Y., and Khazipov, R. (2006). Retinal waves trigger spindle bursts in the neonatal rat visual cortex, *J Neurosci*, 26(25), 6728–36
3. Colonnese, M. T. and Khazipov, R. (2010). “Slow activity transients” in infant rat visual cortex: a spreading synchronous oscillation patterned by retinal waves, *J Neurosci*, 30(12), 4325–37
4. Ackman, J. B., Burbridge, T. J., and Crair, M. C. (2012). Retinal waves coordinate patterned activity throughout the developing visual system, *Nature*, 490(7419), 219–25
5. Siegel, F., Heimel, J. A., Peters, J., and Lohmann, C. (2012). Peripheral and central inputs shape network dynamics in the developing visual cortex in vivo, *Curr Biol*, 22(3), 253–8
6. Vanhatalo, S., Palva, J. M., Andersson, S., Rivera, C., Voipio, J., and Kaila, K. (2005). Slow endogenous activity transients and developmental expression of K⁺-Cl⁻ cotransporter 2 in the immature human cortex, *Eur J Neurosci*, 22(11), 2799–804
7. Colonnese, M. T., Kaminska, A., Minlebaev, M., Milh, M., Bloem, B., Lescure, S., Moriette, G., Chiron, C., Ben-Ari, Y., and Khazipov, R. (2010). A conserved switch in sensory processing prepares developing neocortex for vision, *Neuron*, 67(3), 480–98
8. Weliky, M. and Katz, L. C. (1999). Correlational structure of spontaneous neuronal activity in the developing lateral geniculate nucleus in vivo, *Science*, 285(5427), 599–604
9. Chiu, C. and Weliky, M. (2002). Relationship of correlated spontaneous activity to functional ocular dominance columns in the developing visual cortex, *Neuron*, 35(6), 1123–1134
10. Mooney, R., Penn, A. A., Gallego, R., and Shatz, C. J. (1996). Thalamic relay of spontaneous retinal activity prior to vision, *Neuron*, 17(5), 863–874

11. Zhou, Z. J. and Zhao, D. (2000). Coordinated transitions in neurotransmitter systems for the initiation and propagation of spontaneous retinal waves, *J Neurosci*, 20(17), 6570–6577
12. Zheng, J., Lee, S., and Zhou, Z. J. (2006). A transient network of intrinsically bursting starburst cells underlies the generation of retinal waves, *Nat Neurosci*, 9(3), 363–71
13. Warland, D. K., Huberman, A. D., and Chalupa, L. M. (2006). Dynamics of spontaneous activity in the fetal macaque retina during development of retinogeniculate pathways, *J Neurosci*, 26(19), 5190–7
14. Meister, M., Wong, R. O., Baylor, D. A., and Shatz, C. J. (1991). Synchronous bursts of action potentials in ganglion cells of the developing mammalian retina, *Science*, 252(5008), 939–943
15. Wong, R. O., Meister, M., and Shatz, C. J. (1993). Transient period of correlated bursting activity during development of the mammalian retina, *Neuron*, 11(5), 923–38
16. Feller, M. B., Wellis, D. P., Stellwagen, D., Werblin, F. S., and Shatz, C. J. (1996). Requirement for cholinergic synaptic transmission in the propagation of spontaneous retinal waves, *Science*, 272(5265), 1182–7
17. McLaughlin, T., Torborg, C. L., Feller, M. B., and O’Leary, D. D. M. (2003). Retinotopic map refinement requires spontaneous retinal waves during a brief critical period of development, *Neuron*, 40(6), 1147–1160
18. Xu, H.-p., Furman, M., Mineur, Y. S., Chen, H., King, S. L., Zenisek, D., Zhou, Z. J., Butts, D. A., Tian, N., Picciotto, M. R., and Crair, M. C. (2011). An instructive role for patterned spontaneous retinal activity in mouse visual map development, *Neuron*, 70(6), 1115–27
19. Dhande, O. S., Hua, E. W., Guh, E., Yeh, J., Bhatt, S., Zhang, Y., Ruthazer, E. S., Feller, M. B., and Crair, M. C. (2011). Development of Single Retinofugal Axon Arbors in Normal and beta2 Knock-Out Mice, *J Neurosci*, 31(9), 3384–99
20. Phillips, M. A., Colonnese, M. T., Goldberg, J., Lewis, L. D., Brown, E. N., and Constantine-Paton, M. (2011). A synaptic strategy for consolidation of convergent visuotopic maps, *Neuron*, 71(4), 710–24
21. Shah, R. D. and Crair, M. C. (2008). Retinocollicular synapse maturation and plasticity are regulated by correlated retinal waves, *J Neurosci*, 28(1), 292–303
22. Grubb, M. S., Rossi, F. M., Changeux, J. P., and Thompson, I. D. (2003). Abnormal functional organization in the dorsal lateral geniculate nucleus of mice lacking the beta 2 subunit of the nicotinic acetylcholine receptor, *Neuron*, 40(6), 1161–72
23. Chandrasekaran, A. R., Plas, D. T., Gonzalez, E., and Crair, M. C. (2005). Evidence for an instructive role of retinal activity in retinotopic map refinement in the superior colliculus of the mouse, *J Neurosci*, 25(29), 6929–38
24. Mrsic-Flogel, T. D., Hofer, S. B., Creutzfeldt, C., Cloez-Tayarani, I., Changeux, J.-P., Bonhoeffer, T., and Hubener, M. (2005). Altered map of visual space in the superior colliculus of mice lacking early retinal waves, *J Neurosci*, 25(29), 6921–6928
25. Cang, J., Rentería, R. C., Kaneko, M., Liu, X., Copenhagen, D. R., and Stryker, M. P. (2005). Development of precise maps in visual cortex requires patterned spontaneous activity in the retina, *Neuron*, 48(5), 797–809
26. Triplett, J. W., Owens, M. T., Yamada, J., Lemke, G., Cang, J., Stryker, M. P., and Feldheim, D. A. (2009). Retinal input instructs alignment of visual topographic maps, *Cell*, 139(1), 175–85
27. Huberman, A. D., Speer, C. M., and Chapman, B. (2006). Spontaneous retinal activity mediates development of ocular dominance columns and binocular receptive fields in v1, *Neuron*, 52(2), 247–254
28. Zhang, J., Ackman, J. B., Xu, H.-P., and Crair, M. C. (2011). Visual map development depends on the temporal pattern of binocular activity in mice, *Nat Neurosci*, 15(2), 298–307
29. Ziburkus, J., Dilger, E. K., Lo, F.-S., and Guido, W. (2009). LTD and LTP at the developing retinogeniculate synapse, *J Neurophysiol*, 102(6), 3082–90
30. Duguid, I. and Sjöström, P. J. (2006). Novel presynaptic mechanisms for coincidence detection in synaptic plasticity, *Curr Opin Neurobiol*, 16(3), 312–22
31. Bender, V. A., Bender, K. J., Brasier, D. J., and Feldman, D. E. (2006). Two coincidence detectors for spike timing-dependent plasticity in somatosensory cortex, *J Neurosci*, 26(16), 4166–77

32. Stafford, B. K., Sher, A., Litke, A. M., and Feldheim, D. A. (2009). Spatial-temporal patterns of retinal waves underlying activity-dependent refinement of retinofugal projections, *Neuron*, 64(2), 200–12