

Opinion in Neurobiology

Elsevier Editorial System(tm) for Current
Manuscript Draft

Manuscript Number:

Title: Role of emergent neural activity in visual map development

Article Type: 24 Neural maps

Corresponding Author: Prof. Michael Crair,

Corresponding Author's Institution:

First Author: James B Ackman

Order of Authors: James B Ackman; Michael Crair

- Spontaneous retinal waves drive activation of circuits throughout the developing amniote visual system before the start of visual experience.
- Visual map development in the midbrain, thalamus, and visual cortex is regulated by retinal waves.
- The informational complexity of retinal waves is suitable for establishing multiple intra- and inter-hemispheric visual connections before the start of visuomotor behavior.
- Combinations of new techniques will assist in understanding how emergent activity shapes transcriptional regulation and circuit development.

Title

Role of emergent neural activity in visual map development

Author names and affiliations

James B. Ackman¹ and Michael C. Crair^{1,2,3}

¹Department of Neurobiology, ²Department of Ophthalmology and Visual Science, ³Kavli Institute for Neuroscience, Yale University School of Medicine, New Haven CT, 06510

Corresponding author

michael.crair@yale.edu

+1-203-785-5768

Department of Neurobiology

Yale University School of Medicine

P.O. Box 208001

New Haven, CT 06520-8001

Abstract

The structural and functional development of visual circuits in reptiles, birds, and mammals happens largely independent of sensory experience. Innate genetic programs code for gradients of molecules that provide gross positional information for developing nerve cells, yet much of the cytoarchitectural complexity and synaptogenesis of neurons depends on calcium influx, neurotransmitter release, and neural activity before the start of vision. In fact, specific spatiotemporal patterns of neural activity, or ‘retinal waves’, emerge amidst the development of the earliest connections made between excitable cells in the developing eye. These patterns of spontaneous activity, which have been observed in all amniote retinae examined to date, may be an evolved adaptation for species with long gestational periods before the onset of functional vision, imparting an informational robustness and redundancy to guide development of visual maps across the nervous system. Recent experiments indicate that retinal waves play a crucial role in the development of interconnections between different parts of the visual system, suggesting that these spontaneous patterns serve as a template-matching mechanism to prepare higher-order visually-associative circuits for the onset of visuomotor learning and behavior. Key questions for future studies include determining the exact sources and nature of spontaneous activity during development, characterizing the interactions between neural activity and transcriptional gene regulation, and understanding the extent of circuit connectivity governed by retinal waves within and between sensory-motor systems.

Introduction

The visual system is organized hierarchically such that information relayed through the optic nerve is processed in a set of neural maps distributed throughout the thalamus, midbrain, and cerebral cortex. As with all biological tissues, construction of the neural circuits underlying these visual maps fundamentally depends on innate genetic programs. However, the nervous system is unique in the extent to which it is shaped by changes during development in the external world detected and relayed as nerve impulses by the sensory organs. The extraordinary developmental plasticity of the nervous system is underscored in the visual deprivation experiments conducted by Hubel and Wiesel [1], and is experienced by each of us as we learn our native language as infants but struggle to learn a second

language as adults [2].

Given the many pieces of information that must be parsed from the photic signals in the eye to deconstruct and reconstruct the visual scene, one might think that visual experience is absolutely required to establish the neural circuits that produce complex visual representations in the brain. The debate between Herring and von Helmholtz in the 19th century over whether the binocular coordination of eye movements [3] required for stereoscopic vision was innate or learned highlights the always relevant problem regarding the relative roles for nature and nurture in developmental neuroscience. Though complex brain and behavioral development in the absence of learned experience is not surprising in animals – e.g. infant suckling in newborns – it is still remarkable that a considerable amount of intricate visual circuit structure and function forms before any patterned visual experience. As noted by Hubel and Wiesel [1] based on their recordings from primary visual cortex in newborn monkey:

That area 17 is in so many respects wired up and ready to go when the animal is born is perhaps not so surprising if one remembers that the machinery of area 17 represents building blocks of vision...

Indeed both their studies and subsequent work by others has demonstrated that a number of complex visual circuits are formed by the time of birth or eyelid opening, including cortical maps for retinotopy, orientation selectivity, and ocular dominance. Thus, with so many circuits established before birth, it might seem that by knowing the genetic code and its patterns of expression we would understand all the instructions underlying visual map formation. However, the genetic code does not work in isolation– the information required for organism development stems from the myriad interactions that occur within and between a cell and its environment throughout the course of gestation. A factor particularly relevant to the development and function of the nervous system is the spontaneous electrical activity and neurotransmitter release that emerges among excitable cells in the developing nervous system, even in isolated cultures of immature neurons. Intriguingly, a number of studies have provided experimental evidence that specific *patterns* of neural activity are required for proper visual map formation in rodent, ferret, and cat, indicating that spontaneous activity in the immature nervous system has a distinct function in circuit development. These spontaneous patterns of activity increase the informational complexity that the genome can carry by serving to sculpt neural architecture and synaptic connections so that the nervous system can begin performing computations relevant for visual behavior, learning, and memory at the start of vision. As any mother who has felt their baby kick inside the womb can attest, the developing brain is not quiescent. And pediatric neurologists would direct our attention to the fetal brain malformations that occur when chemicals that interfere with neurotransmission are taken during pregnancy.

In this review we discuss recent work that highlights the function of spontaneous activity in visual system development. We will focus on studies that have examined the nature of spontaneous activity in the developing brain as well as studies that have provided direct experimental evidence for the function of retinal waves in visual map development. For more information about the early role of genes or the later role of experience in visual development, we refer the reader to recent reviews on molecular signaling gradients, such as Eph/Ephrins for visual circuit development [4, 5], as well as on the mechanisms by which patterned spontaneous activity is generated within developing neural circuits [6], and how activity shapes circuit development after the onset of patterned vision [7, 8].

Spontaneous activity in the visual system: What, When, and Where?

In developing networks, it is unknown precisely how spontaneous oscillations of electrical activity arise, but there are numerous examples demonstrating that they do— such as the spontaneous activity found in cultures of isolated neurons [9], as well as immature spinal cord [10], retina [11, 12], hippocampus and cortex [13, 14]. The function of spontaneous correlated activity may be to strengthen the synaptic weights between coactive cells through voltage-dependent calcium influx that mediates downstream changes in transcriptional regulation. Adjustments in gene expression could result in synaptic modifications through changes in ion channel or neurotransmitter receptor function, changes in receptor clustering at existing synapses, or formation of entirely new synapses with nearby cells.

So when and where do these spontaneous activities occur that can mediate these developmental changes? Spontaneous oscillations occur after eye opening and maturation of the visual system is complete, such as standing waves and fast travelling waves [15, 16], which may serve a purpose in experiential pattern replay during sleep [17]. Coordinated waves of spontaneous activity also occur in the visual system before the onset of visual experience in all amniote species that have been examined [18]. These travelling ‘retinal waves’ of spiking activity sweep across the retina before hatching or birth in a wide variety of species including chicken, turtle, rabbit, monkey, rodent and cat (Table 1). In some species that are particularly immature at birth, such as rodent, rabbit, cat and ferret, retinal waves continue postnatally before eyelid opening. Amniote vertebrates have a long gestational time course that provides the opportunity, and potentially the necessity, for these patterns of spontaneous activity to develop. In non-amniote vertebrates, there is only a brief gestational period before the beginning of locomotor and visuomotor behavior, so the roles subserved by spontaneous patterned activity in these species is likely mediated primarily by sensory experience, and indeed spontaneous waves have not been found in non-amniotes [19].

Table 1. Spontaneous activity in developing retina

Species	Age of vision onset	Patterned activity observed?	Experimental method	Age observed	Preparation
fish	3 (zebrafish, oviparous); 21 (guppies, ovoviviparous)	no; ?			
xenopus	3	no	MEA [19]	2–4	in vitro
turtle	60	yes	SU [20]	40	in vitro
chick	21	yes	MCI [21, 22]	8–11 [21]; 13–18 [22]	in vitro
rabbit	42 (E31+P11)	yes	MCI [23], MEA [24]	26–37 [23]; 24–31 [24]	in vitro
mouse, rat	34,36 (E21,E23+P13)	yes	MCI [25, 26], MEA [27, 28], SU [11]	17, 21–34 [25]; 23–32 [28]; 30, 32–34, 36 [27]; 18–21 [11]; 24–30 [26]	in vitro [25, 27, 28], in vivo [11, 26]
ferret	74 (E42+P32)	yes	MCI [29, 30], MEA [12, 31]	47–63 [12]; 42–72 [31]; 46–54 [29]; 42–52 [30]	in vitro
cat	73 (E64+P9)	yes	MEA [12]	52, 65	in vitro
monkey	165 (E165+P0)	yes	MEA [32]	60, 67, 71, 76	in vitro

Notes: Ages reported as total days of embryonic + postnatal development. Age of vision onset refers to age of birth, eye opening, hatching, or retinal function. Abbreviations: MCI, multicellular calcium imaging; MEA, multielectrode array; SU, paired single-units

What kind of activity is in the visual system outside the retina during development? Spontaneous calcium waves have been observed in vitro in several areas of rodent lateral and medial entorhinal cortex, temporal cortex, and fronto-parietal cortex [33, 34, 35, 36], but have not been directly observed in vitro within primary visual cortex. Spontaneous UP-DOWN states have been recorded after eye opening between P14-P21 in mouse visual cortex using calcium imaging and patch recordings in vitro [37].

In the white matter and cortical layer 6, transient populations of pioneer cells called subplate neurons are thought to play a key role in cortical development. Subplate neurons may relay input activity patterns to immature cortical neurons that have yet to form functional synapses with thalamic axons [38]. Several experiments using pharmacology and targeted ablation of subplate neurons indicate that this cell population is important for excitatory and inhibitory synapse maturation as well as the formation of functional visual maps for orientation selectivity and ocular dominance [38]. Evidence that subplate neurons form a disynaptic relay between thalamocortical inputs and developing cortical neurons comes from in vitro recordings in kitten visual cortex [39] and from in vitro recordings in rodent somatosensory cortex [38]. Bursting activity in the subplate layer is known to precede spindle burst oscillations in neonatal rat somatosensory cortex in vivo [40], but the ability for subplate neurons to generate or relay activity in the visual system in vivo is unknown.

Though in vitro experiments are very powerful at dissecting local circuits, ultimately an understanding of the nature and function of neural activity in a complex and extended neural circuit must be achieved in the intact system. To this end, increasingly refined in vivo analyses of activity patterns during early development have been recorded in the visual system of ferrets and rodents (Table 2). Indeed, in EEG recordings from occipital cortex in preterm human infants, slow waves with nested fast oscillations [41, 42] are thought to be homologous to the ‘slow activity transients’ (SATs) in rat visual cortex before eye opening [43]. During early development to before eye opening, most spiking activity in L4 occurs within the field potential oscillations of SATs, which are sensitive to the presence of retinal input [43], suggesting that they are likely triggered by the spiking activity that occurs between local retinal ganglion cells (RGCs) [11] during retinal waves [12, 26].




Figure 1. Retinal waves drive patterned activity in the developing visual system. Image shows summed population activity after ablation of retinal input from the left eye in P6 mouse. Calcium signals are color coded by time of retinal wave front propagation in the SC-L. Notice the lack of activity in SC-R and V1-R. Data are based on unpublished results from [26], and consist of merged fields of view from two subsequent recordings (80 s total; SC-L+SC-R with V1-L or V1-R), overlaid on an Allen Developing Mouse Brain Atlas P4 reference illustration.

Recent calcium imaging studies have proved invaluable for revealing the spatial patterns of spontaneous activity in the visual system. The first in vivo recordings of retinal waves was recently accomplished by recording pre-synaptic calcium signals from RGC terminals in the superior colliculus (SC) in mice. Together with post-synaptic recordings from populations of collicular and cortical neurons, these experiments reveal that retinal waves are the primary source of patterned neural activity throughout the developing visual system before vision [26] (Figure 1), providing spatially consistent information about retinotopic and eye specific organization to multiple hierarchical circuits. These experiments also highlight the importance of examining spontaneous activity in the intact animal, as several fundamental features of retinal waves, such as wave size and direction, are quite different from that observed in reduced (in vitro) preparations. Multicellular two-photon calcium imaging from local clusters of visual cortical neurons also shows that retinal input modulates synchronous calcium signals in cortical neurons [44]. Calcium events with high synchrony among visual cortical neurons can occur independent of retinal input [44], though it's not clear whether anesthetic agents might differentially affect emergent activity patterns, as anesthesia is known to block retinal wave generation [26, 43]. It is also uncertain whether non-retinally driven calcium events might reflect spindle burst oscillations [45] or synchronous bursting activity in V2, which can activate V1 cells in the border regions of V1 [26]. In fact, spindle burst oscillations may be the LFP representation of synchronous, short duration calcium activations recorded in V2 and along the V1/V2 border [26]. In the future, it will be important to

combine targeted extracellular electrophysiological recordings with wide-field and two-photon calcium imaging to establish a coherent understanding of the variety of activity patterns described with the different LFP and imaging approaches. This will help resolve whether the various population activity patterns that have been described in different studies are truly different or just due to differences in experimental methodologies and recording techniques. Some or all of the spindle burst oscillations and continuous cortical activity described in visual cortex [43, 45] may occur independent of retinal waves, and may predominate in secondary visual cortical areas, representing feedback from other sensory areas such as somatosensory cortex where spindle burst oscillations predominate during tactile-motor behavior in the neonatal rat.

Just before and around eye opening, several lines of evidence suggest that the pattern of activity that predominates in the visual system is markedly different than earlier in development. In vivo recordings in ferret show that correlated bursting activity among lateral geniculate nucleus (LGN) and primary visual cortex (V1) neurons is modulated, but not exclusively driven by retinal input [46, 47]. In rat, the SATs encountered earlier in development still predominate, but the patterns split into two populations, a longer duration, higher frequency bursting activity and a short duration, lower frequency bursting activity [43]. Just before eye opening, SATs become less common and are no longer the predominate pattern, with continuous cortical activity and slow wave sleep becoming dominant [43].

Table 2. Spontaneous activity in the developing visual system in vivo

Study	Species	Area	Areal mapping	Observed pattern	Experimental method	Age observed	Anesthetic during recording
[26]	mouse	SC, V1, V2	wave front retinotopy	Retinal waves among RGC axons and cells in SC, V1, V2; retina-independent population synchrony in V2 and around V1 borders	WF-MCI and 2P-MCI	24–30	no
[44]	mouse	V1		Low synchrony events triggered by retina; High synchrony events independent of retina	2P-MCI	29–31	yes
[11]	rat	retina		Spontaneous correlated bursting activity in retina	SU	18–21	yes
[45]	rat	retina, V1	optic nerve stimulation	Spontaneous bursting activity in retina; Cortical bursting activity and ‘spindle burst’ oscillations in V1 triggered by retina	MEA, LFP, MUA, PC	24–29	no
[43]	rat	V1	VEP	‘Slow activity transients’: infra-slow wave and cortical bursting activity triggered by the retina with nested spindle burst oscillations; short duration bursting activity independent of retina	EEG, MEA, MUA, PC	28–30, 32–36	no
[46]	ferret	LGN	VEP	Long duration correlated bursting activity modulated by retinal input; short duration high synchrony activity independent of retina	MEA, LFP, MUA	66–69	no
[48]	ferret	V1	stereotaxic	Long duration correlated ‘macroburst’ activity modulated by retinal input; short duration high synchrony activity independent of retina	MEA, LFP, MUA	64–70	no
[41, 49]	preterm human infant	occipital	stereotaxic	‘Slow activity transients’: infra-slow wave with nested fast oscillations	EEG	224–322	no

Notes: Ages reported as total days of embryonic + postnatal development. Abbreviations: WF- and 2P-MCI, wide-field and two-photon multicellular calcium imaging; EEG, electroencephalography; MEA, multielectrode array; LFP, local field potential; MUA, multiple unit activity; PC, patch-clamp; SU, paired single-units

Regulation of cellular anatomy and physiology by spontaneous activity

Though visual experience has a clear role in directing the refinement of synaptic connectivity [7], much of the topography of neural circuits and the elaboration of cellular morphology are established early in development, before viable sensory-driven activity [50, 51, 52, 53]. Molecular factors, like the signaling gradients provided by Eph receptor tyrosine kinases and their Ephrin ligands, mediate much of this elaboration in the developing visual system [54]. The initial connectivity between axons and their target cells that occurs before vision produces functional neuronal networks that give rise to central features of visual circuitry and processing, such as retinotopy, ocular dominance, and orientation selectivity. However, key aspects of the cellular morphologic and functional synapse maturation that occurs before vision relies on spontaneous neuronal activity. At the retinogeniculate synapse, blockade of spontaneous activity by TTX application to the eye around and before the time of eye opening prevents the normal developmental increase in synapse strength and arrests synaptic pruning [55], while visual deprivation (by delaying eye opening) has no effect. An important model for disrupting spontaneous retinal activity utilizes the *Chrn2* mutant ($\beta 2^{-/-}$) mouse, which lacks a subunit of the nicotinic acetylcholine receptor and is thought to have disrupted retinal waves [25, 56], though some controversy persists on this point [57]. Functional maturation at the retinocollicular synapse is impaired in $\beta 2^{-/-}$ mice during the first week after birth [58], similar to the effect of TTX application to the eye on retinogeniculate synapses, again suggesting that spontaneous retinal activity promotes the maturation of retinogufal synapses before normal vision.

Some of the morphological and functional changes in neuronal properties brought about by obstructing activity during development can be considered homeostatic responses to the withdrawal of synaptic drive [59]. For example, chronic treatment with TTX can produce compensatory changes in synapse number, strength and/or cellular excitability [59], and visual deprivation increases the intrinsic excitability of neurons in the visual cortex [60]. Remarkably, interfering with spontaneous retinal activity, as is thought to occur in $\beta 2^{-/-}$ mice, disrupts retinocollicular synapse number and strength, but the overall visual drive of neurons in the SC remains unaltered [61]. This suggests that developing neurons are sensitive to both local synaptic conditions that modulate individual synapse strength as well as global activity levels in a way that keeps the network balanced and presumably functionally optimal.

The morphological development of neuronal dendrites and axon arbors in the developing visual system is also sensitive to activity before vision. Classic experiments demonstrated that the abolition of spiking activity in the developing cat brain with fetal infusions of TTX prevents eye-specific segregation in the dLGN and results in RGC axon arbors that are abnormally large and diffuse [62, 63]. Equivalent experiments examining the sensitivity of geniculocortical afferents to retinal activity blockade reveals similar results [64, 65], though the interpretation of these experiments is complicated by the observation that nascent ocular dominance columns in the visual cortex are present before activity blockade was begun, indeed before normal vision is possible [52, 53, 66]. RGC axon arbors in the SC, and to a lesser extent the lateral geniculate nucleus, are also disrupted in $\beta 2^{-/-}$ mice [67], and display abnormally diffuse and widespread arborizations. The morphological development of retinorecipient neurons in the SC is shaped by activity [68] through a competitive process [69]. Furthermore, higher order efferent connections from visual cortex (Figure 2) may be regulated by spontaneous retinal waves. For example, retinal input regulates inter- and intra-hemispheric projections during a restricted developmental period in rat and ferret, coinciding with the time when retinal waves occur [70, 71, 72].

What links the morphological and functional development of visual system neurons and synapses?




Figure 2. Efferent connectivity from mouse visual cortex. Reference image (top left) adapted from [Allen Mouse Brain Atlas](#). Axonal projections (right and bottom panels) are color coded by target structure and were rendered using two primary visual cortex injection datasets (black dots indicate injection location) available from the [Allen Mouse Brain Connectivity Atlas](#).

During the developmental window that retinocollicular and retinogeniculate synapse maturation is sensitive to ongoing spontaneous activity, these synapses also exhibit long-term potentiation and depression (LTP/LTD) [58, 73]. Activity-dependent synaptic plasticity in the developing visual system through a Hebb-type process provides a plausible mechanistic link between spontaneous activity and the functional and structural maturation of synapses.

Regulation of visual map structure and function by spontaneous activity

In addition to the maturation of individual neuron synapses, dendrites and axon arbors, the development of large-scale neural circuits or ‘maps’ for visual stimulus features, such as retinotopy, ocular dominance and orientation, are also sensitive to the presence of ongoing spontaneous activity before eye opening. Disrupting retinal waves pharmacologically or genetically interferes with the development of both retinotopy and eye specific segregation in the dLGN and SC of mice [28, 74, 75]. In the SC, retinocollicular target zones and individual axon arbors are enlarged in $\beta 2^{-/-}$ mice, and preferentially elongated along the nasal-temporal axis of the retina, corresponding to the visual field azimuth [74] [67]. Functional response properties of SC neurons are correspondingly impacted, with receptive fields dramatically enlarged, particularly along the visual field azimuth [74, 76]. In the dLGN

of in $\beta 2^{-/-}$ mice, retinal ganglion cell axon arbors are also modestly enlarged [67], but rather than producing expanded receptive fields in individual dLGN neurons, the overall retinotopic map in the dLGN of $\beta 2^{-/-}$ mice is disrupted due to the scatter of receptive field locations, particularly along the visual field azimuth [77]. The biased effect in $\beta 2^{-/-}$ mice on retinotopic maps along the visual field azimuth may stem from the strong bias in retinal wave propagation along this axis [26, 56] and/or differences in the effectiveness of molecular processes in mapping these circuits along the two visual axes. Remarkably, in both the dLGN [77] and SC [61] of $\beta 2^{-/-}$ mice, the organization of response properties associated with ON- or OFF-selectivity, which is not normally observed in mice, emerge *de novo*. In the dLGN of $\beta 2^{-/-}$ mice, ON- and OFF-center cells are spatially segregated [77], while in the SC, neurons become selective to either the onset or the offset of light stimulus, when they normally respond equally well to both [74]. Thus, it appears that circuit organization around new response features emerge in the dLGN and SC of $\beta 2^{-/-}$ mice, possibly because of the delayed functional development of retinofugal synapses [55, 58] and the precocious appearance of glutamate receptor mediated waves in $\beta 2^{-/-}$ mice [25], which have distinct correlation properties in ON- and OFF- retinal ganglion cells [78, 79].

Disruptions in neural circuit development in $\beta 2^{-/-}$ mice are not confined to first order, retinofugal connections. Anatomical and functional response properties in higher order visual circuits, including geniculate-cortical connections [80] and cortico-collicular connections [81] are also perturbed. Moreover, the anatomical and functional receptive field anomalies in $\beta 2^{-/-}$ mice are accompanied by behavioral deficits in visual function [82]. Several other genetic models that appear to disrupt spontaneous retinal activity and visual circuit development in mice have also recently emerged. These manipulations perturb ongoing retinal activity in a variety of ways, and appear to cause corresponding disruptions in visual circuit development [83, 84]. In contrast, disrupting spontaneous retinal activity after eye opening has no effect on circuit development [85, 86]. The results of pharmacological manipulations of retinal activity in the dLGN and SC of mice [74, 87] and the dLGN and visual cortex of ferrets [75] [88] [89] [90] are also generally consistent with genetic manipulations, though some contradictions persist [91]. Optogenetic manipulation of retinal activity before vision suggests that some features of neural circuit development, like retinotopy, are more sensitive to the levels of spontaneous activity, while other features, like eye specific segregation, are sensitive to the timing of activity in the two eyes [86]. Resolution of lingering uncertainties about the precise role of different spatiotemporal features of spontaneous retinal activity in the emergence of visual circuits will require a careful analysis of spontaneous activity *in vivo* in the various genetic and pharmacological models, and a parametric analysis of the role of activity in the development of the diverse features of visual circuitry, including retinotopy and eye segregation.

In addition to maps for retinotopy and ocular dominance, visual circuits for orientation emerge without vision [52, 92, 93]. Binocular matching of orientation preference also occurs without vision in cats and ferrets [52, 92], though in mice orientation matching is degraded without vision [93]. In ferrets, robust direction selectivity requires visual experience [92], but initial experience-independent biases in direction preference are predictive of the future elaboration of direction selectivity brought about by visual experience [94]. In contrast, in mice cortical direction selectivity is independent of vision [95]. Differences between ferret, cat and mice in the apparent role of vision in the emergence of various visual receptive field properties could be due differences in the neural circuits underlying these response features (mice don't have maps for orientation or direction, though they do have orientation and direction selective cells [96] that are tuned to the same orientation in the two eyes [97]). Alternatively, differences in the developmental timing or properties of spontaneous activity relative to neural circuit maturation in these various species may lead to variable effects of visual experience on map development.

Conclusion

It seems increasingly apparent that many, if not all, developing neural circuits display spontaneous activity before they become functionally efficacious. In animals with extended gestation, fundamental neural circuit features emerge without sensory experience, particularly in the visual system. Patterned spontaneous neural activity appears capable of substituting for peripheral sensation in shaping the development of neural circuits. Major open questions in the field concern the degree to which spontaneous activity instructs developing neural circuits, and how activity interacts with gene expression to regulate synapse, neuron and neural circuit development. Hebbian synaptic plasticity mechanisms ('fire-together, wire-together') are well suited to link activity-dependent synapse and neural circuit development, but distinct activity-dependent mechanisms that are sensitive to levels or timing of activity but not correlations in activity likely also play important roles in sculpting developing neural circuits. The increasing availability of modern optical approaches to examine and manipulate activity in developing neural circuits in vivo suggests that a fundamental understanding of the rules that regulate synapse and neural circuit development in the complex mammalian brain is at hand.

Acknowledgements

This work was supported by NIH grants P30 EY000785 and R01 EY015788 to M.C.C.

References

1. Hubel, D. H. & Wiesel, T. N. (1977) Ferrier lecture. Functional architecture of macaque monkey visual cortex. *Proc R Soc Lond B Biol Sci* 198, 1-59.
2. Kuhl, P. K. (2010) Brain mechanisms in early language acquisition. *Neuron* 67, 713-27.
3. King, W. M. & Zhou, W. (2000) New ideas about binocular coordination of eye movements: is there a chameleon in the primate family tree?. *Anat Rec* 261, 153-61.
4. Huberman, A. D., Feller, M. B., & Chapman, B. (2008) Mechanisms underlying development of visual maps and receptive fields. *Annu Rev Neurosci* 31, 479--509.
5. Cang, J. & Feldheim, D. A. (2013) Developmental Mechanisms of Topographic Map Formation and Alignment. *Annu Rev Neurosci* , .
6. Blankenship, A. G. & Feller, M. B. (2010) Mechanisms underlying spontaneous patterned activity in developing neural circuits. *Nat Rev Neurosci* 11, 18-29.
7. Levelt, C. N. & Hübener, M. (2012) Critical-period plasticity in the visual cortex. *Annu Rev Neurosci* 35, 309-30.
8. Espinosa, J. S. & Stryker, M. P. (2012) Development and plasticity of the primary visual cortex. *Neuron* 75, 230-49.
9. Mazzoni, A., Broccard, F. D., Garcia-Perez, E., Bonifazi, P., Ruaro, M. E., & Torre, V. (2007) On the dynamics of the spontaneous activity in neuronal networks. *PLoS ONE* 2, e439.

10. Marder, E. & Rehm, K. J. (2005) Development of central pattern generating circuits. *Curr Opin Neurobiol* 15, 86-93.
11. Maffei, L. & Galli-Resta, L. (1990) Correlation in the discharges of neighboring rat retinal ganglion cells during prenatal life. *Proc Natl Acad Sci U S A* 87, 2861-2864.
12. Meister, M., Wong, R. O., Baylor, D. A., & Shatz, C. J. (1991) Synchronous bursts of action potentials in ganglion cells of the developing mammalian retina. *Science* 252, 939-943.
13. Leinekugel, X., Khazipov, R., Cannon, R., Hirase, H., Ben-Ari, Y., & Buzsáki, G. (2002) Correlated bursts of activity in the neonatal hippocampus in vivo. *Science* 296, 2049-52.
14. Khazipov, R. & Luhmann, H. J. (2006) Early patterns of electrical activity in the developing cerebral cortex of humans and rodents. *Trends Neurosci* 29, 414-8.
15. Benucci, A., Frazor, R. A., & Carandini, M. (2007) Standing waves and traveling waves distinguish two circuits in visual cortex. *Neuron* 55, 103-17.
16. Han, F., Caporale, N., & Dan, Y. (2008) Reverberation of recent visual experience in spontaneous cortical waves. *Neuron* 60, 321--327.
17. Lee, A. K. & Wilson, M. A. (2002) Memory of sequential experience in the hippocampus during slow wave sleep. *Neuron* 36, 1183-94.
18. Wong, R. O. (1999) Retinal waves and visual system development. *Annu Rev Neurosci* 22, 29-47.
19. *Demas, J. A., Payne, H., & Cline, H. T. (2012) Vision drives correlated activity without patterned spontaneous activity in developing *Xenopus* retina. *Dev Neurobiol* 72, 537-46.
Amphibians are visually responsive, but lack spontaneous patterned retinal activity during similar developmental stages in which amniotes lack visual response but show spontaneous retinal activity. This suggests that 'retinal waves' are an evolutionary adaptation in animals that lack vision for extended periods of gestation during visual system development.
20. Sernagor, E. & Grzywacz, N. M. (1999) Spontaneous activity in developing turtle retinal ganglion cells: pharmacological studies. *J Neurosci* 19, 3874-87.
21. Catsicas, M., Bonness, V., Becker, D., & Mobbs, P. (1998) Spontaneous Ca²⁺ transients and their transmission in the developing chick retina. *Curr Biol* 8, 283-6.
22. Wong, W. T., Sanes, J. R., & Wong, R. O. (1998) Developmentally regulated spontaneous activity in the embryonic chick retina. *J Neurosci* 18, 8839-52.
23. Zhou, Z. J. & Zhao, D. (2000) Coordinated transitions in neurotransmitter systems for the initiation and propagation of spontaneous retinal waves. *J Neurosci* 20, 6570--6577.
24. Syed, M. M., Lee, S., He, S., & Zhou, Z. J. (2004) Spontaneous waves in the ventricular zone of developing mammalian retina. *J Neurophysiol* 91, 1999-2009.
25. Bansal, A., Singer, J. H., Hwang, B. J., Xu, W., Beaudet, A., & Feller, M. B. (2000) Mice lacking

specific nicotinic acetylcholine receptor subunits exhibit dramatically altered spontaneous activity patterns and reveal a limited role for retinal waves in forming ON and OFF circuits in the inner retina. *J Neurosci* 20, 7672-81.

26. **Ackman, J. B., Burbridge, T. J., & Crair, M. C. (2012) Retinal waves coordinate patterned activity throughout the developing visual system. *Nature* 490, 219-25.
Retinal waves are present and propagate throughout the visual system in neonatal mice *in vivo*. Unexpectedly, waves are not random, and show biases in their nucleation, propagation and some bilateral correspondence.
27. Demas, J., Eglen, S. J., & Wong, R. O. L. (2003) Developmental loss of synchronous spontaneous activity in the mouse retina is independent of visual experience. *J Neurosci* 23, 2851-60.
28. McLaughlin, T., Torborg, C. L., Feller, M. B., & O'Leary, D. D. M. (2003) Retinotopic map refinement requires spontaneous retinal waves during a brief critical period of development. *Neuron* 40, 1147-1160.
29. Wong, R. O., Chernjavsky, A., Smith, S. J., & Shatz, C. J. (1995) Early functional neural networks in the developing retina. *Nature* 374, 716-8.
30. Feller, M. B., Wellis, D. P., Stellwagen, D., Werblin, F. S., & Shatz, C. J. (1996) Requirement for cholinergic synaptic transmission in the propagation of spontaneous retinal waves. *Science* 272, 1182-7.
31. Wong, R. O., Meister, M., & Shatz, C. J. (1993) Transient period of correlated bursting activity during development of the mammalian retina. *Neuron* 11, 923-38.
32. Warland, D. K., Huberman, A. D., & Chalupa, L. M. (2006) Dynamics of spontaneous activity in the fetal macaque retina during development of retinogeniculate pathways. *J Neurosci* 26, 5190-7.
33. Garaschuk, O., Linn, J., Eilers, J., & Konnerth, A. (2000) Large-scale oscillatory calcium waves in the immature cortex. *Nat Neurosci* 3, 452-9.
34. Peinado, A. (2000) Traveling slow waves of neural activity: a novel form of network activity in developing neocortex. *J Neurosci* 20, RC54.
35. Corlew, R., Bosma, M. M., & Moody, W. J. (2004) Spontaneous, synchronous electrical activity in neonatal mouse cortical neurones. *J Physiol* 560, 377-90.
36. Namiki, S., Norimoto, H., Kobayashi, C., Nakatani, K., Matsuki, N., & Ikegaya, Y. (2013) Layer III Neurons Control Synchronized Waves in the Immature Cerebral Cortex. *J Neurosci* 33, 987-1001.
37. Cossart, R., Aronov, D., & Yuste, R. (2003) Attractor dynamics of network UP states in the neocortex. *Nature* 423, 283-8.
38. Kanold, P. O. & Luhmann, H. J. (2010) The subplate and early cortical circuits. *Annu Rev Neurosci* 33, 23-48.
39. Friauf, E. & Shatz, C. J. (1991) Changing patterns of synaptic input to subplate and cortical plate

during development of visual cortex. *J Neurophysiol* 66, 2059-71.

40. Yang, J.-W., Hanganu-Opatz, I. L., Sun, J.-J., & Luhmann, H. J. (2009) Three patterns of oscillatory activity differentially synchronize developing neocortical networks in vivo. *J Neurosci* 29, 9011-25.
41. Vanhatalo, S., Palva, J. M., Andersson, S., Rivera, C., Voipio, J., & Kaila, K. (2005) Slow endogenous activity transients and developmental expression of K⁺-Cl⁻ cotransporter 2 in the immature human cortex. *Eur J Neurosci* 22, 2799-804.
42. *Colonnese, M. T., Kaminska, A., Minlebaev, M., Milh, M., Bloem, B., Lescure, S., Moriette, G., Chiron, C., Ben-Ari, Y., & Khazipov, R. (2010) A conserved switch in sensory processing prepares developing neocortex for vision. *Neuron* 67, 480-98.
Spontaneous oscillatory activity in the occipital cortex of humans before birth. Intrinsic changes in visual cortex activity patterns occur before the onset of visual experience in both human and rat, likely due to an evolutionarily conserved innate developmental program.
43. **Colonnese, M. T. & Khazipov, R. (2010) "Slow activity transients" in infant rat visual cortex: a spreading synchronous oscillation patterned by retinal waves. *J Neurosci* 30, 4325-37.
Slow activity transients are the predominant spontaneous EEG signals in the developing visual cortex, synchronizing all cortical layers and requiring retinal input. Slow activity transients are likely the electrophysiological signature of retinal waves transmitted to visual cortex.
44. Siegel, F., Heimel, J. A., Peters, J., & Lohmann, C. (2012) Peripheral and central inputs shape network dynamics in the developing visual cortex in vivo. *Curr Biol* 22, 253-8.
45. Hanganu, I. L., Ben-Ari, Y., & Khazipov, R. (2006) Retinal waves trigger spindle bursts in the neonatal rat visual cortex. *J Neurosci* 26, 6728-36.
46. Weliky, M. & Katz, L. C. (1999) Correlational structure of spontaneous neuronal activity in the developing lateral geniculate nucleus in vivo. *Science* 285, 599-604.
47. Chiu, C. & Weliky, M. (2002) Relationship of correlated spontaneous activity to functional ocular dominance columns in the developing visual cortex. *Neuron* 35, 1123--1134.
48. Chiu, C. & Weliky, M. (2001) Spontaneous activity in developing ferret visual cortex in vivo. *J Neurosci* 21, 8906--8914.
49. Tolonen, M., Palva, J. M., Andersson, S., & Vanhatalo, S. (2007) Development of the spontaneous activity transients and ongoing cortical activity in human preterm babies. *Neuroscience* 145, 997-1006.
50. Rakic, P. (1976) Prenatal genesis of connections subserving ocular dominance in the rhesus monkey. *Nature* 261, 467--471.
51. Holt, C. E. & Harris, W. A. (1983) Order in the initial retinotectal map in *Xenopus*: a new technique for labelling growing nerve fibres. *Nature* 301, 150--152.
52. Crair, M. C., Gillespie, D. C., & Stryker, M. P. (1998) The role of visual experience in the

development of columns in cat visual cortex. *Science* 279, 566-70.

53. Crowley, J. C. & Katz, L. C. (2000) Early development of ocular dominance columns. *Science* 290, 1321--1324.
54. Lemke, G. & Reber, M. (2005) Retinotectal mapping: new insights from molecular genetics. *Annu Rev Cell Dev Biol* 21, 551--580.
55. Hooks, B. M. & Chen, C. (2006) Distinct roles for spontaneous and visual activity in remodeling of the retinogeniculate synapse. *Neuron* 52, 281-91.
56. Stafford, B. K., Sher, A., Litke, A. M., & Feldheim, D. A. (2009) Spatial-temporal patterns of retinal waves underlying activity-dependent refinement of retinofugal projections. *Neuron* 64, 200-12.
57. Sun, C., Warland, D. K., Ballesteros, J. M., van der List, D., & Chalupa, L. M. (2008) Retinal waves in mice lacking the beta2 subunit of the nicotinic acetylcholine receptor. *Proc Natl Acad Sci U S A* 105, 13638--13643.
58. Shah, R. D. & Crair, M. C. (2008) Retinocollicular synapse maturation and plasticity are regulated by correlated retinal waves. *J Neurosci* 28, 292-303.
59. Turrigiano, G. (2011) Too many cooks? Intrinsic and synaptic homeostatic mechanisms in cortical circuit refinement. *Annu Rev Neurosci* 34, 89-103.
60. Nataraj, K. & Turrigiano, G. (2011) Regional and temporal specificity of intrinsic plasticity mechanisms in rodent primary visual cortex. *J Neurosci* 31, 17932-40.
61. Chandrasekaran, A. R., Shah, R. D., & Crair, M. C. (2007) Developmental homeostasis of mouse retinocollicular synapses. *J Neurosci* 27, 1746-55.
62. Shatz, C. J. & Stryker, M. P. (1988) Prenatal tetrodotoxin infusion blocks segregation of retinogeniculate afferents. *Science* 242, 87-9.
63. Sretavan, D. W., Shatz, C. J., & Stryker, M. P. (1988) Modification of retinal ganglion cell axon morphology by prenatal infusion of tetrodotoxin. *Nature* 336, 468-71.
64. Stryker, M. P. & Harris, W. A. (1986) Binocular impulse blockade prevents the formation of ocular dominance columns in cat visual cortex. *J Neurosci* 6, 2117--2133.
65. Antonini, A. & Stryker, M. P. (1993) Development of individual geniculocortical arbors in cat striate cortex and effects of binocular impulse blockade. *J Neurosci* 13, 3549-73.
66. Crair, M. C., Horton, J. C., Antonini, A., & Stryker, M. P. (2001) Emergence of ocular dominance columns in cat visual cortex by 2 weeks of age. *J Comp Neurol* 430, 235-49.
67. Dhande, O. S., Hua, E. W., Guh, E., Yeh, J., Bhatt, S., Zhang, Y., Ruthazer, E. S., Feller, M. B., & Crair, M. C. (2011) Development of Single Retinofugal Axon Arbors in Normal and {beta}2 Knock-Out Mice. *J Neurosci* 31, 3384-99.

68. Furman, M. & Crair, M. C. (2012) Synapse maturation is enhanced in the binocular region of the retinocollicular map prior to eye opening. *J Neurophysiol* 107, 3200-16.
69. Furman, M., Xu, H.-P., & Crair, M. C. (2013) Competition driven by retinal waves promotes the morphological and functional synaptic development of neurons in the superior colliculus. *J Neurophysiol* , .
70. Olavarria, J. F. & Safaeian, P. (2006) Development of callosal topography in visual cortex of normal and enucleated rats. *J Comp Neurol* 496, 495-512.
71. Bock, A. S., Kroenke, C. D., Taber, E. N., & Olavarria, J. F. (2012) Retinal input influences the size and corticocortical connectivity of visual cortex during postnatal development in the ferret. *J Comp Neurol* 520, 914-32.
72. Laing, R. J., Bock, A. S., Lasiene, J., & Olavarria, J. F. (2012) Role of retinal input on the development of striate-extrastriate patterns of connections in the rat. *J Comp Neurol* 520, 3256-76.
73. Butts, D. A., Kanold, P. O., & Shatz, C. J. (2007) A burst-based "Hebbian" learning rule at retinogeniculate synapses links retinal waves to activity-dependent refinement. *PLoS Biol* 5, e61.
74. Chandrasekaran, A. R., Plas, D. T., Gonzalez, E., & Crair, M. C. (2005) Evidence for an instructive role of retinal activity in retinotopic map refinement in the superior colliculus of the mouse. *J Neurosci* 25, 6929-38.
75. Stellwagen, D. & Shatz, C. J. (2002) An instructive role for retinal waves in the development of retinogeniculate connectivity. *Neuron* 33, 357-367.
76. Mrsic-Flogel, T. D., Hofer, S. B., Creutzfeldt, C., Cloez-Tayarani, I., Changeux, J.-P., Bonhoeffer, T., & Hubener, M. (2005) Altered map of visual space in the superior colliculus of mice lacking early retinal waves. *J Neurosci* 25, 6921--6928.
77. Grubb, M. S., Rossi, F. M., Changeux, J. P., & Thompson, I. D. (2003) Abnormal functional organization in the dorsal lateral geniculate nucleus of mice lacking the beta 2 subunit of the nicotinic acetylcholine receptor. *Neuron* 40, 1161-72.
78. Kerschensteiner, D. & Wong, R. O. L. (2008) A precisely timed asynchronous pattern of ON and OFF retinal ganglion cell activity during propagation of retinal waves. *Neuron* 58, 851-8.
79. Akrouh, A. & Kerschensteiner, D. (2013) Intersecting Circuits Generate Precisely Patterned Retinal Waves. *Neuron* , .
80. Cang, J., Rentería, R. C., Kaneko, M., Liu, X., Copenhagen, D. R., & Stryker, M. P. (2005) Development of precise maps in visual cortex requires patterned spontaneous activity in the retina. *Neuron* 48, 797-809.
81. Triplett, J. W., Owens, M. T., Yamada, J., Lemke, G., Cang, J., Stryker, M. P., & Feldheim, D. A. (2009) Retinal input instructs alignment of visual topographic maps. *Cell* 139, 175-85.

82. Wang, L., Rangarajan, K. V., Lawhn-Heath, C. A., Sarnaik, R., Wang, B.-S., Liu, X., & Cang, J. (2009) Direction-specific disruption of subcortical visual behavior and receptive fields in mice lacking the beta2 subunit of nicotinic acetylcholine receptor. *J Neurosci* 29, 12909-18.
83. Demas, J., Sagdullaev, B. T., Green, E., Jaubert-Miazza, L., McCall, M. A., Gregg, R. G., Wong, R. O. L., & Guido, W. (2006) Failure to maintain eye-specific segregation in nob, a mutant with abnormally patterned retinal activity. *Neuron* 50, 247-59.
84. Xu, H.-p., Furman, M., Mineur, Y. S., Chen, H., King, S. L., Zenisek, D., Zhou, Z. J., Butts, D. A., Tian, N., Picciotto, M. R., & Crair, M. C. (2011) An instructive role for patterned spontaneous retinal activity in mouse visual map development. *Neuron* 70, 1115-27.
85. *Soto, F., Ma, X., Cecil, J. L., Vo, B. Q., Culican, S. M., & Kerschensteiner, D. (2012) Spontaneous activity promotes synapse formation in a cell-type-dependent manner in the developing retina. *J Neurosci* 32, 5426-39.
Rhythmic hyperactivity of RGCs in Crx-/- mutant mice leads to enhanced synaptogenesis in some, but not all bipolar cell connections with RGCs. Central projections are unaffected. Suggests a cell-type specific modulation of synaptogenesis by spontaneous activity in the retina.
86. **Zhang, J., Ackman, J. B., Xu, H.-P., & Crair, M. C. (2011) Visual map development depends on the temporal pattern of binocular activity in mice. *Nat Neurosci* 15, 298-307.
Using chronic optogenetic techniques to stimulate retinal ganglion cells in mice before they are normally visually responsive, this paper demonstrates that the emergence of eye specific segregation in the lateral geniculate nucleus and superior colliculus of mice is sensitive to the relative timing of activity between the two eyes.
87. Pfeiffenberger, C., Cutforth, T., Woods, G., Yamada, J., Rentería, R. C., Copenhagen, D. R., Flanagan, J. G., & Feldheim, D. A. (2005) Ephrin-As and neural activity are required for eye-specific patterning during retinogeniculate mapping. *Nat Neurosci* 8, 1022-7.
88. Huberman, A. D., Speer, C. M., & Chapman, B. (2006) Spontaneous retinal activity mediates development of ocular dominance columns and binocular receptive fields in v1. *Neuron* 52, 247-254.
89. Chapman, B. (2000) Necessity for afferent activity to maintain eye-specific segregation in ferret lateral geniculate nucleus. *Science* 287, 2479-82.
90. Penn, A. A., Riquelme, P. A., Feller, M. B., & Shatz, C. J. (1998) Competition in retinogeniculate patterning driven by spontaneous activity. *Science* 279, 2108-2112.
91. Huberman, A. D., Wang, G.-Y., Liets, L. C., Collins, O. A., Chapman, B., & Chalupa, L. M. (2003) Eye-specific retinogeniculate segregation independent of normal neuronal activity. *Science* 300, 994--998.
92. Li, Y., Fitzpatrick, D., & White, L. E. (2006) The development of direction selectivity in ferret visual cortex requires early visual experience. *Nat Neurosci* 9, 676--681.
93. *Sarnaik, R., Wang, B.-S., & Cang, J. (2013) Experience-Dependent and Independent Binocular Correspondence of Receptive Field Subregions in Mouse Visual Cortex. *Cereb Cortex*.

Vision is not necessary for the development of substantial overlap in binocular receptive fields and alignment of ON and OFF subregions in visual cortical neurons. However, visual experience enhances the degree of binocular receptive field correspondence.

94. *Van Hooser, S. D., Li, Y., Christensson, M., Smith, G. B., White, L. E., & Fitzpatrick, D. (2012) Initial neighborhood biases and the quality of motion stimulation jointly influence the rapid emergence of direction preference in visual cortex. *J Neurosci* 32, 7258-66.
In ferret visual cortex, the emergence of robust maps for direction preference depends on visual experience. However, local neighborhood biases present before vision appear to nucleate the large scale organization of direction preference after vision.
95. Rochefort, N. L., Narushima, M., Grienberger, C., Marandi, N., Hill, D. N., & Konnerth, A. (2011) Development of direction selectivity in mouse cortical neurons. *Neuron* 71, 425-32.
96. Niell, C. M. & Stryker, M. P. (2008) Highly selective receptive fields in mouse visual cortex. *J Neurosci* 28, 7520--7536.
97. Wang, B.-S., Sarnaik, R., & Cang, J. (2010) Critical period plasticity matches binocular orientation preference in the visual cortex. *Neuron* 65, 246-56.

Figure 1
[Click here to download high resolution image](#)




Figure 2
[Click here to download high resolution image](#)

