

The Dynamics of Gradient Descent in Linear Least Squares Regression

Akiva Lipshitz

June 22, 2017

Arguably, one of the most powerful developments in early modern applied mathematics is that of gradient descent, which is a technique for solving programs of the form

$$\arg \max_w f(w)$$

by solving a dynamical system

$$\mathbf{w}_{t+1} := \mathbf{w} + \alpha \lambda(\mathbf{w} \mid \mathbf{x}, \mathbf{y})$$

where \mathbf{w} represents parameters and α is a learning rate.

In this paper, we show that in gradient descent optimization for regression tasks the dynamical mode (logistic, exponential, or oscillatory, etc.) of the system defined by equation (2) depends on which subset of the real line the learning rate of the system α is an element of. The boundaries of these subsets are significant as they represent bifurcation points in the dynamics of gradient descent. That such results may be theoretically derived shows that after learning rules have been written down, additional analysis must be done to understand the asymptotic behavior and stability dynamics of the dynamical system defined by the learning rules so as to appropriately choose hyperparameters.

In a broad sense, this work sheds light on the space of learning processes which will hopefully inspire new developments in the study of the actual learning process of intelligent learning systems.

Bounds on Learning Rate α for which Learning Converges

Suppose we have already derived the learning rules for a D dimensional regression from the normality assumption. Also, we have removed all constants of

proportionality in the learning equations for the sake of simplicity, which doesn't change the asymptotic behavior of learning.

Let α be a learning rate, \mathbf{x} be a T by D matrix, \mathbf{y} a T by 1 matrix, and \mathbf{w} a D dimensional row vector.

$$\begin{aligned}\Delta w_{i,t} &= -\alpha \sum_j^N (y_i - \hat{y}_j) x_j \\ &= -\alpha \sum_j^N (y_i - w_j x_{ij}) x_{ij} \\ &= -\alpha \mathbf{x}_i \cdot \mathbf{y} \frac{1}{N} + w_{i,t} \alpha \mathbf{x}_i \cdot \mathbf{x}_i \frac{1}{N}\end{aligned}$$

Observe in this linear task the dynamics of each weight w_i is independent of that of any other weight. We can simplify equation (2) by writing $\beta_{i1} = -\alpha \mathbf{x}_i \cdot \mathbf{y} \frac{1}{N}$ and $\beta_{i2} = \alpha \mathbf{x}_i \cdot \mathbf{x}_i \frac{1}{N}$, such that

$$\Delta w_{i,t} = \beta_{i1} + \beta_{i2} w_{it}$$

We would like to analyze the asymptotic behavior of this dynamical system and to do so we need an analytical expression for $w_{i,t}$. First, we will produce an update rule

$$w_{i,t+1} = \beta_{i1} + w_{it}(\beta_{i2} + 1)$$

which we recognize as a one dimensional autoregressive process with an affine term. We can recursively compose equation (5) with itself, using $w_{i,0} \sim \mathcal{N}(\mu, \sigma)$ as initial conditions. This is a bit of a tedious computation that results in a closed form polynomial expression. We simplify the indices in the computation by assuming it holds for all w_i . Thus subscripts in the computation on w refer to iterations, with dimension implied.

$$\begin{aligned}w_0 &= w_0 \\ w_1 &= \beta_1 + w_0(\beta_2 + 1) \\ w_2 &= \beta_1 + \beta_1(\beta_2 + 1) + w_0(\beta_2 + 1)^2 \\ w_3 &= \beta_1 + \beta_1(\beta_2 + 1) + \beta_1(\beta_2 + 1)^2 + w_0(\beta_2 + 1)^3 \\ w_n &= w_0(\beta_2 + 1)^n + \beta_1 \sum_{j=0}^{n-1} (\beta_2 + 1)^j\end{aligned}$$

This leads to somewhat of a closed form expression:

$$w_{it} = (\beta_{i2} + 1)^t w_{i0} + \beta_{i1} \sum_{j=1}^{t-1} (\beta_{i2} + 1)^j$$

We are interested in the limit $t \rightarrow \infty$ as it relates to α .

$$\lim_{t \rightarrow \infty} \left[w_{it} = (\beta_{i2} + 1)^t w_{i0} + \beta_{i1} \sum_{j=1}^{t-1} (\beta_{i2} + 1)^j \right] = ?$$

Both terms in (8) converge if $-1 < \beta_{i2} + 1 < 1$. Recalling from before $\beta_{i2} = \alpha \mathbf{x}_i \cdot \mathbf{x}_i \frac{1}{N}$,

$$\begin{aligned} -1 &< \beta_{i2} + 1 < 1 \\ -1 &< \alpha \frac{\|\mathbf{x}\|^2}{N} + 1 < 1 \end{aligned}$$

There are two cases to consider and we now go through them:

If

$$0 \leq \alpha \frac{\|\mathbf{x}\|^2}{N} + 1 < 1$$

then

$$-1 \leq \alpha \frac{\|\mathbf{x}\|^2}{N} < 0$$

Thus

$$-1 \leq \frac{\alpha}{N} |\mathbf{x}|^2 < 0$$

This leads to the bounds

$$-\frac{N}{\|\mathbf{x}\|^2} \leq \alpha < 0$$

The second case to consider is that of $-1 < \alpha \frac{\mathbf{x}_i \cdot \mathbf{x}_i}{N} + 1 \leq 0$. Here, using a similar thought process

$$-2 \frac{N}{\|\mathbf{x}\|^2} < \alpha \leq -\frac{N}{\|\mathbf{x}\|^2}$$

We now take the union of the sets defined by (17) and (18) as valid α values, naming it A . A is expressed in terms of its components because the inner bound is actually significant as it is the *optimal* α value that leads to convergence in one step.

$$A_i = \left(-2 \frac{N}{\|\mathbf{x}_i\|^2}, -\frac{N}{\|\mathbf{x}\|^2} \right] \cup \left[-\frac{N}{\|\mathbf{x}_i\|^2}, 0 \right)$$

A Closed Form Expression

Equation (12) could have been recognized as a geometric series and is now rewritten as such:

$$w_{it} = (\beta_{i2} + 1)^t w_0 + \beta_{i1} \frac{1 - (\beta_{i2} + 1)^t}{\beta_{i2}}$$

Substitute β values and with some algebra we arrive at the promised closed form expression. That such an equation exists is a rarity. As such, the author believes equation (24) ought to be handled with utmost care and placed deep in a Gringots vault for safekeeping, far from the prying eyes of those nasty adversarial networks.

$$w_{it} = \left[\alpha \frac{\|\mathbf{x}_i\|^2}{N} + 1 \right]^t w_0 - \left[\frac{\mathbf{x}_i \cdot \mathbf{y}}{\|\mathbf{x}_i\|^2} \right] \left[1 - \left(\alpha \frac{\|\mathbf{x}_i\|^2}{N} + 1 \right)^t \right]$$

It is now clear to see if $\alpha \notin A$ then (24) diverges.

We have tested these results in python simulations and have found that indeed with α values above the upper bound $\alpha \leq -\frac{N}{\mathbf{x}_i \cdot \mathbf{x}_i}$, the system converges, and the opposite for $\alpha > -\frac{N}{\|\mathbf{x}_i\|^2}$.

The Dynamics of the Learning Process

Having obtained a nice analytical expression for valid α values, we would like to understand the actual learning dynamics. How is asymptotic convergence affected by the choice of α ? What is the value of the limit in equation (14)?

There are a few interesting initial observations to make.

(1) From equation (4), we can easily see that if

$$\hat{y}_j = y_j$$

then $\Delta w_{i,t} = 0$. Thus the true solution is a stable point regardless of α .

Asymptotic Behavior of Closed form and Simulated Learning: $a = -2.1N/x^2$




Figure 1: Asymptotic Convergence of Gradient Descent for Linear Regression
Least Squares Optimization_12_1

(2) Equation (20) is either monotonically increasing or decreasing. In the limit $t \rightarrow \pm\infty$, all lower order terms drop out and the rate of convergence is of the order $\mathcal{O}(\alpha^t)$.

(3) We can then write the characteristic timescale of convergence $\tau = \frac{1}{\alpha^t}$ which is exponentially small. Thus we will observe very fast convergence.

It is worthwhile as an exercise to study the dynamics of the learning system under the extremal values of α .

$\alpha > 0, \text{unstable}$

From the definition of A , if $\alpha > 0$ the system diverges exponentially.




Figure 2: Asymptotic Convergence of Gradient Descent for Linear Regression Least Squares Optimization_9_1

$\alpha = 0, \text{stable}$

In this case, the weights should diverge linearly. However, because $\beta_{i,1}$ depends on α and β_{i1} is also the constant multiple in the geometric series, the sum itself vanishes and the trajectory is stationary.




Figure 3: png

$$-\frac{N}{\|\mathbf{x}_i\|} < \alpha < 0, \text{ stable}$$




Figure 4: Asymptotic Convergence of Gradient Descent for Linear Regression
Least Squares Optimization_9_3

$$\alpha = -\frac{N}{\|\mathbf{x}_i\|^2}, \text{ stable}$$

Plugging this into (24) yields an expression

$$w_t = 0^t \left(w_0 + \frac{\mathbf{x}_i \cdot \mathbf{y}}{\|\mathbf{x}_i\|^2} \right) + \frac{\mathbf{x}_i \cdot \mathbf{y}}{\|\mathbf{x}_i\|^2}$$

This is actually interesting because the system converges in one iteration. The first term vanishes for $t > 0$, such that the closed form solution is $\frac{\mathbf{x}_i \cdot \mathbf{y}}{\|\mathbf{x}_i\|^2}$




Figure 5: png

$$\lim_{k \rightarrow -2^+} \alpha = -k \frac{N}{\|\mathbf{x}_i\|}, \text{ stable}$$

Recall from the definition of A that its left bound is open. As such, the dynamics of learning are convergent for values of α infinitesimally close to 2.




Figure 6: png

$$\alpha = -2 \frac{N}{\|\mathbf{x}_i\|^2}, \text{ *unstable*}$$

By plugging in this value of α , we get an non-converging oscillator.

$$w_t = (-1)^t w_0 + \frac{\mathbf{x}_i \cdot \mathbf{y}}{\|\mathbf{x}_i\|^2} ((-1)^t - 1)$$

By neglecting the terms with constant magnitude, we can rewrite (26) to emphasize its nature as a divergent oscillator. If we look back at our work, (26) oscillates only because gradient descent looks for the direction of descent, which is the negative of the error gradient with respect to weights.




Figure 7: png

Dynamical Bifurcations

At this point we have identified some significant α values and studied the dynamics of the system under such values. To review, we observed stationary dynamics for $\alpha = 0$, logistic growth or decay for $\alpha = -N/\|x\|^2$, and oscillatory divergence for $\alpha = -2N/\|x\|^2$. Now, notice our equations are continuous for all values of α . Thus, with equation (24) we can continuously interpolate between these the distinct dynamical regimes.

The boundary points of the set A are dynamical bifurcation points. Suppose the boundary points of A are used to dissect the real line into disjoint subsets. The qualitative behavior of learning dynamics is distinct for values of α picked from each of these subsets.

Conclusion

We have derived exact analytical bounds on α values which lead to learning convergence. Although linear regression has a closed form solution, that such results exist is exciting. Secondly, auxiliary calculations reveal exponentially small convergence timescales. It shows that understanding the learning behavior of gradient descent dynamical systems is actually quite a tractable problem. This ought to inspire efforts to understand the learning process of more complex optimization tasks. This is practically useful as with deeper understanding comes more powerful algorithms. In the longer run, it will be extremely valuable to the effort to decipher the fundamental algorithms underlying intelligent, learning, systems.