

Logic Review

Basic terms:

- ▶ Symbols: \neg (not), \wedge (and), \vee (or), \rightarrow (implies)

Logic Review

Basic terms:

- ▶ Symbols: \neg (not), \wedge (and), \vee (or), \rightarrow (implies)
- ▶ $p \rightarrow q$ – False only if p True and q False, True otherwise

Logic Review

Basic terms:

- ▶ Symbols: \neg (not), \wedge (and), \vee (or), \rightarrow (implies)
 - ▶ $p \rightarrow q$ – False only if p True and q False, True otherwise
- ▶ **contrapositive**: $p \rightarrow q \iff \neg q \rightarrow \neg p$

Logic Review

Basic terms:

- ▶ Symbols: \neg (not), \wedge (and), \vee (or), \rightarrow (implies)
 - ▶ $p \rightarrow q$ – False only if p True and q False, True otherwise
- ▶ **contrapositive**: $p \rightarrow q \iff \neg q \rightarrow \neg p$
- ▶ **negation**: distribute logical \neg over the expression

Logic Review

Basic terms:

- ▶ Symbols: \neg (not), \wedge (and), \vee (or), \rightarrow (implies)
 - ▶ $p \rightarrow q$ – False only if p True and q False, True otherwise
- ▶ **contrapositive**: $p \rightarrow q \iff \neg q \rightarrow \neg p$
- ▶ **negation**: distribute logical \neg over the expression
- ▶ **DeMorgan's Law**: $\neg(p \wedge q) = \neg p \vee \neg q$,
 $\neg(p \vee q) = \neg p \wedge \neg q$

Example 1

For which values of p , q , and r is the following expression true?
Give a succinct description (not the full truth table):

$$(\neg q \vee r) \wedge (p \rightarrow q) \wedge (\neg r \vee \neg p)$$

Example 2

Show that the following two expressions are not logically equivalent:

$$(q \rightarrow r) \vee p$$

$$q \rightarrow (r \vee p)$$

Example 3

What is the contrapositive of the statement: “If it rains, then the ground gets wet”?

Example 4

Negate the statement: “If I study and sleep well, then I will pass the exam.”

Example 5

Use DeMorgan's Law to simplify: $\neg(p \vee \neg q)$

Example 6

Which of the following are logically equivalent?

A. $\neg(p \wedge q)$

B. $\neg p \vee \neg q$

C. $\neg p \wedge \neg q$

Review: Sets

- ▶ A **set** is a collection of objects without ordering or duplicates.

Review: Sets

- ▶ A **set** is a collection of objects without ordering or duplicates.
- ▶ The **only** thing you can do with a set is see what's in it!

Review: Sets

- ▶ A **set** is a collection of objects without ordering or duplicates.
- ▶ The **only** thing you can do with a set is see what's in it!
 - ▶ $A = \{2, 4, 3, 5, 1\}$

Review: Sets

- ▶ A **set** is a collection of objects without ordering or duplicates.
- ▶ The **only** thing you can do with a set is see what's in it!
 - ▶ $A = \{2, 4, 3, 5, 1\}$
 - ▶ Is 2 in the set A ?

Review: Sets

- ▶ A **set** is a collection of objects without ordering or duplicates.
- ▶ The **only** thing you can do with a set is see what's in it!
 - ▶ $A = \{2, 4, 3, 5, 1\}$
 - ▶ Is 2 in the set A ? ✓

Review: Sets

- ▶ A **set** is a collection of objects without ordering or duplicates.
- ▶ The **only** thing you can do with a set is see what's in it!
 - ▶ $A = \{2, 4, 3, 5, 1\}$
 - ▶ Is 2 in the set A ? ✓
 - ▶ Is 0 in the set A ?

Review: Sets

- ▶ A **set** is a collection of objects without ordering or duplicates.
- ▶ The **only** thing you can do with a set is see what's in it!
 - ▶ $A = \{2, 4, 3, 5, 1\}$
 - ▶ Is 2 in the set A ? ✓
 - ▶ Is 0 in the set A ? ✗

Review: Sets

- ▶ A **set** is a collection of objects without ordering or duplicates.
- ▶ The **only** thing you can do with a set is see what's in it!
 - ▶ $A = \{2, 4, 3, 5, 1\}$
 - ▶ Is 2 in the set A ? ✓
 - ▶ Is 0 in the set A ? ✗
 - ▶ What is the first element of A ?

Review: Sets

- ▶ A **set** is a collection of objects without ordering or duplicates.
- ▶ The **only** thing you can do with a set is see what's in it!
 - ▶ $A = \{2, 4, 3, 5, 1\}$
 - ▶ Is 2 in the set A ? ✓
 - ▶ Is 0 in the set A ? ✗
 - ▶ What is the first element of A ? Invalid question!

Review: Sets

- ▶ A **set** is a collection of objects without ordering or duplicates.
- ▶ The **only** thing you can do with a set is see what's in it!
 - ▶ $A = \{2, 4, 3, 5, 1\}$
 - ▶ Is 2 in the set A ? ✓
 - ▶ Is 0 in the set A ? ✗
 - ▶ What is the first element of A ? **Invalid question!**
- ▶ 3 standard ways to define sets:

Review: Sets

- ▶ A **set** is a collection of objects without ordering or duplicates.
- ▶ The **only** thing you can do with a set is see what's in it!
 - ▶ $A = \{2, 4, 3, 5, 1\}$
 - ▶ Is 2 in the set A ? ✓
 - ▶ Is 0 in the set A ? ✗
 - ▶ What is the first element of A ? **Invalid question!**
- ▶ 3 standard ways to define sets:
 1. *Precisely* describe what they contain in English: Let S denote the set of all positive even integers.

Review: Sets

- ▶ A **set** is a collection of objects without ordering or duplicates.
- ▶ The **only** thing you can do with a set is see what's in it!
 - ▶ $A = \{2, 4, 3, 5, 1\}$
 - ▶ Is 2 in the set A ? ✓
 - ▶ Is 0 in the set A ? ✗
 - ▶ What is the first element of A ? **Invalid question!**
- ▶ 3 standard ways to define sets:
 1. *Precisely* describe what they contain in English: Let S denote the set of all positive even integers.
 2. List out their elements (with ellipses if necessary and clear): $S = \{2, 4, 6, \dots\}$.

Review: Sets

- ▶ A **set** is a collection of objects without ordering or duplicates.
- ▶ The **only** thing you can do with a set is see what's in it!
 - ▶ $A = \{2, 4, 3, 5, 1\}$
 - ▶ Is 2 in the set A ? ✓
 - ▶ Is 0 in the set A ? ✗
 - ▶ What is the first element of A ? **Invalid question!**
- ▶ 3 standard ways to define sets:
 1. *Precisely* describe what they contain in English: Let S denote the set of all positive even integers.
 2. List out their elements (with ellipses if necessary and clear):
 $S = \{2, 4, 6, \dots\}$.
 3. Set builder notation: $S = \{2k \mid \underbrace{k \in \mathbb{Z}, k > 0}_{\text{Condition}}\}$

Review: Sets

- ▶ A **set** is a collection of objects without ordering or duplicates.
- ▶ The **only** thing you can do with a set is see what's in it!
 - ▶ $A = \{2, 4, 3, 5, 1\}$
 - ▶ Is 2 in the set A ? ✓
 - ▶ Is 0 in the set A ? ✗
 - ▶ What is the first element of A ? **Invalid question!**
- ▶ 3 standard ways to define sets:
 1. *Precisely* describe what they contain in English: Let S denote the set of all positive even integers.
 2. List out their elements (with ellipses if necessary and clear): $S = \{2, 4, 6, \dots\}$.
 3. Set builder notation: $S = \{2k \mid \underbrace{k \in \mathbb{Z}, k > 0}_{\text{Condition}}\}$
- ▶ Special notation: $\emptyset = \{\}$, the empty set.

Review: Sets

- ▶ $(A \subseteq B)$ A is a subset of B : $\forall a \in A, a \in B$
 $\{1, 2, 4\} \subseteq \{1, 2, 3, 4\}$

Review: Sets

- ▶ $(A \subseteq B)$ A is a subset of B : $\forall a \in A, a \in B$
 $\{1, 2, 4\} \subseteq \{1, 2, 3, 4\}$
- ▶ Equality: $A = B \iff A \subseteq B$ and $B \subseteq A$

Review: Sets

- ▶ $(A \subseteq B)$ A is a subset of B : $\forall a \in A, a \in B$
 $\{1, 2, 4\} \subseteq \{1, 2, 3, 4\}$
- ▶ Equality: $A = B \iff A \subseteq B$ and $B \subseteq A$
- ▶ Union: $A \cup B = \{x \mid x \in A \text{ or } x \in B\}$
 $\{\text{milk, cheese, eggs}\} \cup \{\text{yogurt, milk, carrots}\} =$
 $\{\text{milk, yogurt, cheese, eggs, carrots}\}$

Review: Sets

- ▶ $(A \subseteq B)$ A is a subset of B : $\forall a \in A, a \in B$
 $\{1, 2, 4\} \subseteq \{1, 2, 3, 4\}$
- ▶ Equality: $A = B \iff A \subseteq B$ and $B \subseteq A$
- ▶ Union: $A \cup B = \{x \mid x \in A \text{ or } x \in B\}$
 $\{\text{milk, cheese, eggs}\} \cup \{\text{yogurt, milk, carrots}\} =$
 $\{\text{milk, yogurt, cheese, eggs, carrots}\}$
- ▶ Intersection: $A \cap B = \{x \mid x \in A \text{ and } x \in B\}$
 $\{\text{milk, cheese, eggs}\} \cap \{\text{yogurt, milk, carrots}\} = \{\text{milk}\}$

Review: Sets

- ▶ $(A \subseteq B)$ A is a subset of B : $\forall a \in A, a \in B$
 $\{1, 2, 4\} \subseteq \{1, 2, 3, 4\}$
- ▶ Equality: $A = B \iff A \subseteq B$ and $B \subseteq A$
- ▶ Union: $A \cup B = \{x \mid x \in A \text{ or } x \in B\}$
 $\{\text{milk, cheese, eggs}\} \cup \{\text{yogurt, milk, carrots}\} =$
 $\{\text{milk, yogurt, cheese, eggs, carrots}\}$
- ▶ Intersection: $A \cap B = \{x \mid x \in A \text{ and } x \in B\}$
 $\{\text{milk, cheese, eggs}\} \cap \{\text{yogurt, milk, carrots}\} = \{\text{milk}\}$
 - ▶ What can you say about $A, A \cup B$, and $A \cap B$?

Review: Sets

- ▶ $(A \subseteq B)$ A is a subset of B : $\forall a \in A, a \in B$
 $\{1, 2, 4\} \subseteq \{1, 2, 3, 4\}$
- ▶ Equality: $A = B \iff A \subseteq B$ and $B \subseteq A$
- ▶ Union: $A \cup B = \{x \mid x \in A \text{ or } x \in B\}$
 $\{\text{milk, cheese, eggs}\} \cup \{\text{yogurt, milk, carrots}\} =$
 $\{\text{milk, yogurt, cheese, eggs, carrots}\}$
- ▶ Intersection: $A \cap B = \{x \mid x \in A \text{ and } x \in B\}$
 $\{\text{milk, cheese, eggs}\} \cap \{\text{yogurt, milk, carrots}\} = \{\text{milk}\}$
 - ▶ What can you say about $A, A \cup B$, and $A \cap B$?
 - ▶ $A \cap B \subseteq A \subseteq A \cup B$

Review: Sets

- ▶ $(A \subseteq B)$ A is a subset of B : $\forall a \in A, a \in B$
 $\{1, 2, 4\} \subseteq \{1, 2, 3, 4\}$
- ▶ Equality: $A = B \iff A \subseteq B$ and $B \subseteq A$
- ▶ Union: $A \cup B = \{x \mid x \in A \text{ or } x \in B\}$
 $\{\text{milk, cheese, eggs}\} \cup \{\text{yogurt, milk, carrots}\} =$
 $\{\text{milk, yogurt, cheese, eggs, carrots}\}$
- ▶ Intersection: $A \cap B = \{x \mid x \in A \text{ and } x \in B\}$
 $\{\text{milk, cheese, eggs}\} \cap \{\text{yogurt, milk, carrots}\} = \{\text{milk}\}$
 - ▶ What can you say about $A, A \cup B$, and $A \cap B$?
 - ▶ $A \cap B \subseteq A \subseteq A \cup B$
- ▶ Cardinality: Given a (finite) set A , $|A|$ is the number of elements in the set.

Review: Sets

- ▶ $(A \subseteq B)$ A is a subset of B : $\forall a \in A, a \in B$
 $\{1, 2, 4\} \subseteq \{1, 2, 3, 4\}$
- ▶ Equality: $A = B \iff A \subseteq B$ and $B \subseteq A$
- ▶ Union: $A \cup B = \{x \mid x \in A \text{ or } x \in B\}$
 $\{\text{milk, cheese, eggs}\} \cup \{\text{yogurt, milk, carrots}\} =$
 $\{\text{milk, yogurt, cheese, eggs, carrots}\}$
- ▶ Intersection: $A \cap B = \{x \mid x \in A \text{ and } x \in B\}$
 $\{\text{milk, cheese, eggs}\} \cap \{\text{yogurt, milk, carrots}\} = \{\text{milk}\}$
 - ▶ What can you say about $A, A \cup B$, and $A \cap B$?
 - ▶ $A \cap B \subseteq A \subseteq A \cup B$
- ▶ Cardinality: Given a (finite) set A , $|A|$ is the number of elements in the set.
- ▶ Cartesian Product: $A \times B = \{(a, b) \mid a \in A, b \in B\}$

Review: Sets

- ▶ $(A \subseteq B)$ A is a subset of B : $\forall a \in A, a \in B$
 $\{1, 2, 4\} \subseteq \{1, 2, 3, 4\}$
- ▶ Equality: $A = B \iff A \subseteq B$ and $B \subseteq A$
- ▶ Union: $A \cup B = \{x \mid x \in A \text{ or } x \in B\}$
 $\{\text{milk, cheese, eggs}\} \cup \{\text{yogurt, milk, carrots}\} =$
 $\{\text{milk, yogurt, cheese, eggs, carrots}\}$
- ▶ Intersection: $A \cap B = \{x \mid x \in A \text{ and } x \in B\}$
 $\{\text{milk, cheese, eggs}\} \cap \{\text{yogurt, milk, carrots}\} = \{\text{milk}\}$
 - ▶ What can you say about $A, A \cup B$, and $A \cap B$?
 - ▶ $A \cap B \subseteq A \subseteq A \cup B$
- ▶ Cardinality: Given a (finite) set A , $|A|$ is the number of elements in the set.
- ▶ Cartesian Product: $A \times B = \{(a, b) \mid a \in A, b \in B\}$
- ▶ Powerset (set of all subsets): $\mathbb{P}(A) = \{S \mid S \subseteq A\}$

Important Points

- Sets can contain sets! The set $A = \{1, 2, 3\}$ is distinct from the set $B = \{A\} = \{\{1, 2, 3\}\}$

Important Points

- ▶ Sets can contain sets! The set $A = \{1, 2, 3\}$ is distinct from the set $B = \{A\} = \{\{1, 2, 3\}\}$
- ▶ Sets need not have the same “type” of element: $\{1, (3, \text{“no”}), f\}$ is a perfectly valid set.

Practice Problems I

1. What is $|A|$, where $A = \{1, 2, 4, 1, 5\}$?

Practice Problems I

1. What is $|A|$, where $A = \{1, 2, 4, 1, 5\}$?
2. What is $\emptyset \times \{1, 2, 3\}$?

Practice Problems I

1. What is $|A|$, where $A = \{1, 2, 4, 1, 5\}$?
2. What is $\emptyset \times \{1, 2, 3\}$?
3. What is $\{\emptyset\} \times \{1, 2, 3\}$?

Practice Problems I

1. What is $|A|$, where $A = \{1, 2, 4, 1, 5\}$?
2. What is $\emptyset \times \{1, 2, 3\}$?
3. What is $\{\emptyset\} \times \{1, 2, 3\}$?
4. For sets A and B , what is $|A \times B|$, in terms of $|A|$ and $|B|$?

Practice Problems I

1. What is $|A|$, where $A = \{1, 2, 4, 1, 5\}$?
2. What is $\emptyset \times \{1, 2, 3\}$?
3. What is $\{\emptyset\} \times \{1, 2, 3\}$?
4. For sets A and B , what is $|A \times B|$, in terms of $|A|$ and $|B|$?
5. What is $|A \cup B|$?

Practice Problems I

1. What is $|A|$, where $A = \{1, 2, 4, 1, 5\}$?
2. What is $\emptyset \times \{1, 2, 3\}$?
3. What is $\{\emptyset\} \times \{1, 2, 3\}$?
4. For sets A and B , what is $|A \times B|$, in terms of $|A|$ and $|B|$?
5. What is $|A \cup B|$?
6. What is $|\mathbb{P}(A)|$?

Practice Problems I

1. What is $|A|$, where $A = \{1, 2, 4, 1, 5\}$?
2. What is $\emptyset \times \{1, 2, 3\}$?
3. What is $\{\emptyset\} \times \{1, 2, 3\}$?
4. For sets A and B , what is $|A \times B|$, in terms of $|A|$ and $|B|$?
5. What is $|A \cup B|$?
6. What is $|\mathbb{P}(A)|$?
7. For any set A , what element is *always* in $\mathbb{P}(A)$?

Practice Problems I

1. What is $|A|$, where $A = \{1, 2, 4, 1, 5\}$?
2. What is $\emptyset \times \{1, 2, 3\}$?
3. What is $\{\emptyset\} \times \{1, 2, 3\}$?
4. For sets A and B , what is $|A \times B|$, in terms of $|A|$ and $|B|$?
5. What is $|A \cup B|$?
6. What is $|\mathbb{P}(A)|$?
7. For any set A , what element is *always* in $\mathbb{P}(A)$?
8. What is $\mathbb{P}(\{1, 2, 3\})$?

Set Proofs

Proving Inclusion

Let $a \in A$.

(logic...)

Thus, $a \in B$, so $A \subseteq B$.

Set Proofs

Proving Inclusion

Let $a \in A$.

(logic...)

Thus, $a \in B$, so $A \subseteq B$.

How would you prove that $A \not\subseteq B$? How about $A \subsetneq B$?

Practice Problems II

1. Prove that $\{6x \mid x \in \mathbb{N}\} \subseteq \{2y \mid y \in \mathbb{N}\}$

Practice Problems II

1. Prove that $\{6x \mid x \in \mathbb{N}\} \subseteq \{2y \mid y \in \mathbb{N}\}$
2. Prove that $\{p > 2 \mid p \text{ prime}\} \subseteq \{x : 2 \nmid x \in \mathbb{N}\}$

Practice Problems II

1. Prove that $\{6x \mid x \in \mathbb{N}\} \subseteq \{2y \mid y \in \mathbb{N}\}$
2. Prove that $\{p > 2 \mid p \text{ prime}\} \subseteq \{x : 2 \nmid x \in \mathbb{N}\}$
3. Prove that $\{x^2 \mid x \in \mathbb{R}\} = \{y \geq 0 \mid y \in \mathbb{R}\}$

Counting Elements in Sets

- ▶ Number of ways to order n objects: $n!$

Counting Elements in Sets

- ▶ Number of ways to order n objects: $n!$
- ▶ Number of ways to order k objects from a set of n : $\frac{n!}{(n-k)!}$

Counting Elements in Sets

- ▶ Number of ways to order n objects: $n!$
- ▶ Number of ways to order k objects from a set of n : $\frac{n!}{(n-k)!}$
- ▶ (!!!) Number of ways to choose k objects from a collection of n : $\binom{n}{k} = \frac{n!}{k!(n-k)!}$

Counting Elements in Sets

- ▶ Number of ways to order n objects: $n!$
- ▶ Number of ways to order k objects from a set of n : $\frac{n!}{(n-k)!}$
- ▶ (!!!) Number of ways to choose k objects from a collection of n : $\binom{n}{k} = \frac{n!}{k!(n-k)!}$
- ▶ Stars and Bars: How many ways to partition n identical elements into k bins: $\binom{k+n-1}{k-1}$

Practice Problems III

1. In a race with 6 contestants, how many possible top 3 players are there (order matters)?

Practice Problems III

1. In a race with 6 contestants, how many possible top 3 players are there (order matters)?
2. How many ways are there to divide $2n$ students into two even groups?

Practice Problems III

1. In a race with 6 contestants, how many possible top 3 players are there (order matters)?
2. How many ways are there to divide $2n$ students into two even groups?
3. How many 10-bit binary strings are there with exactly 3 0 bits?

Practice Problems III

1. In a race with 6 contestants, how many possible top 3 players are there (order matters)?
2. How many ways are there to divide $2n$ students into two even groups?
3. How many 10-bit binary strings are there with exactly 3 0 bits?
4. How many triples of nonnegative numbers (x, y, z) are there such that $x + y + z = 10$?

Practice Problems III

1. In a race with 6 contestants, how many possible top 3 players are there (order matters)?
2. How many ways are there to divide $2n$ students into two even groups?
3. How many 10-bit binary strings are there with exactly 3 0 bits?
4. How many triples of nonnegative numbers (x, y, z) are there such that $x + y + z = 10$?
5. Same question but for *positive* numbers.

Practice Problems III

1. In a race with 6 contestants, how many possible top 3 players are there (order matters)?
2. How many ways are there to divide $2n$ students into two even groups?
3. How many 10-bit binary strings are there with exactly 3 0 bits?
4. How many triples of nonnegative numbers (x, y, z) are there such that $x + y + z = 10$?
5. Same question but for *positive* numbers.
6. (Summer 2024 Review Question) Let $A = \{(a, b) \in \mathbb{R}^2 \mid a = 3 - b^2\}$, $B = \{(x, y) \in \mathbb{R}^2 \mid |x| \geq 1 \text{ or } |y| \geq 1\}$. Prove that $A \subseteq B$.

Questions/Examples

Functions: Review

- Functions map between sets. Given an input, it gives exactly one output (duh).

Functions: Review

- ▶ Functions map between sets. Given an input, it gives exactly one output (duh).
- ▶ Functions are defined *both* by what they do to the input and their (co)domain.

Functions: Review

- ▶ Functions map between sets. Given an input, it gives exactly one output (duh).
- ▶ Functions are defined *both* by what they do to the input and their (co)domain.
- ▶ $f : \mathbb{R} \rightarrow \mathbb{R}, f(x) = x^2$ and $g : \mathbb{R} \rightarrow [0, \infty), g(x) = x^2$ are *different* functions (only one is onto), even though they have the same formula.

Functions: Review

- ▶ Functions map between sets. Given an input, it gives exactly one output (duh).
- ▶ Functions are defined *both* by what they do to the input and their (co)domain.
- ▶ $f : \mathbb{R} \rightarrow \mathbb{R}, f(x) = x^2$ and $g : \mathbb{R} \rightarrow [0, \infty), g(x) = x^2$ are *different* functions (only one is onto), even though they have the same formula.
- ▶ Can be specified with a precise English description, a formula (if working with numbers), or by defining each input/output pair individually.

Functions: Review

- ▶ Functions map between sets. Given an input, it gives exactly one output (duh).
- ▶ Functions are defined *both* by what they do to the input and their (co)domain.
- ▶ $f : \mathbb{R} \rightarrow \mathbb{R}, f(x) = x^2$ and $g : \mathbb{R} \rightarrow [0, \infty), g(x) = x^2$ are *different* functions (only one is onto), even though they have the same formula.
- ▶ Can be specified with a precise English description, a formula (if working with numbers), or by defining each input/output pair individually.
- ▶ These are all the same function:

Functions: Review

- ▶ Functions map between sets. Given an input, it gives exactly one output (duh).
- ▶ Functions are defined *both* by what they do to the input and their (co)domain.
- ▶ $f : \mathbb{R} \rightarrow \mathbb{R}, f(x) = x^2$ and $g : \mathbb{R} \rightarrow [0, \infty), g(x) = x^2$ are *different* functions (only one is onto), even though they have the same formula.
- ▶ Can be specified with a precise English description, a formula (if working with numbers), or by defining each input/output pair individually.
- ▶ These are all the same function:
 - ▶ $f : \{2, 3, 4\} \rightarrow \mathbb{N}$ $f(n)$ returns the n^{th} prime number

Functions: Review

- ▶ Functions map between sets. Given an input, it gives exactly one output (duh).
- ▶ Functions are defined *both* by what they do to the input and their (co)domain.
- ▶ $f : \mathbb{R} \rightarrow \mathbb{R}, f(x) = x^2$ and $g : \mathbb{R} \rightarrow [0, \infty), g(x) = x^2$ are *different* functions (only one is onto), even though they have the same formula.
- ▶ Can be specified with a precise English description, a formula (if working with numbers), or by defining each input/output pair individually.
- ▶ These are all the same function:
 - ▶ $f : \{2, 3, 4\} \rightarrow \mathbb{N}$ $f(n)$ returns the n^{th} prime number
 - ▶ $f : \mathbb{N} \cap [2, 4] \rightarrow \mathbb{N}$ $f(2) = 3, f(3) = 5, f(4) = 7$

Functions: Review

- ▶ Functions map between sets. Given an input, it gives exactly one output (duh).
- ▶ Functions are defined *both* by what they do to the input and their (co)domain.
- ▶ $f : \mathbb{R} \rightarrow \mathbb{R}, f(x) = x^2$ and $g : \mathbb{R} \rightarrow [0, \infty), g(x) = x^2$ are *different* functions (only one is onto), even though they have the same formula.
- ▶ Can be specified with a precise English description, a formula (if working with numbers), or by defining each input/output pair individually.
- ▶ These are all the same function:
 - ▶ $f : \{2, 3, 4\} \rightarrow \mathbb{N}$ $f(n)$ returns the n^{th} prime number
 - ▶ $f : \mathbb{N} \cap [2, 4] \rightarrow \mathbb{N}$ $f(2) = 3, f(3) = 5, f(4) = 7$
 - ▶ $f : \{2, 3, 4\} \rightarrow \mathbb{N}$ $f(x) = 2x - 1$

Onto, One-to-one and Bijective

- ▶ A function $f : A \rightarrow B$ is **onto** if for every $b \in B$, there is some $a \in A$ where $f(a) = b$.

Onto, One-to-one and Bijective

- ▶ A function $f : A \rightarrow B$ is **onto** if for every $b \in B$, there is some $a \in A$ where $f(a) = b$.
- ▶ A function $f : A \rightarrow B$ is **one-to-one** if $f(x) = f(y)$ implies $x = y$ for all $x, y \in A$.

Onto, One-to-one and Bijective

- ▶ A function $f : A \rightarrow B$ is **onto** if for every $b \in B$, there is some $a \in A$ where $f(a) = b$.
- ▶ A function $f : A \rightarrow B$ is **one-to-one** if $f(x) = f(y)$ implies $x = y$ for all $x, y \in A$.
- ▶ A nice way to think about it: onto means every $b \in B$ has at *least* one preimage; one-to-one means every $b \in B$ has at *most* one preimage.

Onto, One-to-one and Bijective

- ▶ A function $f : A \rightarrow B$ is **onto** if for every $b \in B$, there is some $a \in A$ where $f(a) = b$.
- ▶ A function $f : A \rightarrow B$ is **one-to-one** if $f(x) = f(y)$ implies $x = y$ for all $x, y \in A$.
- ▶ A nice way to think about it: onto means every $b \in B$ has at *least* one preimage; one-to-one means every $b \in B$ has at *most* one preimage.
- ▶ So f being **bijective** (onto and one-to-one) means every $b \in B$ has *exactly* one preimage.

Practice Questions I

Which of these functions are onto, one-to-one, or both?

Signature	$f(x)$	Onto?	One-to-one?
$\mathbb{R} \rightarrow \mathbb{R}$	x^2		
$[0, \infty) \rightarrow [0, \infty)$	x^2		
$\mathbb{N} \rightarrow \mathbb{Z}$	$(-1)^x x$		
$\{2, 5, 6\} \rightarrow \{3, 6, 7, 8\}$	$x + 1$		
$\mathbb{N} \cup \{-1\} \rightarrow \mathbb{N}$	$2x + 1$		

Practice Questions II

1. Let $g : \mathbb{N} \rightarrow \mathbb{N}$ be onto, and define $f : \mathbb{N}^2 \rightarrow \mathbb{Z}$ by $f(n, m) = (m - 1)g(n)$. Prove f is onto.

Practice Questions II

1. Let $g : \mathbb{N} \rightarrow \mathbb{N}$ be onto, and define $f : \mathbb{N}^2 \rightarrow \mathbb{Z}$ by $f(n, m) = (m - 1)g(n)$. Prove f is onto.
2. (SU24 Review) Let $f : \mathbb{Z}^+ \times \mathbb{Z}^+ \rightarrow \mathbb{Q}^2$, $f(x, y) = (\frac{x}{y}, x + y)$. Show that f is one-to-one.

Practice Questions II

1. Let $g : \mathbb{N} \rightarrow \mathbb{N}$ be onto, and define $f : \mathbb{N}^2 \rightarrow \mathbb{Z}$ by $f(n, m) = (m - 1)g(n)$. Prove f is onto.
2. (SU24 Review) Let $f : \mathbb{Z}^+ \times \mathbb{Z}^+ \rightarrow \mathbb{Q}^2$, $f(x, y) = (\frac{x}{y}, x + y)$. Show that f is one-to-one.
3. Suppose we have finite sets A and B with $|A| = |B| = n \in \mathbb{N}$. Show that if $f : A \rightarrow B$ is one-to-one, then f is onto. How many such functions are there, for fixed A and B ?

Practice Questions II

1. Let $g : \mathbb{N} \rightarrow \mathbb{N}$ be onto, and define $f : \mathbb{N}^2 \rightarrow \mathbb{Z}$ by $f(n, m) = (m - 1)g(n)$. Prove f is onto.
2. (SU24 Review) Let $f : \mathbb{Z}^+ \times \mathbb{Z}^+ \rightarrow \mathbb{Q}^2$, $f(x, y) = (\frac{x}{y}, x + y)$. Show that f is one-to-one.
3. Suppose we have finite sets A and B with $|A| = |B| = n \in \mathbb{N}$. Show that if $f : A \rightarrow B$ is one-to-one, then f is onto. How many such functions are there, for fixed A and B ?
4. Suppose $A \subseteq \mathbb{R}$ and $f : A \rightarrow A$, $f(x) = \sqrt{2}x$. For what sets A is f a bijection?

Practice Questions II

1. Let $g : \mathbb{N} \rightarrow \mathbb{N}$ be onto, and define $f : \mathbb{N}^2 \rightarrow \mathbb{Z}$ by $f(n, m) = (m - 1)g(n)$. Prove f is onto.
2. (SU24 Review) Let $f : \mathbb{Z}^+ \times \mathbb{Z}^+ \rightarrow \mathbb{Q}^2$, $f(x, y) = (\frac{x}{y}, x + y)$. Show that f is one-to-one.
3. Suppose we have finite sets A and B with $|A| = |B| = n \in \mathbb{N}$. Show that if $f : A \rightarrow B$ is one-to-one, then f is onto. How many such functions are there, for fixed A and B ?
4. Suppose $A \subseteq \mathbb{R}$ and $f : A \rightarrow A$, $f(x) = \sqrt{2}x$. For what sets A is f a bijection?
5. Suppose that $f \circ g$ is one-to-one. Does f have to be one-to-one? Does g ?

Questions/Examples

Questions/Examples

Questions/Examples

Questions/Examples

Recurrences: Review

- We can specify functions with a recursive formula:

$$T(n) = \begin{cases} c & n \leq B \\ \langle \text{Formula with smaller arguments to } T \rangle & n > B \end{cases}$$

Recurrences: Review

- We can specify functions with a recursive formula:

$$T(n) = \begin{cases} c & n \leq B \\ \langle \text{Formula with smaller arguments to } T \rangle & n > B \end{cases}$$

- Some examples:

Recurrence		Closed Formula
$T(n) = \begin{cases} 1 & n \leq 1 \\ 2T(n-1) & n > 1 \end{cases}$		$T(n) = 2^n$
$T(n) = \begin{cases} 1 & n \leq 1 \\ 2T(n/2) + 1 & n > 1 \end{cases}$		$T(n) = 2n - 1$
$T(n) = \begin{cases} 0 & n \leq 1 \\ 2T(n/2) + n & n > 1 \end{cases}$		$T(n) = n \log(n)$

Solving A Recurrence

The general strategy for solving a recurrence is a **recurrence tree**.

- ▶ Each node has a value equal to the **nonrecursive** work done.

Solving A Recurrence

The general strategy for solving a recurrence is a **recurrence tree**.

- ▶ Each node has a value equal to the **nonrecursive** work done.
- ▶ Each node has a child for every recursive call it makes.

Solving A Recurrence


The general strategy for solving a recurrence is a **recurrence tree**.

- ▶ Each node has a value equal to the **nonrecursive** work done.
- ▶ Each node has a child for every recursive call it makes.

Solving A Recurrence

The general strategy for solving a recurrence is a **recurrence tree**.

- ▶ Each node has a value equal to the **nonrecursive** work done.
- ▶ Each node has a child for every recursive call it makes.



Solving A Recurrence

1. At level k ...

Solving A Recurrence

1. At level k ...
 - ▶ How many nodes are there?

Solving A Recurrence

1. At level k ...
 - ▶ How many nodes are there?
 - ▶ What is the input?

Solving A Recurrence

1. At level k ...

- ▶ How many nodes are there?
- ▶ What is the input?
- ▶ How much work is done by each node?

Solving A Recurrence

1. At level k ...

- ▶ How many nodes are there?
- ▶ What is the input?
- ▶ How much work is done by each node?
- ▶ What is the total nonrecursive work? ($W(k)$)

Solving A Recurrence

1. At level k ...
 - ▶ How many nodes are there?
 - ▶ What is the input?
 - ▶ How much work is done by each node?
 - ▶ What is the total nonrecursive work? ($W(k)$)
2. What is the height of the tree (i.e. at what level h do we reach the base case input)?

Solving A Recurrence

1. At level k ...
 - ▶ How many nodes are there?
 - ▶ What is the input?
 - ▶ How much work is done by each node?
 - ▶ What is the total nonrecursive work? ($W(k)$)
2. What is the height of the tree (i.e. at what level h do we reach the base case input)?
3. What is the work done by the leaf nodes?
 $L = [\# \text{ of leaf nodes}] \times [\text{work done by each}]$

Solving A Recurrence

1. At level k ...
 - ▶ How many nodes are there?
 - ▶ What is the input?
 - ▶ How much work is done by each node?
 - ▶ What is the total nonrecursive work? ($W(k)$)
2. What is the height of the tree (i.e. at what level h do we reach the base case input)?
3. What is the work done by the leaf nodes?
 $L = [\# \text{ of leaf nodes}] \times [\text{work done by each}]$

4. The total work is $\underbrace{\sum_{k=0}^{h-1} W(k)}_{\text{internal work}} + \underbrace{L}_{\text{leaf work}}$

Recurrence Practice I

$$T(n) = \begin{cases} c & n \leq 1 \\ 3T(n/2) + n & n > 1 \end{cases}$$

What is $T(n)$ for n a power of 2?

Recurrence Practice I

$$T(n) = \begin{cases} c & n \leq 1 \\ 3T(n/2) + n & n > 1 \end{cases}$$

What is $T(n)$ for n a power of 2?

1. At level k ...

- ▶ There are _____ nodes.
- ▶ The input is _____.
- ▶ Each node does _____ work.
- ▶ The total nonrecursive work $W(k) =$ _____

2. $h =$ _____

3. What is the work done by the leaf nodes?

$$L = [\# \text{ of leaf nodes}] \times [\text{work done by each}] = \underline{\hspace{2cm}}$$

4.
$$\underbrace{\sum_{k=0}^{h-1} W(k)}_{\text{internal work}} + \underbrace{L}_{\text{leaf work}} = \underline{\hspace{2cm}}$$

Recurrence Practice I

$$T(n) = \begin{cases} c & n \leq 1 \\ 3T(n/2) + n & n > 1 \end{cases}$$

What is $T(n)$ for n a power of 2?

1. At level k ...

- ▶ There are _____ nodes.
- ▶ The input is _____.
- ▶ Each node does _____ work.
- ▶ The total nonrecursive work $W(k) =$ _____

2. $h =$ _____

3. What is the work done by the leaf nodes?

$$L = [\# \text{ of leaf nodes}] \times [\text{work done by each}] = \underline{\hspace{2cm}}$$

4. $\underbrace{\sum_{k=0}^{h-1} W(k)}_{\text{internal work}} + \underbrace{L}_{\text{leaf work}} = \underline{\hspace{2cm}}$

Answer: $T(n) = (c + 2)n^{\log_2 3} - 2n$

Recurrence Practice II

$$T(n) = \begin{cases} c & n \leq 2 \\ 2T(n-1) + d & n > 2 \end{cases}$$

What is $T(n)$ for $n \geq 2$?

Recurrence Practice II

$$T(n) = \begin{cases} c & n \leq 2 \\ 2T(n-1) + d & n > 2 \end{cases}$$

What is $T(n)$ for $n \geq 2$?

Answer: $T(n) = (c + d)2^{n-2} - d$

Recurrence Practice III

$$T(n) = \begin{cases} c & n \leq 1 \\ T(n/2) + n^2 & n > 1 \end{cases}$$

What is $T(n)$ for n a power of 2?

Recurrence Practice III

$$T(n) = \begin{cases} c & n \leq 1 \\ T(n/2) + n^2 & n > 1 \end{cases}$$

What is $T(n)$ for n a power of 2?

Answer: $T(n) = \frac{4}{3}(n^2 - 1) + c$

What about Θ ?

- ▶ If all we care about is what T is Θ of, the specific constants don't matter.

What about Θ ?

- ▶ If all we care about is what T is Θ of, the specific constants don't matter.
- ▶ In the previous slide, $T(n) = \frac{4}{3}(n^2 - 1) + c$ is always $\Theta(n^2)$, no matter what c is.

What about Θ ?

- ▶ If all we care about is what T is Θ of, the specific constants don't matter.
- ▶ In the previous slide, $T(n) = \frac{4}{3}(n^2 - 1) + c$ is always $\Theta(n^2)$, no matter what c is.
- ▶ Similar for $T(n) = (c + d)2^{n-2} - d$ is $\Theta(2^n)$ and $T(n) = (c + 2)n^{\log_2 3} - 2n$ is $\Theta(n^{\log_2 3})$.

Recurrence Practice V

What is the Big-Theta running time of SHAKE?

```
SHAKE( $A[1..n]$ )  
  if  $n \leq 1$ :  
    return  $A$   
   $m \leftarrow \lfloor n/2 \rfloor$   
   $A_1 \leftarrow \text{SHAKE}(A[1..m])$   
   $A_2 \leftarrow \text{SHAKE}(A[m+1..n])$   
   $B \leftarrow []$   
  for  $i$  in  $1..m$ :  
     $B.\text{add}(A_1[i])$   
     $B.\text{add}(A_2[i])$   
  return  $B$ 
```

Recurrence Practice V

What is the Big-Theta running time of SHAKE?

```
SHAKE( $A[1..n]$ )  
  if  $n \leq 1$ :  
    return  $A$   
   $m \leftarrow \lfloor n/2 \rfloor$   
   $A_1 \leftarrow \text{SHAKE}(A[1..m])$   
   $A_2 \leftarrow \text{SHAKE}(A[m + 1..n])$   
   $B \leftarrow []$   
  for  $i$  in  $1..m$ :  
     $B.\text{add}(A_1[i])$   
     $B.\text{add}(A_2[i])$   
  return  $B$ 
```

Answer: $\Theta(n \log n)$

Recurrence Practice VI: HARD

$$T(n) = \begin{cases} c & n \leq 1 \\ T(n/2) + T(n/3) + T(n/6) + n & n > 1 \end{cases}$$

What is $T(n)$ Big-Theta of?

Recurrence Practice VI: HARD

$$T(n) = \begin{cases} c & n \leq 1 \\ T(n/2) + T(n/3) + T(n/6) + n & n > 1 \end{cases}$$

What is $T(n)$ Big-Theta of?

Answer: $\Theta(n \log n)$

Questions/Examples

Questions/Examples

Questions/Examples

Questions/Examples

Induction!!!

- ▶ Fundamental course concept; closely related to recursive definition in programming languages

Induction!!!

- ▶ Fundamental course concept; closely related to recursive definition in programming languages
- ▶ **Goal:** Prove a mathematical statement is true for all natural numbers.

Induction!!!

- ▶ Fundamental course concept; closely related to recursive definition in programming languages
- ▶ **Goal:** Prove a mathematical statement is true for all natural numbers.
- ▶ Outline: base case, inductive hypothesis, inductive step

Induction Proofs

The general strategy for proving a claim by induction is to

- (a) define the **base case(s)** and show the claim holds for them
- (b) state the **inductive hypothesis** assuming that the claim holds true for all $n < k$ ($n \in \mathbb{N}$)
- (c) prove that the claim holds for $n = k$ in the rest of the **inductive step**

Induction Proofs

The general strategy for proving a claim by induction is to

- (a) define the **base case(s)** and show the claim holds for them
 - (b) state the **inductive hypothesis** assuming that the claim holds true for all $n < k$ ($n \in \mathbb{N}$)
 - (c) prove that the claim holds for $n = k$ in the rest of the **inductive step**
- You can also do $n \leq k$ and then $n = k + 1$.

Induction Proofs

The general strategy for proving a claim by induction is to

- (a) define the **base case(s)** and show the claim holds for them
- (b) state the **inductive hypothesis** assuming that the claim holds true for all $n < k$ ($n \in \mathbb{N}$)
- (c) prove that the claim holds for $n = k$ in the rest of the **inductive step**
 - ▶ You can also do $n \leq k$ and then $n = k + 1$.
 - ▶ **Always use strong induction!** Your inductive hypothesis must hold for *all* values up to k .

How many base cases?

Make sure to justify all the base cases that are necessary to establish the claim.

How many base cases?

Make sure to justify all the base cases that are necessary to establish the claim.

- ▶ What base cases do you need to prove this claim:
 $\forall n, f_n < 2^n$, where $f_{n+1} = f_n + f_{n-1}$?

Induction Example I

Prove that the following holds for all natural numbers n .

$$\sum_{i=0}^n x^i = \frac{x^{n+1} - 1}{x - 1}$$

Induction Example I

Prove that the following holds for all natural numbers n .

$$\sum_{i=0}^n x^i = \frac{x^{n+1} - 1}{x - 1}$$

► Base case(s): For _____, we have

► Inductive hypothesis: Suppose $\left[\right]$

► Inductive step: Consider $n = \underline{\hspace{2cm}}$. We want to show that $\left[\right]$.

Therefore $\left[\right]$, which is what we needed to show.

Induction Example I

- ▶ Inductive step

$$\left(\sum_{i=0}^n x^i = \frac{x^{n+1}-1}{x-1} \right)$$

Induction Example I (Solution)

Prove that the following holds for all natural numbers n .

$$\sum_{i=0}^n x^i = \frac{x^{n+1} - 1}{x - 1}$$

- ▶ Base case(s): For $n = 0$, we have $\sum_{i=0}^n x^i = x^0 = 1$ and $\frac{x^{n+1}-1}{x-1} = \frac{x^{0+1}-1}{x-1} = \frac{x-1}{x-1} = 1$. So $\sum_{i=0}^n x^i = \frac{x^{n+1}-1}{x-1}$.
- ▶ Inductive hypothesis: Suppose $\sum_{i=0}^n x^i = \frac{x^{n+1}-1}{x-1}$ for $n = 0, 1, \dots, k$.
- ▶ Inductive step (next slide)

Induction Example I (Solution)

- Inductive step: Consider $n = k + 1$. We want to show that

$$\sum_{i=0}^n x^i = \frac{x^{n+1}-1}{x-1}.$$

$$\sum_{i=0}^{k+1} x^i = \sum_{i=0}^{k+1} x^i = x^{k+1} + \sum_{i=0}^k x^i.$$

By the inductive hypothesis, $\sum_{i=0}^k x^i = \frac{x^{k+1}-1}{x-1}$. So:

$$\begin{aligned}\sum_{i=0}^{k+1} x^i &= x^{k+1} + \sum_{i=0}^k x^i = x^{k+1} + \frac{x^{k+1}-1}{x-1} \\ &= \frac{(x-1)x^{k+1}}{x-1} + \frac{x^{k+1}-1}{x-1} \\ &= \frac{x^{k+2} - x^{k+1} + x^{k+1} - 1}{x-1} \\ &= \frac{x^{k+2} - 1}{x-1}.\end{aligned}$$

Therefore $\sum_{i=0}^{k+1} x^i = \frac{x^{k+2}-1}{x-1}$, which is what we needed to show.

Inequality Induction

- ▶ Very similar to ‘equality’ induction! Don’t overthink it! Just manipulating inequalities instead of equations, still primarily algebra

Inequality Induction

- ▶ Very similar to ‘equality’ induction! Don’t overthink it! Just manipulating inequalities instead of equations, still primarily algebra
- ▶ The only ‘trick’ is that you may have to simplify or change to a term to something even smaller/bigger (resp. the inequality) to make your algebra match your ‘goal’. (Confused? The next example should illuminate this more clearly.)

Inequality Induction Example I

Let f_n be the n th Fibonacci number (i.e. $f_{n+1} = f_n + f_{n-1}$).
Prove that $f_n \geq (\frac{3}{2})^{n-2}$.

Inequality Induction Example I

Let f_n be the n th Fibonacci number (i.e. $f_{n+1} = f_n + f_{n-1}$ where $f_1 = f_2 = 1$). Prove that $f_n \geq (\frac{3}{2})^{n-2}$.

- ▶ Base cases: For $n = 1$, we have $f_1 = 1$, $(\frac{3}{2})^{1-2} = \frac{2}{3}$. $1 \geq \frac{2}{3}$.
For $n = 2$, we have $f_2 = 1$, $(\frac{3}{2})^{2-2} = 1$. $1 \geq 1$.
- ▶ Inductive hypothesis: Suppose that $f_n \geq (\frac{3}{2})^{n-2}$ for $n = 1, 2, \dots, k-1$.

- ▶ Inductive step: Consider $n = k$. We want to show that $f_k \geq (\frac{3}{2})^{k-2}$.
 $f_k = \underline{\hspace{2cm}}$

By the inductive hypothesis, we have $f_{k-1} \geq (\frac{3}{2})^{k-3}$ and $f_{k-2} \geq (\frac{3}{2})^{k-4}$.

So $\underline{\hspace{2cm}}$

Therefore $f_n \geq (\frac{3}{2})^{n-2}$, which is what we needed to show.

Inequality Induction Example I

- Inductive step: Consider $n = k$. We want to show that $f_n \geq \left(\frac{3}{2}\right)^{n-2}$.

$$f_k = f_{k-1} + f_{k-2}$$

By the inductive hypothesis, we have $f_{k-1} \geq \left(\frac{3}{2}\right)^{k-3}$ and $f_{k-2} \geq \left(\frac{3}{2}\right)^{k-4}$. So

$$\begin{aligned} f_n &\geq \left(\frac{3}{2}\right)^{n-3} + \left(\frac{3}{2}\right)^{n-4} \\ &\geq \left(\frac{3}{2}\right)^{n-4} \left(\frac{3}{2} + 1\right) = \left(\frac{3}{2}\right)^{n-4} \left(\frac{5}{2}\right) \\ &\geq \left(\frac{3}{2}\right)^{n-4} \left(\frac{9}{4}\right) = \left(\frac{3}{2}\right)^{n-4} \left(\frac{3}{2}\right)^2 \\ &\geq \left(\frac{3}{2}\right)^{n-2} \end{aligned}$$

Therefore $f_n \geq \left(\frac{3}{2}\right)^{n-2}$, which is what we needed to show.

Tree Induction

- ▶ Most important detail to remember: induct on **height of the tree**, *never* the number of nodes/leaves/etc.

Tree Induction

- ▶ Most important detail to remember: induct on **height of the tree**, *never* the number of nodes/leaves/etc.
- ▶ Also critical: build your tree from the ‘bottom-up’ (adding root nodes at the top, never by adding leaves)

Tree Induction

- ▶ Most important detail to remember: induct on **height of the tree**, *never* the number of nodes/leaves/etc.
- ▶ Also critical: build your tree from the ‘bottom-up’ (adding root nodes at the top, never by adding leaves)
- ▶ May have to do casework, e.g. if node has different values depending on number/properties of children

Important Tree Terminology

- ▶ **binary**: each node has 0, 1, or 2 children
- ▶ **n -ary**: each node has between 0 and n children
- ▶ **full**: each node has strictly either 0 or n children
- ▶ **complete**: every level, except possibly the last, is completely filled
- ▶ **perfect**: full and complete; all levels filled, all internal nodes have 2 children, all leaves at same depth

Tree Induction Example

Define a Filbert tree to be a binary tree containing 2D points such that:

- ▶ Each leaf node contains $(3, 1)$, $(-2, -5)$, or $(2, 2)$.
- ▶ An internal node with one child labeled (a, b) has label $(a + 1, b - 1)$.
- ▶ An internal node with two children labeled (x, y) and (a, b) has label $(x + a, y + b)$.

Prove that the point in the root node of any Filbert tree is on or below the line $x = y$.

Tree Induction Example

Prove that the point in the root node of any Filbert tree is on or below the line $x = y$.

Tree Induction Example (Solution)

Prove that the point in the root node of any Filbert tree is on or below the line $x = y$.

Proof by induction on h , where h is the height of the tree.

- ▶ Base case(s): For Filbert tree where $h = 0$, the root node is a leaf and so contains $(3, 1)$, $(-2, -5)$, or $(2, 2)$, all of which are on or below the line $x = y$.
- ▶ Inductive hypothesis: Suppose that the point in the root node of any Filbert tree is on or below the line $x = y$ for trees of height $h = 0, 1, \dots, k - 1$ ($k \geq 1$).

Tree Induction Example (Solution)

- Inductive step: Let T be a Filbert tree of height k . There are 2 cases.

Case 1: The root of T has one child subtree, whose root contains (a, b) . The root of T contains $(a + 1, b - 1)$. By the inductive hypothesis, (a, b) is on or below $x = y$, i.e. $b \leq a$. Since $b \leq a$, $b - 1 \leq a + 1$, so this point is on or below $x = y$.

Case 2: The root of T has two child subtrees, whose roots contain (x, y) and (a, b) . Then the root of T contains $(x + a, y + b)$. By the inductive hypothesis, $y \leq x$ and $b \leq a$. So $y + b \leq x + a \implies (x + a, y + b)$ is on or below $x = y$. In all cases the root node contains a point on or below $x = y$, which is what we needed to show.

(Another) Inequality Induction Example

Prove by induction that for any two lists of nonnegative numbers (x_1, \dots, x_n) and (y_1, \dots, y_n) ,

$$\left(\sum_{i=1}^n x_i^2 \right) \left(\sum_{i=1}^n y_i^2 \right) \geq \left(\sum_{i=1}^n x_i y_i \right)^2$$

You may use the AM-GM inequality: For any real numbers $a, b \geq 0$, $\frac{a+b}{2} \geq \sqrt{ab}$.

(Another) Inequality Induction Example

$$\left(\sum_{i=1}^n x_i^2\right)\left(\sum_{i=1}^n y_i^2\right) \geq \left(\sum_{i=1}^n x_i y_i\right)^2, \\ \frac{a+b}{2} \geq \sqrt{ab}$$

Questions/Examples

Questions/Examples

Questions/Examples

Questions/Examples

Graphs: Review

- (Abstract Def) An (undirected) **graph** is a tuple $G = (V, E)$ where V is any set and $E \subseteq \{\{u, v\} : u, v \in V\}$.

Graphs: Review

- ▶ (Abstract Def) An (undirected) **graph** is a tuple $G = (V, E)$ where V is any set and $E \subseteq \{\{u, v\} : u, v \in V\}$.
- ▶ (Usable Def) A graph is a set of vertices and edges between them.




Figure: A graph with vertices $V = \{a, b, c\}$ and edges $E = \{\{a, b\}, \{b, c\}\}$.





Figure: A graph with edges if nodes are Big-Theta of each other


Special Graphs

- The graph K_n has n nodes and an edge between every pair of vertices:




Special Graphs

- ▶ The graph K_n has n nodes and an edge between every pair of vertices:




- ▶ The graph C_n has n nodes in a single cycle.




Bipartite

A graph is **bipartite** if its vertices can be divided into disjoint sets L and R such that every edge is between L and R .

****Bipartite \Leftrightarrow 2-Colorable \Leftrightarrow No odd cycles****



Bipartite



Not bipartite


Paths, Walks, and Cycles

A **walk** in a graph is a sequence of vertices (v_1, v_2, \dots, v_k) and edges $(e_1, e_2, \dots, e_{k-1})$ where each edge connects the two vertices on either end of it ($e_i = \{v_i, v_{i+1}\}$).

A **path** is a walk that doesn't repeat vertices.


A **cycle** is a path where the start and end vertices are connected with an edge ($\{v_n, v_1\} \in E$).

The **distance** between two vertices is the length of the shortest path between them.




Connected Components

A **connected component** of a graph is a maximal set of vertices where each pair has a connecting walk. The graph below can be split into 3 connected components.



Isomorphisms

- Two graphs $G = (V, E)$ and $G' = (V', E')$ are **isomorphic** if there is a bijection $f : V \rightarrow V'$ such that $\{v_1, v_2\} \in E$ if and only if $\{f(v_1), f(v_2)\} \in E'$. (What type of relation is this?)



Practice Questions I

- For which n is K_n bipartite? C_n ?

Practice Questions I



- ▶ For which n is K_n bipartite? C_n ?
- ▶ For a graph with n vertices, what is an upper bound on the number of distinct paths?

Practice Questions I

- ▶ For which n is K_n bipartite? C_n ?
- ▶ For a graph with n vertices, what is an upper bound on the number of distinct paths?
- ▶ How many edges are in K_n ? C_n ?


Practice Questions I

- ▶ For which n is K_n bipartite? C_n ?
- ▶ For a graph with n vertices, what is an upper bound on the number of distinct paths?
- ▶ How many edges are in K_n ? C_n ?
- ▶ (SU24) Are these graphs isomorphic? What is the diameter of the left graph? What is the distance between a and c ?



Coloring


- ▶ A (proper) k -coloring of a graph $G = (V, E)$ is an function $f : V \rightarrow \{1, 2, \dots, k\}$ such that for every edge $\{u, v\} \in E$, $f(u) \neq f(v)$.



A 3-coloring of this graph (say, red means 1, blue means 2, and green means 3)

Coloring

- ▶ A (proper) k -coloring of a graph $G = (V, E)$ is an function $f : V \rightarrow \{1, 2, \dots, k\}$ such that for every edge $\{u, v\} \in E$, $f(u) \neq f(v)$.
- ▶ The **chromatic number** of a graph G is $\chi(G)$, the smallest k such that G is k -colorable.



A 3-coloring of this graph (say, red means 1, blue means 2, and green means 3)

Coloring

- ▶ K_n has an n -coloring, and no smaller number works.

Coloring

- ▶ K_n has an n -coloring, and no smaller number works.
- ▶ If G has K_n as a subgraph, $\chi(G) \geq n$.

Coloring


- ▶ K_n has an n -coloring, and no smaller number works.
- ▶ If G has K_n as a subgraph, $\chi(G) \geq n$.
- ▶ If G can be colored with k vertices, then $\chi(G) \leq k$.

Practice Questions II

- ▶ What is an upper bound for $\chi(G)$, given that G has n vertices?

Practice Questions II

- ▶ What is an upper bound for $\chi(G)$, given that G has n vertices?
- ▶ (SU24) What is the chromatic number of the graph below? Prove it.



Practice Questions II

- ▶ The *degree* of a vertex is the number of edges incident to it.
In a graph $G = (V, E)$, what is $\sum_{v \in V} \deg(v)$?

Practice Questions II

- ▶ The *degree* of a vertex is the number of edges incident to it. In a graph $G = (V, E)$, what is $\sum_{v \in V} \deg(v)$?
- ▶ (Past Examlet) In a tree of height h , which best describes the diameter of the tree?
(a) h (b) $2h$ (c) $\leq 2h$ (d) $h + 1$ (e) $\leq h$

Tree Induction

Given a tree T and a function $w : T \rightarrow \mathbb{R}_+$, the *weighted sum* of the tree is $\sum_{v \in T} w(v)2^{-h(v)}$, where $h(v)$ is the depth of the node.

A weight function is *fair* if, for every node $v \in T$,

1. If the node has one child c , $w(u) = w(c)$
2. If the node has two children c_1 and c_2 ,
$$w(u) = w(c_1) + w(c_2)$$

Let T be a tree and r be its root. Prove by (strong) induction that for any fair weight function w , the weighed sum of the tree is no more than $2w(r)$.

Questions/Examples

Questions/Examples

Questions/Examples

Questions/Examples