

# JOYAL'S CYLINDER CONJECTURE

ALEXANDER CAMPBELL

**ABSTRACT.** For each pair of simplicial sets  $A$  and  $B$ , the category  $\mathbf{Cyl}(A, B)$  of cylinders (also called correspondences) from  $A$  to  $B$  admits a model structure induced from Joyal's model structure for quasi-categories. In this paper, we prove Joyal's conjecture that a cylinder  $X \in \mathbf{Cyl}(A, B)$  is fibrant if and only if the canonical morphism  $X \rightarrow A \star B$  is an inner fibration, and that a morphism between fibrant cylinders in  $\mathbf{Cyl}(A, B)$  is a fibration if and only if it is an inner fibration. We use this result to give a new proof of a characterisation of covariant equivalences due to Lurie, which avoids the use of the straightening theorem. In an appendix, we introduce a new family of model structures on the slice categories  $\mathbf{sSet}/B$  whose fibrant objects are the inner fibrations with codomain  $B$ , which we use to prove some new results about inner anodyne extensions and inner fibrations.

## CONTENTS

1. Introduction	1
2. Model structures for cylinders	3
3. Cylinders as presheaves	6
4. Change of base	13
5. Joyal's cylinder conjecture	15
6. Covariant equivalences	17
Appendix A. The parametrised Joyal model structure	18
References	24

## 1. INTRODUCTION

Recall that the *collage*<sup>1</sup> of a profunctor  $M: A \rightarrow B$  (i.e. a functor  $M: A^{\text{op}} \times B \rightarrow \mathbf{Set}$ ) is the category  $C(M)$  whose set of objects is the disjoint union  $\text{ob } C(M) = \text{ob } A + \text{ob } B$ , whose hom-sets are given by

$$C(M)(x, y) = \begin{cases} A(x, y) & \text{if } x, y \in A \\ B(x, y) & \text{if } x, y \in B \\ M(x, y) & \text{if } x \in A, y \in B \\ \emptyset & \text{if } x \in B, y \in A, \end{cases}$$

and whose identities and composition are defined in the evident way by those of the categories  $A$  and  $B$ , and by the action of  $M$  on morphisms. There is a unique functor  $C(M) \rightarrow \mathbf{2} = \{0 < 1\}$  whose fibres above 0 and 1 are  $A$  and  $B$  respectively. Bénabou observed that the collage construction defines an equivalence between the category of profunctors (between arbitrary categories) and the slice category  $\mathbf{Cat}/\mathbf{2}$  (see [Str01]).

In quasi-category theory, the category of *cylinders* (or *correspondences*) is defined to be the slice category  $\mathbf{sSet}/\Delta[1]$ . By analogy with the previous paragraph, a cylinder  $p: X \rightarrow \Delta[1]$  may be thought of as a model for the collage of a quasi-categorical profunctor from  $\partial_0 X := p^{-1}(0)$  to  $\partial_1 X := p^{-1}(1)$ . (See [Joy08b, Chapter 7], [Joy08a, §14], [Lur09, §2.3.1], and [Ste18a] for further details and intuition concerning cylinders/correspondences.)

---

*Date:* 31st October 2019.

These are notes in progress for a series of talks given at the Australian Category Seminar in October 2019.

<sup>1</sup>This construction is due to Bénabou [Bén72, Bén73]; its name is due to Walters (see [Str04]).

For each pair of simplicial sets  $A$  and  $B$ , the category  $\mathbf{Cyl}(A, B)$  of *cylinders from  $A$  to  $B$* , or  $(A, B)$ -*cylinders*, is defined to be the fibre of the functor

$$(\partial_0, \partial_1): \mathbf{sSet}/\Delta[1] \longrightarrow \mathbf{sSet} \times \mathbf{sSet}$$

over the object  $(A, B)$ . Thus an object of  $\mathbf{Cyl}(A, B)$  is a simplicial set  $X$  (the *underlying simplicial set* of the cylinder) equipped with a map  $X \longrightarrow \Delta[1]$  whose fibres above 0 and 1 are  $A$  and  $B$  respectively, as displayed below.

$$\begin{array}{ccccc} A & \longrightarrow & X & \longleftarrow & B \\ \downarrow \lrcorner & & \downarrow & & \lrcorner \downarrow \\ \{0\} & \longrightarrow & \Delta[1] & \longleftarrow & \{1\} \end{array}$$

Note that the initial and terminal objects of  $\mathbf{Cyl}(A, B)$  are the disjoint union  $A+B$  and join  $A \star B$  respectively, equipped with the manifest structure maps. Hence every cylinder  $X \in \mathbf{Cyl}(A, B)$  is equipped with canonical morphisms  $A+B \longrightarrow X$  and  $X \longrightarrow A \star B$ .

In [Joy08a, §14.6], Joyal described a model structure on  $\mathbf{Cyl}(A, B)$  – which we call the *Joyal model structure* on  $\mathbf{Cyl}(A, B)$  – created<sup>2</sup> by the forgetful functor  $\mathbf{Cyl}(A, B) \longrightarrow \mathbf{sSet}$  from the Joyal model structure for quasi-categories on  $\mathbf{sSet}$ , about which he made the following conjecture. (Note that, by definition, an object  $X \in \mathbf{Cyl}(A, B)$  is fibrant in this model structure if and only if the canonical morphism  $X \longrightarrow A \star B$  is a fibration in the Joyal model structure for quasi-categories on  $\mathbf{sSet}$ .)

**1.1. Conjecture (Joyal).** *A cylinder  $X \in \mathbf{Cyl}(A, B)$  is fibrant if and only if the canonical morphism  $X \longrightarrow A \star B$  is an inner fibration, and a morphism between fibrant cylinders in  $\mathbf{Cyl}(A, B)$  is a fibration if and only if it is an inner fibration.*

The main goal of this paper is to prove this conjecture (see Theorem 5.4; see §1.3 below for an outline of our proof). Note that the special case of this conjecture in which  $A$  and  $B$  are quasi-categories is easily proven (see [Ste18a, Lemmas 3.7 and 3.8] or Lemma 5.1 below); we prove it for *every* pair of simplicial sets  $A$  and  $B$ .

**1.2. Remark.** For many years, it was an open question whether every monic bijective-on-0-simplices weak categorical equivalence is inner anodyne (see [Joy08a, §2.10]). Were this so, the general case of Joyal’s conjecture would be as easy to prove as the special case in which  $A$  and  $B$  are quasi-categories. However, the author recently proved [Cam20] that this is not so; hence a different argument is required to prove Joyal’s conjecture.

Our proof of Joyal’s conjecture is contained in §§2–5 of this paper. (The contents of each section may be gleaned from its opening paragraph.) In the final section §6, we use this result to give a new proof of Lurie’s characterisation of covariant equivalences (see Theorem 6.5), which avoids the use of the straightening theorem [Lur09, Theorem 2.2.1.2]. In an appendix (Appendix A), we introduce a new family of model structures on the slice categories  $\mathbf{sSet}/B$  (which we call the *parametrised Joyal model structures*) whose fibrant objects are the inner fibrations with codomain  $B$ , using which we prove some new results about inner anodyne extensions and inner fibrations.

**1.3. Outline of proof.** Our proof of Joyal’s conjecture may be outlined as follows.

- (1) We construct (in Theorem 2.11) a model structure on  $\mathbf{Cyl}(A, B)$ , which we call the *ambivariant model structure*, which has the same cofibrations as the Joyal model structure (i.e. the monomorphisms), but whose (fibrations between) fibrant objects are precisely as described in Joyal’s conjecture.

---

<sup>2</sup>A model structure on a category  $\mathcal{A}$  is said to be *created* by a functor  $F: \mathcal{A} \rightarrow \mathcal{C}$  from a model structure on  $\mathcal{C}$  if a morphism  $f$  of  $\mathcal{A}$  is a cofibration, weak equivalence, or fibration in the model structure on  $\mathcal{A}$  if and only if the morphism  $Ff$  is a cofibration, weak equivalence, or fibration respectively in the model structure on  $\mathcal{C}$ .

- (2) We observe that Joyal's conjecture is therefore equivalent to the statement that, on the category  $\mathbf{Cyl}(A, B)$ , the Joyal model structure and the ambivariant model structure coincide. In particular, we know that these two model structures do coincide if  $A$  and  $B$  are quasi-categories (see Corollary 5.2).
- (3) Since every fibration in the Joyal model structure is in particular an inner fibration, it follows that every ambivariant equivalence in  $\mathbf{Cyl}(A, B)$  is a weak categorical equivalence. It remains to prove the converse.
- (4) For each pair of weak categorical equivalences  $u: A \rightarrow A'$  and  $v: B \rightarrow B'$ , we prove that the pushforward functor

$$(u, v)_!: \mathbf{Cyl}(A, B) \rightarrow \mathbf{Cyl}(A', B')$$

- (a) preserves weak categorical equivalences (see Proposition 4.2), and
  - (b) reflects ambivariant equivalences (see Theorem 4.7).
- (5) It then remains to argue as follows (see Theorem 5.3). Let  $u: A \rightarrow A'$  and  $v: B \rightarrow B'$  be weak categorical equivalences in  $\mathbf{sSet}$  such that  $A'$  and  $B'$  are quasi-categories. For any morphism  $f$  in  $\mathbf{Cyl}(A, B)$ ,  $f$  a weak categorical equivalence  $\implies (u, v)_!(f)$  a weak categorical equivalence (by (4)(a))  $\implies (u, v)_!(f)$  an ambivariant equivalence (by (2), since  $A'$  and  $B'$  are quasi-categories)  $\implies f$  an ambivariant equivalence (by (4)(b)). This completes the proof.

## 2. MODEL STRUCTURES FOR CYLINDERS

The goal of this section is to carry out step (1) of the proof of Joyal's conjecture outlined in §1.3. For each pair of simplicial sets  $A$  and  $B$ , we construct both the *Joyal model structure* (Theorem 2.10) and the *ambivariant model structure* (Theorem 2.11) on the category  $\mathbf{Cyl}(A, B)$  described in §1. (Note that the construction of the latter model structure involves the parametrised Joyal model structures introduced in Appendix A.) We shall construct both of these model structures by the following general technique.

**2.1. Restricting model structures.** We say that a model structure on a category  $\mathcal{C}$  *restricts* to a model structure on a full subcategory  $\mathcal{A}$  of  $\mathcal{C}$  if the full inclusion  $\mathcal{A} \rightarrow \mathcal{C}$  creates a model structure on  $\mathcal{A}$  from the model structure on  $\mathcal{C}$ , that is, if the classes consisting of those morphisms in  $\mathcal{A}$  which are cofibrations, weak equivalences, and fibrations respectively in the model structure on  $\mathcal{C}$  form a model structure on  $\mathcal{A}$ .

**2.2. Proposition.** *Let  $\mathcal{C}$  be a cofibrantly generated model category, and let  $\mathcal{A}$  be a full subcategory of  $\mathcal{C}$  which is both reflective and coreflective via an adjoint triple  $L \dashv I \dashv R$ . Then the model structure on  $\mathcal{C}$  restricts to one on  $\mathcal{A}$  if and only if the adjunction*

$$\mathcal{C} \begin{array}{c} \xleftarrow{IL} \\ \perp \\ \xrightarrow{IR} \end{array} \mathcal{C}$$

*is a Quillen adjunction.*

*Proof.* If the model structure on  $\mathcal{C}$  restricts to one on  $\mathcal{A}$ , then the adjunctions  $L \dashv I$  and  $I \dashv R$  are Quillen adjunctions, and hence so is their composite  $IL \dashv IR$ .

Conversely, suppose that the adjunction  $IL \dashv IR$  is a Quillen adjunction. Note that the category  $\mathcal{A}$  is complete and cocomplete, since it is a (co)reflective subcategory of the complete and cocomplete category  $\mathcal{C}$ . By [DCH19, Theorem 2.3], the category  $\mathcal{A}$  admits a model structure in which a morphism  $f$  is a fibration (resp. weak equivalence) if and only if  $If$  is a fibration (resp. weak equivalence) in the model category  $\mathcal{C}$ ; furthermore, the adjunctions  $L \dashv I$  and  $I \dashv R$  are Quillen adjunctions with respect to these model structures.

It remains to show that a morphism  $f$  in  $\mathcal{A}$  is a cofibration in this model structure on  $\mathcal{A}$  if and only if  $If$  is a cofibration in  $\mathcal{C}$ . Necessity follows from the foregoing fact that the functor  $I$  is left Quillen, while sufficiency follows from the fact that  $L$  is left Quillen and the assumption that  $I$  is fully faithful.  $\square$

2.3. *Remark.* One can show (by using [GKR18, Corollary 2.7] and arguing as in the proof above) that the conclusion of Proposition 2.2 also holds under the alternative hypothesis that  $\mathcal{C}$  is an *accessible* model category (in the sense of [Ros17]).

2.4. **Two model structures on the factorisation category  $(A + B)/\mathbf{sSet}/(A \star B)$ .** Let  $A$  and  $B$  be a pair of simplicial sets. We shall use Proposition 2.2 to induce the “Joyal” and “ambivariant” model structures on  $\mathbf{Cyl}(A, B)$  from the following two model structures on the category  $(A + B)/\mathbf{sSet}/(A \star B)$  of factorisations<sup>3</sup> in  $\mathbf{sSet}$  of the canonical inclusion  $A + B \longrightarrow A \star B$ .

The first of these model structures on the factorisation category  $(A + B)/\mathbf{sSet}/(A \star B)$  is the one created by the forgetful functor

$$(A + B)/\mathbf{sSet}/(A \star B) \longrightarrow \mathbf{sSet}$$

from the Joyal model structure for quasi-categories on  $\mathbf{sSet}$ . The existence of this created model structure follows from [Hir18, §2] (which corrects [Hir03, Theorem 7.6.5(3)]).

The second model structure on  $(A + B)/\mathbf{sSet}/(A \star B)$  is induced from the parametrised Joyal model structure on  $\mathbf{sSet}/(A \star B)$  (introduced in Appendix A, see Theorem A.7) as follows. By [Hir03, Theorem 7.6.5(1)] – applied to the inclusion  $A + B \longrightarrow A \star B$  as an object of the category  $\mathbf{sSet}/(A \star B)$  equipped with the parametrised Joyal model structure – there exists a model structure on  $(A + B)/\mathbf{sSet}/(A \star B)$  created by the forgetful functor

$$(A + B)/\mathbf{sSet}/(A \star B) \longrightarrow \mathbf{sSet}/(A \star B)$$

from the parametrised Joyal model structure on  $\mathbf{sSet}/(A \star B)$ .

Since both the Joyal model structure on  $\mathbf{sSet}$  and the parametrised Joyal model structure on  $\mathbf{sSet}/(A \star B)$  are cofibrantly generated, it follows by [Hir05] that both of the above model structures on the factorisation category  $(A + B)/\mathbf{sSet}/(A \star B)$  are cofibrantly generated.

2.5.  **$\mathbf{Cyl}(A, B)$  as a reflective and coreflective subcategory of  $(A + B)/\mathbf{sSet}/(A \star B)$ .** Recall that the disjoint union  $A + B$  and join  $A \star B$  are the initial and terminal objects respectively of the category  $\mathbf{Cyl}(A, B)$ . As observed in [Joy08a, §14.6], the forgetful functor  $\mathbf{Cyl}(A, B) \longrightarrow \mathbf{sSet}$  lifts to a fully faithful functor  $\mathbf{Cyl}(A, B) \longrightarrow (A + B)/\mathbf{sSet}/(A \star B)$ . This full embedding has both a left adjoint, given by the composite

$$(A + B)/\mathbf{sSet}/(A \star B) \longrightarrow \mathbf{sSet}/(A \star B) \xrightarrow{L} \mathbf{Cyl}(A, B)$$

of the forgetful functor and the reflection  $L$  described in §2.6 below, and a right adjoint, given by the composite

$$(A + B)/\mathbf{sSet}/(A \star B) \longrightarrow (A + B)/\mathbf{sSet} \xrightarrow{R} \mathbf{Cyl}(A, B)$$

of the (other) forgetful functor and the coreflection  $R$  described in §2.7 below.

2.6.  **$\mathbf{Cyl}(A, B)$  as a reflective subcategory of  $\mathbf{sSet}/(A \star B)$ .** As described in [Ste18a, Remark 3.5], the fully faithful functor  $\mathbf{Cyl}(A, B) \longrightarrow \mathbf{sSet}/(A \star B)$  has a left adjoint  $L$ , which sends an object  $X \longrightarrow A \star B$  of  $\mathbf{sSet}/(A \star B)$  to the  $(A, B)$ -cylinder  $L(X)$  defined by the pushout below,

$$\begin{array}{ccc} \partial_0 X + \partial_1 X & \longrightarrow & A + B \\ \downarrow & & \downarrow \\ X & \xrightarrow{\quad \quad} & L(X) \end{array} \quad \lrcorner$$

with the induced structure map  $L(X) \longrightarrow A \star B \longrightarrow \Delta[1]$ . It follows that a morphism of  $(A, B)$ -cylinders is a monomorphism in  $\mathbf{Cyl}(A, B)$  if and only if its underlying morphism of simplicial sets is a monomorphism. The essential image of the full embedding  $\mathbf{Cyl}(A, B) \longrightarrow \mathbf{sSet}/(A \star B)$  consists of those morphisms  $X \longrightarrow A \star B$  whose pullback along the inclusion  $A + B \longrightarrow A \star B$  is an isomorphism.

<sup>3</sup>Beware that an object of the factorisation category  $(A + B)/\mathbf{sSet}/(A \star B)$  is not an arbitrary composable pair of morphisms  $A + B \longrightarrow X \longrightarrow A \star B$ , but rather a pair whose composite is the canonical inclusion  $A + B \longrightarrow A \star B$ .

**2.7.  $\mathbf{Cyl}(A, B)$  as a coreflective subcategory of  $(A + B)/\mathbf{sSet}$ .** The fully faithful functor  $\mathbf{Cyl}(A, B) \rightarrow (A + B)/\mathbf{sSet}$  has a right adjoint  $R$  given by the “quasi-categorical collage construction” described in [RV19, Appendix F]. (We shall not need to know anything about this right adjoint beyond its existence.) The essential image of the full embedding  $\mathbf{Cyl}(A, B) \rightarrow (A + B)/\mathbf{sSet}$  consists of those cospans of simplicial sets

$$A \longrightarrow C \longleftarrow B$$

whose left leg  $A \rightarrow C$  is a sieve inclusion and whose right leg  $B \rightarrow C$  is the complementary cosieve inclusion (in the sense of [Joy08b, §7.2] and [Joy08a, §14.5]).

**2.8. Restricted model structures on  $\mathbf{Cyl}(A, B)$ .** The following proposition verifies the necessary and sufficient condition of Proposition 2.2 in our two cases of interest.

**2.9. Proposition.** *The reflection  $L: \mathbf{sSet}/(A \star B) \rightarrow \mathbf{Cyl}(A, B)$  preserves monomorphisms, inner anodyne extensions, and weak categorical equivalences, and inverts any morphism between objects of  $\mathbf{sSet}/(A \star B)$  whose structure maps factor through the inclusion  $A + B \rightarrow A \star B$ .*

*Proof.* By definition (see §2.6), the functor  $L$  sends a morphism  $f: X \rightarrow Y$  in  $\mathbf{sSet}/(A \star B)$  to the morphism  $L(X) \rightarrow L(Y)$  induced by pushout from the diagram below, in which, we note, the left-pointing maps are monomorphisms and the left-hand square is a pullback.

$$\begin{array}{ccccc} X & \longleftarrow & \partial_0 X + \partial_1 X & \longrightarrow & A + B \\ f \downarrow & & \lrcorner \downarrow \partial_0 f + \partial_1 f & & \parallel \\ Y & \longleftarrow & \partial_0 Y + \partial_1 Y & \longrightarrow & A + B \end{array}$$

It follows from the exactness of pushouts of monomorphisms in the presheaf category  $\mathbf{sSet}$  that  $L$  preserves monomorphisms. If  $f$  is inner anodyne, then [Joy08b, Lemma 3.21] implies that the morphism  $\partial_0 f + \partial_1 f$  is also inner anodyne, whence [Ste18b, Lemma 2.5] implies that  $Lf$  is inner anodyne. It follows similarly from [Joy08b, Corollary 7.10] and the gluing lemma (see [Ree74]) that  $L$  preserves weak categorical equivalences.

By construction, any object  $(X, p)$  of  $\mathbf{sSet}/(A \star B)$  whose structure map  $p: X \rightarrow A \star B$  factors through the inclusion  $A + B \rightarrow A \star B$  is sent by  $L$  to the initial  $(A, B)$ -cylinder  $A + B$ . Hence  $L$  inverts any morphism between two such objects.  $\square$

We are now ready to construct the two model structures on  $\mathbf{Cyl}(A, B)$  described in §1, and thus complete step (1) of the proof of Joyal’s conjecture outlined in §1.3. The existence of the first model structure was stated by Joyal [Joy08a, §14.6]; an alternative proof of its existence is given in [Ste18a, Theorem 3.9].

**2.10. Theorem** (the Joyal model structure on  $\mathbf{Cyl}(A, B)$ ). *There exists a model structure on  $\mathbf{Cyl}(A, B)$  created by the forgetful functor  $\mathbf{Cyl}(A, B) \rightarrow \mathbf{sSet}$  from the Joyal model structure for quasi-categories on  $\mathbf{sSet}$ .*

*Proof.* The existence of this model structure is proven by an application of Proposition 2.2 to the first model structure on the factorisation category  $(A + B)/\mathbf{sSet}/(A \star B)$  described in §2.4 and the adjoint triple described in §2.5. The necessary and sufficient condition of Proposition 2.2 follows from the fact, proved in Proposition 2.9, that the reflection  $L: \mathbf{sSet}/(A \star B) \rightarrow \mathbf{Cyl}(A, B)$  preserves monomorphisms and weak categorical equivalences.  $\square$

We call the model structure of Theorem 2.10 the *Joyal model structure* on  $\mathbf{Cyl}(A, B)$ . Note that (by the remark on monomorphisms made in §2.6) a morphism in  $\mathbf{Cyl}(A, B)$  is a cofibration in this model structure if and only if it is a monomorphism in  $\mathbf{Cyl}(A, B)$ . Since, for any cylinder  $X \in \mathbf{Cyl}(A, B)$ , the canonical morphism  $A + B \rightarrow X$  is a monomorphism, we see that every object of  $\mathbf{Cyl}(A, B)$  is cofibrant in this model structure; the same is true of the following model structure.

**2.11. Theorem** (the ambivariant model structure on  $\mathbf{Cyl}(A, B)$ ). *There exists a model structure on  $\mathbf{Cyl}(A, B)$  whose cofibrations are the monomorphisms and whose fibrant objects are those*

*cylinders*  $X \in \mathbf{Cyl}(A, B)$  for which the canonical morphism  $X \rightarrow A \star B$  is an inner fibration. A morphism between fibrant objects in  $\mathbf{Cyl}(A, B)$  is a fibration if and only if it is an inner fibration.

*Proof.* The first statement is proven by an application of Proposition 2.2 to the second model structure on  $(A + B)/\mathbf{sSet}/(A \star B)$  described in §2.4 and the adjoint triple described in §2.5, which implies that there exists a model structure on  $\mathbf{Cyl}(A, B)$  created by the forgetful functor  $\mathbf{Cyl}(A, B) \rightarrow \mathbf{sSet}/(A \star B)$  from the parametrised Joyal model structure on  $\mathbf{sSet}/(A \star B)$ . The necessary and sufficient condition of Proposition 2.2 follows from Proposition A.10 and Proposition 2.9, the latter of which implies that the reflection  $L$  preserves monomorphisms and inner anodyne extensions, and inverts the morphism  $(J, x \circ \text{pr}) \rightarrow (\Delta[0], x)$  in  $\mathbf{sSet}/(A \star B)$  for every 0-simplex  $x$  of  $A \star B$ .

By the construction of this model structure and by Theorem A.7, a morphism between fibrant objects in  $\mathbf{Cyl}(A, B)$  is a fibration if and only if it is an inner fibration and a fibrewise isofibration over  $A \star B$  (see Definition A.3). But this latter property is trivially satisfied, since if an object  $(X, p)$  of  $\mathbf{sSet}/(A \star B)$  is an  $(A, B)$ -cylinder, then the fibre of  $p: X \rightarrow A \star B$  over any vertex of  $A \star B$  is the terminal simplicial set. This proves the second statement.  $\square$

We call the model structure of Theorem 2.11 the *ambivariant model structure* on  $\mathbf{Cyl}(A, B)$ . We say that an object of  $\mathbf{Cyl}(A, B)$  is *ambifibrant* if it is a fibrant object in this model structure, and that a morphism in  $\mathbf{Cyl}(A, B)$  is an *ambivariant equivalence* if it is a weak equivalence in this model structure.

The following proposition gives a recognition principle for left Quillen functors from the ambivariant model structure on  $\mathbf{Cyl}(A, B)$ .

**2.12. Proposition.** *Let  $\mathcal{C}$  be a model category, and let  $F: \mathbf{Cyl}(A, B) \rightarrow \mathcal{C}$  be a cocontinuous functor that sends monomorphisms to cofibrations. Then  $F$  sends every ambivariant equivalence in  $\mathbf{Cyl}(A, B)$  to a weak equivalence in  $\mathcal{C}$  if and only if it sends every inner anodyne extension in  $\mathbf{Cyl}(A, B)$  to a weak equivalence in  $\mathcal{C}$ .*

*Proof.* This follows by a standard argument from Proposition 2.9, Theorem 2.11, [JT07, Lemma 7.14], and [JT07, Proposition 7.15].  $\square$

We conclude this section with an observation about duality, which will help to simplify the exposition of the following two sections.

**2.13. Observation** (duality and the ambivariant model structure). Recall that the category of simplicial sets bears an involution  $(-)^{\text{op}}: \mathbf{sSet} \rightarrow \mathbf{sSet}$  (induced from the non-trivial involution on the category  $\Delta$ ), which sends a simplicial set  $X$  to its *opposite*  $X^{\text{op}}$ . For each pair of simplicial sets  $A$  and  $B$ , this involution defines an isomorphism

$$\mathbf{Cyl}(A, B) \cong \mathbf{Cyl}(B^{\text{op}}, A^{\text{op}})$$

between the category of  $(A, B)$ -cylinders and the category of  $(B^{\text{op}}, A^{\text{op}})$ -cylinders. Since the involution  $(-)^{\text{op}}: \mathbf{sSet} \rightarrow \mathbf{sSet}$  preserves inner fibrations, it follows that this isomorphism respects the ambivariant model structures on these two categories.

### 3. CYLINDERS AS PRESHEAVES

In this section, we make Joyal's observation (see §3.1) that the category  $\mathbf{Cyl}(A, B)$  of  $(A, B)$ -cylinders is equivalent to the category of presheaves over  $\Delta/A \times \Delta/B$  the basis for a deeper analysis of the ambivariant model structure on  $\mathbf{Cyl}(A, B)$  introduced in §2. We use the related equivalence of categories

$$\mathbf{Cyl}(A, B) \simeq [(\Delta/A)^{\text{op}}, \mathbf{sSet}/B]$$

to construct (Proposition 3.16) a *Reedy model structure* on  $\mathbf{Cyl}(A, B)$ , induced from the *covariant model structure* on  $\mathbf{sSet}/B$ . We prove (Theorem 3.20) that the ambivariant model structure on  $\mathbf{Cyl}(A, B)$  is a *Bousfield localisation* of this Reedy model structure, for which the *local objects* are those  $(A, B)$ -cylinders whose corresponding functors  $(\Delta/A)^{\text{op}} \rightarrow \mathbf{sSet}/B$  send the *final vertex maps* in  $\Delta/A$  to covariant equivalences in  $\mathbf{sSet}/B$ . (A dual result relates the ambivariant

model structure on  $\mathbf{Cyl}(A, B)$  to the *contravariant* model structure on  $\mathbf{sSet}/A$  and the *initial* vertex maps in  $\Delta/B$ ; see Remark 3.22.) We will use this result to prove the main theorem (Theorem 4.7) of §4.

**3.1.  $\mathbf{Cyl}(A, B)$  as a presheaf category.** We begin with Joyal's observation that, for each pair of simplicial sets  $A$  and  $B$ , the category  $\mathbf{Cyl}(A, B)$  of  $(A, B)$ -cylinders is equivalent to the category of presheaves over the product  $\Delta/A \times \Delta/B$  of the categories of simplices of  $A$  and  $B$ .

$$\mathbf{Cyl}(A, B) \simeq [(\Delta/A \times \Delta/B)^{\mathrm{op}}, \mathbf{Set}] \quad (3.2)$$

Under this equivalence, an  $(A, B)$ -cylinder  $X$  with structure map  $p: X \rightarrow A \star B$  corresponds to the presheaf on  $\Delta/A \times \Delta/B$  whose value at the object  $(([m], \alpha), ([n], \beta)) \in \Delta/A \times \Delta/B$  is the set of  $(m+1+n)$ -simplices  $\sigma$  of  $X$  such that  $p(\sigma) = \alpha \star \beta$ , i.e. such that the diagram below commutes.

$$\begin{array}{ccc} & & X \\ & \nearrow \sigma & \downarrow p \\ \Delta[m] \star \Delta[n] & \xrightarrow{\alpha \star \beta} & A \star B \end{array}$$

We refer the reader to [Joy08b, Chapter 7] for further details of this equivalence.

Combining the equivalence (3.2) with the standard equivalences

$$\mathbf{sSet}/C \simeq [(\Delta/C)^{\mathrm{op}}, \mathbf{Set}], \quad (3.3)$$

we obtain further equivalences between  $\mathbf{Cyl}(A, B)$  and the functor categories displayed below.

$$[(\Delta/A)^{\mathrm{op}}, \mathbf{sSet}/B] \simeq \mathbf{Cyl}(A, B) \simeq [(\Delta/B)^{\mathrm{op}}, \mathbf{sSet}/A] \quad (3.4)$$

(Note that, in the sequel, we shall use the word *vertical* (resp. *horizontal*) for those properties of  $(A, B)$ -cylinders that are most naturally expressed in terms of their corresponding functors  $(\Delta/A)^{\mathrm{op}} \rightarrow \mathbf{sSet}/B$  (resp.  $(\Delta/B)^{\mathrm{op}} \rightarrow \mathbf{sSet}/A$ ), etc.)

These equivalences of categories enable us to employ various standard constructions (drawn from [JT07, §7], [Ara14, §3], and [RV14, §4], to which we refer the reader for further details) in our analysis of the ambivariant model structure on  $\mathbf{Cyl}(A, B)$ . Following a brief survey of these constructions, the main argument of this section begins in earnest with Proposition 3.14, in which we use these constructions, together with a result of Stevenson (Lemma 3.13), to give an alternative characterisation of the ambifibrant objects of  $\mathbf{Cyl}(A, B)$ .

**3.5. Exterior (Leibniz) products.** Recall that we may define the *exterior product* bifunctor

$$[(\Delta/A)^{\mathrm{op}}, \mathbf{Set}] \times [(\Delta/B)^{\mathrm{op}}, \mathbf{Set}] \xrightarrow{\boxtimes} [(\Delta/A \times \Delta/B)^{\mathrm{op}}, \mathbf{Set}]$$

in the way made manifest by the formula  $(X \boxtimes Y)_{\alpha, \beta} = X_{\alpha} \times Y_{\beta}$ . Furthermore, we may define the corresponding *exterior Leibniz product* (or *exterior pushout-product*) bifunctor between arrow categories

$$[(\Delta/A)^{\mathrm{op}}, \mathbf{Set}]^2 \times [(\Delta/B)^{\mathrm{op}}, \mathbf{Set}]^2 \xrightarrow{\widehat{\boxtimes}} [(\Delta/A \times \Delta/B)^{\mathrm{op}}, \mathbf{Set}]^2,$$

which sends a pair of morphisms  $(f: M \rightarrow N, g: S \rightarrow T)$  to the pushout-corner map

$$f \widehat{\boxtimes} g: (M \boxtimes T) \cup_{M \boxtimes S} (N \boxtimes S) \rightarrow N \boxtimes T$$

of the commutative square displayed below.

$$\begin{array}{ccc} M \boxtimes S & \xrightarrow{1 \boxtimes g} & M \boxtimes T \\ f \boxtimes 1 \downarrow & & \downarrow f \boxtimes 1 \\ N \boxtimes S & \xrightarrow{1 \boxtimes g} & N \boxtimes T \end{array}$$

It is straightforward to show that, under the equivalences (3.2) and (3.3), the exterior product bifunctor corresponds to the composite bifunctor

$$\mathbf{sSet}/A \times \mathbf{sSet}/B \xrightarrow{\star} \mathbf{sSet}/(A \star B) \xrightarrow{L} \mathbf{Cyl}(A, B),$$

where  $L$  denotes the reflection described in §2.6. Furthermore, the exterior Leibniz product bifunctor corresponds under these equivalences to the composite bifunctor

$$(\mathbf{sSet}/A)^2 \times (\mathbf{sSet}/B)^2 \xrightarrow{\hat{\star}} (\mathbf{sSet}/(A \star B))^2 \xrightarrow{L^2} \mathbf{Cyl}(A, B)^2,$$

whose first factor is the *Leibniz join* (or *pushout-join*) bifunctor. We shall henceforth denote these two composite bifunctors by  $\boxtimes$  and  $\hat{\boxtimes}$  respectively, so that  $(X, p) \boxtimes (Y, q) = L(X \star Y, p \star q)$  and  $f \hat{\boxtimes} g = L(f \hat{\star} g)$ .

**3.6. Left and right division.** The exterior product bifunctor

$$\mathbf{sSet}/A \times \mathbf{sSet}/B \xrightarrow{\boxtimes} \mathbf{Cyl}(A, B)$$

is cocontinuous in each variable, and therefore forms part of a two-variable adjunction

$$(\mathbf{sSet}/B)(S, M \setminus X) \cong (\mathbf{Cyl}(A, B))(M \boxtimes S, X) \cong (\mathbf{sSet}/A)(M, X/S),$$

where, following [JT07, §7], we denote<sup>4</sup> the two right adjoint bifunctors

$$(\mathbf{sSet}/A)^{\text{op}} \times \mathbf{Cyl}(A, B) \xrightarrow{\setminus} \mathbf{sSet}/B \quad \mathbf{Cyl}(A, B) \times (\mathbf{sSet}/B)^{\text{op}} \xrightarrow{/} \mathbf{sSet}/A$$

by the symbols of left division and right division, as displayed.

The Yoneda lemma implies the important observation that the functor  $(\Delta/A)^{\text{op}} \rightarrow \mathbf{sSet}/B$  to which an object  $X \in \mathbf{Cyl}(A, B)$  corresponds under the equivalence (3.4) is naturally isomorphic to the composite

$$(\Delta/A)^{\text{op}} \longrightarrow (\mathbf{sSet}/A)^{\text{op}} \xrightarrow{-\setminus X} \mathbf{sSet}/B$$

of the Yoneda embedding and the “left division of  $X$ ” functor.

Furthermore, under the equivalences of §3.1, these “division” bifunctors correspond to the evident *weighted limit* bifunctors, as indicated below.

$$[(\Delta/A)^{\text{op}}, \mathbf{Set}]^{\text{op}} \times [(\Delta/A)^{\text{op}}, \mathbf{sSet}/B] \xrightarrow{\setminus} \mathbf{sSet}/B$$

$$[(\Delta/B)^{\text{op}}, \mathbf{sSet}/A] \times [(\Delta/B)^{\text{op}}, \mathbf{Set}]^{\text{op}} \xrightarrow{/} \mathbf{sSet}/A$$

Thus, for  $M \in \mathbf{sSet}/A$  and  $X \in \mathbf{Cyl}(A, B)$ , the object  $M \setminus X \in \mathbf{sSet}/B$  is the limit of the functor  $X: (\Delta/A)^{\text{op}} \rightarrow \mathbf{sSet}/B$  weighted by the functor  $M: (\Delta/A)^{\text{op}} \rightarrow \mathbf{Set}$ .

**3.7. Observation** (Leibniz joins and lifting problems). Let  $X \in \mathbf{Cyl}(A, B)$  with structure map  $p: X \rightarrow A \star B$ , and consider a pair of morphisms in  $\mathbf{sSet}/A$  and  $\mathbf{sSet}/B$  as displayed below.

$$\begin{array}{ccc} M & \xrightarrow{f} & N \\ & \searrow nf & \swarrow n \\ & A & \end{array} \quad \begin{array}{ccc} S & \xrightarrow{g} & T \\ & \searrow tg & \swarrow t \\ & B & \end{array}$$

By adjointness (see e.g. [JT07, Proposition 7.6], [Ara14, Proposition 3.3], or [RV14, Observation 4.11]) and the formula  $f \hat{\boxtimes} g = L(f \hat{\star} g)$  (see §3.5 above), the following are equivalent.

<sup>4</sup>As in [JT07], so too in this paper is there no risk of confusing these constructions with the similarly denoted slice constructions defined in [Joy02, §3], since the latter do not here appear.



(i) Any lifting problem in  $\mathbf{sSet}$  of the form

$$\begin{array}{ccc} (M \star T) \cup_{M \star S} (N \star S) & \xrightarrow{\quad} & X \\ \widehat{f \star g} \downarrow & \nearrow & \downarrow p \\ N \star T & \xrightarrow{n \star t} & A \star B \end{array}$$

has a solution.

- (ii) The cylinder  $X$  has the right lifting property in  $\mathbf{Cyl}(A, B)$  with respect to the exterior Leibniz product  $f \boxtimes g: (M \boxtimes T) \cup_{M \boxtimes S} (N \boxtimes S) \rightarrow N \boxtimes T$ .
- (iii) The morphism  $f \backslash X: N \backslash X \rightarrow M \backslash X$  has the right lifting property in  $\mathbf{sSet}/B$  with respect to the morphism  $g: (S, tg) \rightarrow (T, t)$ .
- (iv) The morphism  $X/g: X/T \rightarrow X/S$  has the right lifting property in  $\mathbf{sSet}/A$  with respect to the morphism  $f: (M, nf) \rightarrow (N, n)$ .

**3.8. Observation** (cellular presentations of monomorphisms). Every monomorphism in  $\mathbf{sSet}/A$  can be expressed as a countable composite of pushouts of coproducts of the boundary inclusions  $b_m: (\partial\Delta[m], \partial\alpha) \rightarrow (\Delta[m], \alpha)$  for  $([m], \alpha) \in \Delta/A$ . Furthermore, every monomorphism in  $\mathbf{Cyl}(A, B)$  can be expressed as a countable composite of pushouts of coproducts of the exterior Leibniz products

$$b_m \boxtimes b_n: (\Delta[m], \alpha) \boxtimes (\partial\Delta[n], \partial\beta) \cup (\partial\Delta[m], \partial\alpha) \boxtimes (\Delta[n], \beta) \rightarrow (\Delta[m], \alpha) \boxtimes (\Delta[n], \beta)$$

for  $([m], \alpha) \in \Delta/A$  and  $([n], \beta) \in \Delta/B$ . These two statements follow from the Eilenberg–Zilber lemma applied to the categories  $\Delta/A$  and  $\Delta/A \times \Delta/B$  equipped with their Eilenberg–Zilber Reedy structures inherited from the standard Eilenberg–Zilber Reedy structure on  $\Delta$  (see e.g. [Cis19, §1.3]).

Before we can state Proposition 3.14, we must first make a couple of definitions.

**3.9. Recall** (the covariant model structure). There exists a model structure (due to Joyal; see [Joy08b, Chapter 8]) on the slice category  $\mathbf{sSet}/B$  whose cofibrations are the monomorphisms, and whose fibrant objects are the left fibrations with codomain  $B$ , which we shall sometimes call the *left fibrant* objects of  $\mathbf{sSet}/B$ . A morphism between left fibrant objects is a fibration if and only if it is a left fibration. This model structure is called the *covariant model structure* on  $\mathbf{sSet}/B$ . A morphism in  $\mathbf{sSet}/B$  is said to be a *covariant equivalence* if it is a weak equivalence in this model structure.

**3.10. Definition** (Reedy fibrant cylinders). A cylinder  $X \in \mathbf{Cyl}(A, B)$  is said to be *vertically Reedy left fibrant* if the morphism on the left below is a left fibration in  $\mathbf{sSet}/B$  for all  $([m], \alpha) \in \Delta/A$ .

$$b_m \backslash X: (\Delta[m], \alpha) \backslash X \rightarrow (\partial\Delta[m], \partial\alpha) \backslash X \quad X/b_n: X/(\Delta[n], \beta) \rightarrow X/(\partial\Delta[n], \partial\beta)$$

Dually, an object  $X \in \mathbf{Cyl}(A, B)$  is said to be *horizontally Reedy right fibrant* if the morphism on the right above is a right fibration in  $\mathbf{sSet}/A$  for all  $([n], \beta) \in \Delta/B$ .

**3.11. Observation** (Reedy fibrant cylinders recast). It follows from Observation 3.8 that a cylinder  $X \in \mathbf{Cyl}(A, B)$  is vertically Reedy left fibrant if and only if the morphism  $f \backslash X: N \backslash X \rightarrow M \backslash X$  is a left fibration in  $\mathbf{sSet}/B$  for every monomorphism  $f: M \rightarrow N$  in  $\mathbf{sSet}/A$ . In particular, if  $X \in \mathbf{Cyl}(A, B)$  is vertically Reedy left fibrant, then the object  $M \backslash X \in \mathbf{sSet}/B$  is left fibrant for every object  $M \in \mathbf{sSet}/A$ .

Furthermore, Observation 3.7 implies that a cylinder  $X \in \mathbf{Cyl}(A, B)$  is horizontally Reedy right fibrant if and only if the morphism

$$h_m^k \backslash X: (\Delta[m], \alpha) \backslash X \rightarrow (\Lambda^k[m], \Lambda^k[\alpha]) \backslash X,$$

induced by the horn inclusion  $h_m^k: \Lambda^k[m] \rightarrow \Delta[m]$ , is a trivial fibration in  $\mathbf{sSet}/A$  for all  $m \geq 1$ ,  $0 < k \leq m$ , and  $([m], \alpha) \in \Delta/A$ .

For each  $m \geq 1$ , we let  $i_m: \Delta[0] \rightarrow \Delta[m]$  denote the *final vertex map*, which picks out the final vertex  $m$  of  $\Delta[m]$ .

**3.12. Definition** (vertically right local cylinders). A cylinder  $X \in \mathbf{Cyl}(A, B)$  is said to be *vertically right local* if the morphism

$$i_m \backslash X: (\Delta[m], \alpha) \backslash X \longrightarrow (\Delta[0], \alpha_m) \backslash X$$

is a covariant equivalence in  $\mathbf{sSet}/B$  for all  $m \geq 1$  and  $([m], \alpha) \in \Delta/A$ .

Recall that a class  $\mathbf{C}$  of monomorphisms in a category is said to have the *right cancellation property* if  $(u \in \mathbf{C} \text{ and } vu \in \mathbf{C}) \implies v \in \mathbf{C}$  for any composable pair of monomorphisms  $u$  and  $v$ .

**3.13. Lemma** (Stevenson). *Let  $\mathbf{C}$  be a class of monomorphisms of simplicial sets which is closed under composition, stable under pushout, and has the right cancellation property. Then the following are equivalent.*

- (i)  $\mathbf{C}$  contains the horn inclusion  $\Lambda^k[m] \longrightarrow \Delta[m]$  for every  $m \geq 1$  and  $0 < k \leq m$ .
- (ii)  $\mathbf{C}$  contains the final vertex map  $i_m: \Delta[0] \longrightarrow \Delta[m]$  for every  $m \geq 1$ .

*Proof.* See [Ste17, Proposition 2.6]. □

**3.14. Proposition.** *Let  $X \in \mathbf{Cyl}(A, B)$ . The following are equivalent.*

- (i) *The canonical morphism  $X \longrightarrow A \star B$  is an inner fibration.*
- (ii)  *$X$  is vertically Reedy left fibrant and horizontally Reedy right fibrant.*
- (iii)  *$X$  is vertically Reedy left fibrant and vertically right local.*

*Proof.* We first prove the equivalence (i)  $\iff$  (ii). By definition, the canonical morphism  $X \longrightarrow A \star B$  is an inner fibration if and only if every lifting problem of the form displayed below,

$$\begin{array}{ccc} \Lambda^k[l] & \longrightarrow & X \\ \downarrow & \nearrow & \downarrow \\ \Delta[l] & \xrightarrow{\sigma} & A \star B \end{array}$$

where  $l \geq 2$  and  $0 < k < l$ , has a solution. These lifting problems fall into two cases. If  $\sigma: \Delta[l] \longrightarrow A \star B$  factors through the inclusion  $A + B \longrightarrow A \star B$ , then the lifting problem is solved trivially, since the pullback of  $X \longrightarrow A \star B$  along  $A + B \longrightarrow A \star B$  is an isomorphism (see §2.6). Otherwise,  $\sigma$  is of the form  $\alpha \star \beta: \Delta[m] \star \Delta[n] \longrightarrow A \star B$  for some  $([m], \alpha) \in \Delta/A$  and  $([n], \beta) \in \Delta/B$ , in which case the displayed inner horn inclusion is either (see [Joy02, Lemma 3.3]) the Leibniz join

$$(\Lambda^k[m] \longrightarrow \Delta[m]) \hat{\star} (\partial\Delta[n] \longrightarrow \Delta[n])$$

if  $0 < k \leq m$ , or the Leibniz join

$$(\partial\Delta[m] \longrightarrow \Delta[m]) \hat{\star} (\Lambda^{k-m-1}[n] \longrightarrow \Delta[n])$$

if  $m+1 \leq k < m+1+n$ .

It thus follows by Observation 3.7 that the canonical morphism  $X \longrightarrow A \star B$  is an inner fibration if and only if  $(\Delta[m], \alpha) \backslash X \longrightarrow (\partial\Delta[m], \partial\alpha) \backslash X$  is a left fibration for all  $([m], \alpha) \in \Delta/A$  and  $X/(\Delta[n], \beta) \longrightarrow X/(\partial\Delta[n], \partial\beta)$  is a right fibration for all  $([n], \beta) \in \Delta/B$ , that is, if and only if  $X$  is vertically Reedy left fibrant and horizontally Reedy right fibrant.

We now prove the equivalence (ii)  $\iff$  (iii). Suppose that  $X$  is vertically Reedy left fibrant. Let  $\mathbf{C}$  denote the class of monomorphisms  $f: M \longrightarrow N$  in  $\mathbf{sSet}$  with the property that, for any morphism  $n: N \longrightarrow A$ , the induced morphism  $f \backslash X: (N, n) \backslash X \longrightarrow (M, nf) \backslash X$  is a covariant equivalence, or equivalently a trivial fibration (since  $X$  is vertically Reedy left fibrant), in  $\mathbf{sSet}/B$ . This class of monomorphisms  $\mathbf{C}$  is closed under composition, stable under pushout, and has the right cancellation property. Hence Lemma 3.13 implies (via Observation 3.11) that  $X$  is horizontally Reedy right fibrant if and only if it is vertically right local. □

We now use the equivalence  $\mathbf{Cyl}(A, B) \simeq [(\Delta/A)^{\text{op}}, \mathbf{sSet}/B]$  (see §3.1) to construct a “Reedy” model structure on  $\mathbf{Cyl}(A, B)$ , whose weak equivalences are described in the following definition.

**3.15. Definition** (vertical covariant equivalences). A morphism  $f: X \rightarrow Y$  in  $\mathbf{Cyl}(A, B)$  is said to be a *vertical covariant equivalence* if the morphism

$$(\Delta[m], \alpha) \backslash f: (\Delta[m], \alpha) \backslash X \rightarrow (\Delta[m], \alpha) \backslash Y$$

is a covariant equivalence in  $\mathbf{sSet}/B$  for all  $([m], \alpha) \in \Delta/A$ .

**3.16. Proposition.** *There exists a model structure on the category  $\mathbf{Cyl}(A, B)$  whose cofibrations are the monomorphisms, whose weak equivalences are the vertical covariant equivalences, and whose fibrant objects are the vertically Reedy left fibrant objects.*

*Proof.* By for instance [RV14, Theorem 4.18], the Reedy structure on  $\Delta/A$  (inherited from the standard Reedy structure on  $\Delta$ ) and the covariant model structure on  $\mathbf{sSet}/B$  (see Recollection 3.9) cooperate to endow the functor category  $[(\Delta/A)^{\text{op}}, \mathbf{sSet}/B]$  with a Reedy model structure, which transports along the equivalence (3.4) to a model structure on  $\mathbf{Cyl}(A, B)$ . By construction, the weak equivalences of this model structure are the vertical covariant equivalences; by [RV14, Corollary 6.7] and Observation 3.8, the cofibrations are the monomorphisms.

By construction, an object  $X \in \mathbf{Cyl}(A, B)$  is fibrant in this model structure if and only if the morphism

$$b_m \backslash X: (\Delta[m], \alpha) \backslash X \rightarrow (\partial\Delta[m], \partial\alpha) \backslash X$$

is a fibration in the covariant model structure on  $\mathbf{sSet}/B$  for every  $([m], \alpha) \in \Delta/A$ . But it follows (as in Observation 3.11) that  $X$  is fibrant if and only if the morphism  $b_m \backslash X$  is a fibration *between fibrant objects* in the covariant model structure on  $\mathbf{sSet}/B$  for every  $([m], \alpha) \in \Delta/A$ . Hence, by Observation 3.11 and Recollection 3.9, an object of  $\mathbf{Cyl}(A, B)$  is fibrant in this model structure if and only if it is vertically Reedy left fibrant.  $\square$

We call the model structure of Proposition 3.16 the *vertical Reedy covariant model structure* on  $\mathbf{Cyl}(A, B)$ .

The following theorem is phrased in terms of the theory of Bousfield localisations of model categories (see e.g. [Joy08b, Appendix E]), from which we recall a few basic definitions and results.

**3.17. Definition** (Bousfield localisation). Let  $\mathbf{M}$  and  $\mathbf{N}$  be model structures on a category  $\mathcal{C}$ . The model structure  $\mathbf{N}$  is said to be a *Bousfield localisation* of the model structure  $\mathbf{M}$  if  $\mathbf{N}$  has the same class of cofibrations as  $\mathbf{M}$ , and if every fibrant object of  $\mathbf{N}$  is fibrant in  $\mathbf{M}$ .

**3.18. Definition** (local objects). Let  $\mathbf{M}$  and  $\mathbf{N}$  be model structures on a category  $\mathcal{C}$ , and suppose that  $\mathbf{N}$  is a Bousfield localisation of  $\mathbf{M}$ . An object of  $\mathcal{C}$  is said to be *local* (with respect to this Bousfield localisation) if it is weakly equivalent in  $\mathbf{M}$  to a fibrant object of  $\mathbf{N}$ .

**3.19. Lemma.** *Let  $\mathbf{M}$  and  $\mathbf{N}$  be model structures on a category  $\mathcal{C}$ , and suppose that  $\mathbf{N}$  is a Bousfield localisation of  $\mathbf{M}$ . Every weak equivalence in  $\mathbf{M}$  is a weak equivalence in  $\mathbf{N}$ . Conversely, any weak equivalence in  $\mathbf{N}$  between local objects is a weak equivalence in  $\mathbf{M}$ .*

*Proof.* See e.g. [Joy08b, Proposition E.1.10] and [Joy08b, Proposition E.2.21].  $\square$

We now use Proposition 3.14 to prove the main result of this section.

**3.20. Theorem.** *On the category  $\mathbf{Cyl}(A, B)$ , the ambivariant model structure is a Bousfield localisation of the vertical Reedy covariant model structure. An object of  $\mathbf{Cyl}(A, B)$  is local with respect to this Bousfield localisation if and only if it is vertically right local.*

*Proof.* In both model structures, the cofibrations are precisely the monomorphisms. By Proposition 3.14, every ambifibrant object of  $\mathbf{Cyl}(A, B)$  is vertically Reedy left fibrant. This proves the first statement.

As indicated by the diagram below, it is immediate from the definitions and the two-out-of-three property that an object of  $\mathbf{Cyl}(A, B)$  is vertically right local if it is weakly equivalent in

the vertical Reedy covariant model structure to a vertically right local object.

$$\begin{array}{ccc} (\Delta[m], \alpha) \backslash X & \xrightarrow{i_m \backslash X} & (\Delta[0], \alpha_m) \backslash X \\ \sim \downarrow & & \downarrow \sim \\ (\Delta[m], \alpha) \backslash Y & \xrightarrow{i_m \backslash Y} & (\Delta[0], \alpha_m) \backslash Y \end{array}$$

Hence, to prove the second statement, it suffices to prove that a vertically Reedy left fibrant object of  $\mathbf{Cyl}(A, B)$  is ambifibrant if and only if it is vertically right local. But this was shown in Proposition 3.14.  $\square$

**3.21. Corollary.** *In the category  $\mathbf{Cyl}(A, B)$ , every vertical covariant equivalence is an ambivariant equivalence. Conversely, any ambivariant equivalence between vertically right local objects of  $\mathbf{Cyl}(A, B)$  is a vertical covariant equivalence.*

*Proof.* This follows from Theorem 3.20 by Lemma 3.19.  $\square$

This completes the main argument of this section. We conclude this section with a few remarks which, though they will play no role in the sequel, may be of interest to the reader.

**3.22. Remark** (dual results). By Observation 2.13, each of the results in this section has a dual result. For example, to Proposition 3.14 we may add the further equivalent condition

(iv)  $X$  is horizontally Reedy right fibrant and horizontally left local,

where the latter property means that the morphism

$$X/i_0: X/(\Delta[n], \beta) \longrightarrow X/(\Delta[0], \beta_0),$$

induced by the *initial* vertex map  $i_0: \Delta[0] \longrightarrow \Delta[n]$ , is a *contravariant* equivalence in  $\mathbf{sSet}/A$  for all  $n \geq 1$  and  $([n], \beta) \in \Delta/B$ .

**3.23. Remark** (cylinders and bisimplicial sets). (In this remark only, we use the symbol  $\boxtimes$  to denote the exterior product bifunctor  $\boxtimes: \mathbf{sSet} \times \mathbf{sSet} \longrightarrow \mathbf{ssSet}$ .)

As observed by Joyal in [Joy08b, Chapter 7], the equivalence (3.2) yields an equivalence between  $\mathbf{Cyl}(A, B)$  and the category  $\mathbf{ssSet}/(A \boxtimes B)$  of bisimplicial sets over the exterior product  $A \boxtimes B$ . Under this equivalence, the ambifibrant objects of  $\mathbf{Cyl}(A, B)$  correspond to the ambifibrations with codomain  $A \boxtimes B$ , where a morphism of bisimplicial sets is said to be an *ambifibration* if it has the right lifting property with respect to the exterior Leibniz products  $h_m^k \hat{\boxtimes} b_n$  (for  $m \geq 1$ ,  $0 < k \leq m$ ,  $n \geq 0$ ) and  $b_m \hat{\boxtimes} h_n^k$  (for  $m \geq 0$ ,  $n \geq 1$ ,  $0 \leq k < n$ ).

Note that the category of bisimplicial sets bears an involution  $(-)^{\text{vop}}$ , induced from the involution  $(-)^{\text{op}} \times \text{id}$  on  $\Delta \times \Delta$ , which sends a bisimplicial set  $X$  to its *vertical opposite*  $X^{\text{vop}}$ . Note also that  $(A \boxtimes B)^{\text{vop}} = A^{\text{op}} \boxtimes B$ . This involution on  $\mathbf{ssSet}$  interchanges the class of ambifibrations and the class of *left bifibrations* defined in [Cis19, Definition 5.5.10]. It follows that, under the composite equivalence

$$\mathbf{Cyl}(A, B) \xrightarrow{\simeq} \mathbf{ssSet}/(A \boxtimes B) \xrightarrow[\cong]{(-)^{\text{vop}}} \mathbf{ssSet}/(A^{\text{op}} \boxtimes B),$$

the ambivariant model structure on  $\mathbf{Cyl}(A, B)$  corresponds to the *bicovariant* model structure on  $\mathbf{ssSet}/(A^{\text{op}} \boxtimes B)$  constructed in [Cis19, Theorem 5.5.13].

**3.24. Remark** (alternative constructions of the ambivariant model structure). The results of this section suggest a few alternative constructions of the ambivariant model structure on  $\mathbf{Cyl}(A, B)$ . To give just two such alternatives, we could construct it as the Bousfield localisation of the vertical Reedy covariant model structure of Proposition 3.16 at the set of exterior Leibniz products  $h_m^k \hat{\boxtimes} b_n$  for  $m \geq 1$ ,  $0 < k \leq m$ , and  $n \geq 0$  (or at yet some other set of morphisms), or by transporting the bicovariant model structure on  $\mathbf{ssSet}/(A^{\text{op}} \boxtimes B)$  along the composite equivalence displayed in Remark 3.23.

Despite these alternatives, we have chosen to give the construction presented in §2 because it arrives directly at the defining properties of the ambivariant model structure which make

it central to our proof of Joyal's conjecture (viz. it has the same cofibrations as the Joyal model structure on  $\mathbf{Cyl}(A, B)$  and its (fibrations between) fibrant objects are precisely as described in Joyal's conjecture), because it parallels the natural construction of the Joyal model structure on  $\mathbf{Cyl}(A, B)$ , and because it uses, and has thus provided an opportunity to expose, the parametrised Joyal model structures introduced in Appendix A, which are of independent interest.

#### 4. CHANGE OF BASE

The goal of this section is to carry out step (4) of the proof of Joyal's conjecture outlined in §1.3. We prove (Proposition 4.2) that, for each pair of morphisms of simplicial sets  $u$  and  $v$ , the *pushforward–pullback adjunction*  $(u, v)_! \dashv (u, v)^*$  (see §4.1) is a Quillen adjunction with respect to the Joyal and ambivariant model structures (constructed in §2). Furthermore, we use the results of §3 to prove (Theorem 4.7) that, if  $u$  and  $v$  are weak categorical equivalences, then the pushforward–pullback adjunction  $(u, v)_! \dashv (u, v)^*$  is a Quillen equivalence with respect to the ambivariant model structures.

**4.1. The pushforward–pullback adjunction.** Let  $u: A \rightarrow A'$  and  $v: B \rightarrow B'$  be a pair of morphisms of simplicial sets. Recall from [Joy08a, §14.6] that there is an adjunction

$$\mathbf{Cyl}(A, B) \begin{array}{c} \xrightarrow{(u, v)_!} \\ \perp \\ \xleftarrow{(u, v)^*} \end{array} \mathbf{Cyl}(A', B')$$

whose left adjoint sends an  $(A, B)$ -cylinder  $X$  to the *pushforward*  $(A', B')$ -cylinder  $(u, v)_!(X)$  defined by the pushout square on the left below,

$$\begin{array}{ccc} A + B & \xrightarrow{u+v} & A' + B' \\ \downarrow & & \downarrow \\ X & \xrightarrow{\quad} & (u, v)_!(X) \end{array} \quad \begin{array}{ccc} (u, v)^*(Y) & \xrightarrow{\quad} & Y \\ \downarrow \lrcorner & & \downarrow \\ A \star B & \xrightarrow{u \star v} & A' \star B' \end{array}$$

and whose right adjoint sends an  $(A', B')$ -cylinder  $Y$  to the *pullback*  $(A, B)$ -cylinder  $(u, v)^*(Y)$  defined by the pullback square on the right above.

**4.2. Proposition.** *The pushforward–pullback adjunction*

$$\mathbf{Cyl}(A, B) \begin{array}{c} \xrightarrow{(u, v)_!} \\ \perp \\ \xleftarrow{(u, v)^*} \end{array} \mathbf{Cyl}(A', B')$$

is a Quillen adjunction between the Joyal (resp. ambivariant) model structures on  $\mathbf{Cyl}(A, B)$  and  $\mathbf{Cyl}(A', B')$ . In particular, the pushforward functor  $(u, v)_!: \mathbf{Cyl}(A, B) \rightarrow \mathbf{Cyl}(A', B')$  preserves weak categorical equivalences and ambivariant equivalences.

*Proof.* It suffices to show that the left adjoint  $(u, v)_!$  preserves monomorphisms, weak categorical equivalences, and (by Proposition 2.12) inner anodyne extensions. The proof is a reiteration of the proof of Proposition 2.9. The second statement follows from the first because every object of  $\mathbf{Cyl}(A, B)$  is cofibrant in both the Joyal model structure and the ambivariant model structure.  $\square$

**4.3. Observation** (pushforward as left Kan extension). Under the equivalences (3.2), the pushforward–pullback adjunction  $(u, v)_! \dashv (u, v)^*$  corresponds to the adjunction

$$[(\Delta/A \times \Delta/B)^{\text{op}}, \mathbf{Set}] \begin{array}{c} \xrightarrow{(\Delta/u \times \Delta/v)_!} \\ \perp \\ \xleftarrow{(\Delta/u \times \Delta/v)^*} \end{array} [(\Delta/A' \times \Delta/B')^{\text{op}}, \mathbf{Set}]$$

defined by left Kan extension and restriction along (the opposite of) the functor

$$\Delta/A \times \Delta/B \xrightarrow{\Delta/u \times \Delta/v} \Delta/A' \times \Delta/B'.$$

In the special case in which  $u$  is the identity morphism  $1_A$ , the pushforward–pullback adjunction corresponds under the equivalences (3.4) to the adjunction

$$[(\Delta/A)^{\text{op}}, \mathbf{sSet}/B] \begin{array}{c} \xrightarrow{[1, v_!]} \\ \perp \\ \xleftarrow{[1, v^*]} \end{array} [(\Delta/A)^{\text{op}}, \mathbf{sSet}/B']$$

induced from the adjunction

$$\mathbf{sSet}/B \begin{array}{c} \xrightarrow{v_!} \\ \perp \\ \xleftarrow{v^*} \end{array} \mathbf{sSet}/B'$$

whose left and right adjoints are given by composition with, and pullback along, the morphism  $v: B \rightarrow B'$  respectively. Hence, by §3.6, there exists an isomorphism

$$(\Delta[m], \alpha) \setminus (1_A, v)_!(X) \cong v_!((\Delta[m], \alpha) \setminus X) \quad (4.4)$$

in  $\mathbf{sSet}/B'$ , natural in  $([m], \alpha) \in \Delta/A$  and  $X \in \mathbf{Cyl}(A, B)$ .

We now use the results and constructions of §3 to deduce the main theorem of this section from the following theorem of Joyal.

**4.5. Theorem (Joyal).** *If  $v: B \rightarrow B'$  is a weak categorical equivalence, then the adjunction*

$$\mathbf{sSet}/B \begin{array}{c} \xrightarrow{v_!} \\ \perp \\ \xleftarrow{v^*} \end{array} \mathbf{sSet}/B'$$

*is a Quillen equivalence between the covariant model structures on  $\mathbf{sSet}/B$  and  $\mathbf{sSet}/B'$ .*

*Proof.* See [Joy08b, Theorem 10.2] that this adjunction is a Quillen adjunction, and see [Cis19, Theorem 5.2.14] for a proof<sup>5</sup> that it is moreover a Quillen equivalence.  $\square$

**4.6. Lemma.** *For any simplicial set  $A$  and any morphism of simplicial sets  $v: B \rightarrow B'$ , the pushforward functor  $(1_A, v)_!: \mathbf{Cyl}(A, B) \rightarrow \mathbf{Cyl}(A, B')$  preserves vertically right local objects.*

*Proof.* Let  $X$  be a vertically right local object of  $\mathbf{Cyl}(A, B)$  (see Definition 3.12). For each object  $([m], \alpha) \in \Delta/A$ , the natural isomorphism (4.4) implies that the morphism

$$i_m \setminus (1_A, v)_!(X): (\Delta[m], \alpha) \setminus (1_A, v)_!(X) \rightarrow (\Delta[0], \alpha_m) \setminus (1_A, v)_!(X)$$

is isomorphic in  $\mathbf{sSet}/B'$  to the morphism

$$v_!(i_m \setminus X): v_!((\Delta[m], \alpha) \setminus X) \rightarrow v_!((\Delta[0], \alpha_m) \setminus X).$$

But this latter morphism is a covariant equivalence in  $\mathbf{sSet}/B'$ , since  $X$  is vertically right local by assumption and since the functor  $v_!: \mathbf{sSet}/B \rightarrow \mathbf{sSet}/B'$  preserves covariant equivalences by [Joy08b, Theorem 10.2]. Hence  $(1_A, v)_!(X)$  is a vertically right local object of  $\mathbf{Cyl}(A, B')$ .  $\square$

**4.7. Theorem.** *Suppose  $u: A \rightarrow A'$  and  $v: B \rightarrow B'$  are weak categorical equivalences. Then the pushforward–pullback adjunction*

$$\mathbf{Cyl}(A, B) \begin{array}{c} \xrightarrow{(u, v)_!} \\ \perp \\ \xleftarrow{(u, v)^*} \end{array} \mathbf{Cyl}(A', B')$$

*is a Quillen equivalence between the ambivariant model structures on  $\mathbf{Cyl}(A, B)$  and  $\mathbf{Cyl}(A', B')$ . In particular, the pushforward functor  $(u, v)_!: \mathbf{Cyl}(A, B) \rightarrow \mathbf{Cyl}(A', B')$  reflects ambivariant equivalences.*

<sup>5</sup>This proof largely follows (and improves upon) Joyal's original, unpublished proof of this theorem.

*Proof.* By Proposition 4.2, the pushforward–pullback adjunction is a Quillen adjunction between the ambivariant model structures.

Observe that the pushforward–pullback adjunction is isomorphic to the following composite adjunction.

$$\mathbf{Cyl}(A, B) \begin{array}{c} \xrightarrow{(1_A, v)_!} \\ \perp \\ \xleftarrow{(1_A, v)^*} \end{array} \mathbf{Cyl}(A, B') \begin{array}{c} \xrightarrow{(u, 1_{B'})_!} \\ \perp \\ \xleftarrow{(u, 1_{B'})^*} \end{array} \mathbf{Cyl}(A', B')$$

Hence it suffices to consider the case in which  $u$  is an identity and the case in which  $v$  is an identity. We shall prove the Quillen equivalence in the first case, from which the second case follows by duality (see Observation 2.13).

We first prove that the pushforward functor  $(1_A, v)_! : \mathbf{Cyl}(A, B) \rightarrow \mathbf{Cyl}(A, B')$  reflects ambivariant equivalences. Let  $f : X \rightarrow Y$  be a morphism in  $\mathbf{Cyl}(A, B)$  and suppose that the morphism

$$(1_A, v)_!(f) : (1_A, v)_!(X) \rightarrow (1_A, v)_!(Y)$$

is an ambivariant equivalence in  $\mathbf{Cyl}(A, B')$ . Since the pushforward functor  $(1_A, v)_!$  preserves ambivariant equivalences by Proposition 4.2, we may suppose without loss of generality that  $X$  and  $Y$  are ambifibrant, and hence vertically right local by Proposition 3.14. Hence it follows from Lemma 4.6 that  $(1_A, v)_!(X)$  and  $(1_A, v)_!(Y)$  are vertically right local objects of  $\mathbf{Cyl}(A, B')$ . It then follows from Corollary 3.21 that the morphism  $(1_A, v)_!(f)$  is a vertical covariant equivalence in  $\mathbf{Cyl}(A, B')$ , i.e. that the morphism

$$(\Delta[m], \alpha) \setminus (1_A, v)_!(f) : (\Delta[m], \alpha) \setminus (1_A, v)_!(X) \rightarrow (\Delta[m], \alpha) \setminus (1_A, v)_!(Y)$$

is a covariant equivalence in  $\mathbf{sSet}/B'$  for each  $([m], \alpha) \in \Delta/A$ . By the natural isomorphism (4.4), this latter morphism is isomorphic in  $\mathbf{sSet}/B'$  to the morphism

$$v_!((\Delta[m], \alpha) \setminus f) : v_!((\Delta[m], \alpha) \setminus X) \rightarrow v_!((\Delta[m], \alpha) \setminus Y),$$

which is therefore also a covariant equivalence in  $\mathbf{sSet}/B'$ . Therefore, since the functor  $v_! : \mathbf{sSet}/B \rightarrow \mathbf{sSet}/B'$  reflects covariant equivalences by Theorem 4.5, we may deduce that the morphism  $f : X \rightarrow Y$  is a vertical covariant equivalence, and hence an ambivariant equivalence in  $\mathbf{Cyl}(A, B)$  by Corollary 3.21.

It remains to prove that, for each ambifibrant object  $Y \in \mathbf{Cyl}(A, B')$ , the counit morphism

$$\varepsilon_Y : (1_A, v)_!(1_A, v)^*(Y) \rightarrow Y$$

is an ambivariant equivalence in  $\mathbf{Cyl}(A, B')$ . Indeed, this morphism is a vertical covariant equivalence (and hence an ambivariant equivalence by Corollary 3.21), i.e., for each object  $([m], \alpha) \in \Delta/A$ , the morphism

$$(\Delta[m], \alpha) \setminus \varepsilon_Y : (\Delta[m], \alpha) \setminus (1_A, v)_!(1_A, v)^*(Y) \rightarrow (\Delta[m], \alpha) \setminus Y$$

is a covariant equivalence in  $\mathbf{sSet}/B'$ . For, by Observation 4.3, this latter morphism is isomorphic in  $\mathbf{sSet}/B'$  to the counit morphism

$$v_!v^*((\Delta[m], \alpha) \setminus Y) \rightarrow (\Delta[m], \alpha) \setminus Y$$

of the adjunction  $v_! \dashv v^*$ , and is therefore a covariant equivalence by Theorem 4.5, since  $(\Delta[m], \alpha) \setminus Y$  is a left fibrant object of  $\mathbf{sSet}/B$  by Observation 3.11, Proposition 3.14, and the assumption that  $Y$  is ambifibrant.

Hence the pushforward–pullback adjunction is a Quillen equivalence between the ambivariant model structures. This implies the second statement because every object of  $\mathbf{Cyl}(A, B)$  is cofibrant in the ambivariant model structure.  $\square$

## 5. JOYAL'S CYLINDER CONJECTURE

In this section, we use the results of the preceding sections to complete the proof of Joyal's conjecture outlined in §1.3.

We begin with a recollection from [Ste18a, Lemmas 3.7 and 3.8] of the proof of the easy case of Joyal's conjecture.

**5.1. Lemma.** *Suppose  $A$  and  $B$  are quasi-categories. Then, in the Joyal model structure on  $\mathbf{Cyl}(A, B)$ , an object  $X$  is fibrant if and only if the canonical morphism  $X \rightarrow A \star B$  is an inner fibration, and a morphism between fibrant objects is a fibration if and only if it is an inner fibration.*

*Proof.* Recall from Theorem 2.10 that a morphism in  $\mathbf{Cyl}(A, B)$  is a fibration in the Joyal model structure on  $\mathbf{Cyl}(A, B)$  if and only if it (more precisely, its underlying morphism of simplicial sets) is a fibration in the Joyal model structure for quasi-categories on  $\mathbf{sSet}$ . Since every fibration in the Joyal model structure for quasi-categories is in particular an inner fibration, it remains to prove the sufficiency of each condition.

Let  $X \in \mathbf{Cyl}(A, B)$ , and suppose that the canonical morphism  $p: X \rightarrow A \star B$  is an inner fibration. Since  $p$  is an inner fibration between quasi-categories (by [Joy08b, Corollary 3.23]), to prove that  $p$  is a fibration, it suffices to prove that the induced functor between homotopy categories  $\mathrm{ho}(p): \mathrm{ho}(X) \rightarrow \mathrm{ho}(A \star B)$  is an isofibration. In fact, this functor is a *discrete* isofibration (i.e. has the *unique* isomorphism lifting property): for any isomorphism in the category  $\mathrm{ho}(A \star B) \cong \mathrm{ho}(A) \star \mathrm{ho}(B)$  must belong to either of the full subcategories  $\mathrm{ho}(A)$  or  $\mathrm{ho}(B)$ , but the pullback of  $\mathrm{ho}(p)$  along either full subcategory inclusion  $\mathrm{ho}(A) \rightarrow \mathrm{ho}(A \star B)$  or  $\mathrm{ho}(B) \rightarrow \mathrm{ho}(A \star B)$  is an isomorphism (since  $X \in \mathbf{Cyl}(A, B)$ ). Hence  $X$  is a fibrant object in the Joyal model structure on  $\mathbf{Cyl}(A, B)$ .

Now, let  $f: X \rightarrow Y$  be a morphism between fibrant objects in the Joyal model structure on  $\mathbf{Cyl}(A, B)$ , and suppose that  $f$  is an inner fibration. Once again, it suffices to prove that the functor  $\mathrm{ho}(f)$  is an isofibration. Let  $q: Y \rightarrow A \star B$  denote the canonical morphism from  $Y$  to the terminal object of  $\mathbf{Cyl}(A, B)$ . By the previous paragraph, the functors  $\mathrm{ho}(q)$  and  $\mathrm{ho}(qf) = \mathrm{ho}(q) \circ \mathrm{ho}(f)$  are discrete isofibrations, whence the functor  $\mathrm{ho}(f)$  is also a discrete isofibration. Hence  $f$  is a fibration in the Joyal model structure on  $\mathbf{Cyl}(A, B)$ .  $\square$

**5.2. Corollary.** *Suppose  $A$  and  $B$  are quasi-categories. Then, on the category  $\mathbf{Cyl}(A, B)$ , the Joyal model structure and the ambivariant model structure coincide. In particular, a morphism in  $\mathbf{Cyl}(A, B)$  is a weak categorical equivalence if and only if it is an ambivariant equivalence.*

*Proof.* The two model structures have the same cofibrations (namely the monomorphisms) and, by Lemma 5.1, the same fibrant objects. Hence the two model structures coincide by [Joy08b, Proposition E.1.10].  $\square$

We now use the results of the preceding sections to deduce the general case of Joyal's conjecture from this special case.

**5.3. Theorem.** *Let  $A$  and  $B$  be a pair of simplicial sets. On the category  $\mathbf{Cyl}(A, B)$ , the Joyal model structure and the ambivariant model structure coincide.*

*Proof.* Since the two model structures have the same cofibrations (namely the monomorphisms), it suffices to show that they have the same weak equivalences, i.e., that a morphism in  $\mathbf{Cyl}(A, B)$  is a weak categorical equivalence if and only if it is an ambivariant equivalence. Since every fibration in the Joyal model structure is in particular an inner fibration, it follows by [Joy08b, Proposition E.1.10] that every ambivariant equivalence in  $\mathbf{Cyl}(A, B)$  is a weak categorical equivalence. It remains to prove the converse.

Let  $f$  be a weak categorical equivalence in  $\mathbf{Cyl}(A, B)$ . Let  $u: A \rightarrow A'$  and  $v: B \rightarrow B'$  be weak categorical equivalences in  $\mathbf{sSet}$  such that  $A'$  and  $B'$  are quasi-categories (as may be constructed by the small object argument). By Proposition 4.2, the morphism  $(u, v)_!(f)$  is a weak categorical equivalence in  $\mathbf{Cyl}(A', B')$ . Then, since  $A'$  and  $B'$  are quasi-categories, Corollary 5.2 implies that  $(u, v)_!(f)$  is an ambivariant equivalence in  $\mathbf{Cyl}(A', B')$ . Finally, since  $u$  and  $v$  are weak categorical equivalences, Theorem 4.7 implies that  $f$  is an ambivariant equivalence in  $\mathbf{Cyl}(A, B)$ .  $\square$

Therefore, for each pair of simplicial sets  $A$  and  $B$ , everything we have proved about the ambivariant model structure on  $\mathbf{Cyl}(A, B)$  is true also of the Joyal model structure on  $\mathbf{Cyl}(A, B)$ , for the two are one. In particular, we may deduce the following corollary.



**5.4. Theorem** (Joyal's cylinder conjecture). *Let  $A$  and  $B$  be a pair of simplicial sets. In the Joyal model structure on  $\mathbf{Cyl}(A, B)$ , an object  $X$  is fibrant if and only if the canonical morphism  $X \rightarrow A \star B$  is an inner fibration, and a morphism between fibrant objects is a fibration if and only if it is an inner fibration.*

*Proof.* Since the Joyal and ambivariant model structures on  $\mathbf{Cyl}(A, B)$  coincide by Theorem 5.3, they must have the same fibrant objects and the same fibrations between fibrant objects. By the description of the ambivariant model structure given in Theorem 2.11, this proves the theorem.  $\square$

## 6. COVARIANT EQUIVALENCES

In this final section, we use our proof of Joyal's cylinder conjecture to give a new, direct proof (see Theorem 6.5) of a characterisation of covariant equivalences due to Lurie [Lur09, Chapter 2], which avoids the use of the straightening theorem [Lur09, Theorem 2.2.1.2].

**6.1. The left cone functor.** Let  $B$  be a simplicial set. Recall from [Lur09, Definition 2.4.1.2] the *left cone functor*  $C^\triangleleft: \mathbf{sSet}/B \rightarrow \mathbf{sSet}$ , which sends an object  $p: X \rightarrow B$  of  $\mathbf{sSet}/B$  to its *left cone*, that is, the simplicial set  $C^\triangleleft(X, p)$  defined by the pushout square below.

$$\begin{array}{ccc} X & \xrightarrow{p} & B \\ \downarrow & & \downarrow \\ \Delta[0] \star X & \xrightarrow{\quad} & C^\triangleleft(X, p) \end{array}$$

Observe that  $C^\triangleleft(X, p)$  is the underlying simplicial set of a  $(\Delta[0], B)$ -cylinder; indeed, the left cone functor is none other than the composite

$$\mathbf{sSet}/B \xrightarrow{\Delta[0] \boxtimes (-)} \mathbf{Cyl}(\Delta[0], B) \rightarrow \mathbf{sSet}$$

of the displayed exterior product functor (see §3.5; here  $A = \Delta[0]$ ) and the forgetful functor.

**6.2. Lurie's characterisation of covariant equivalences.** In [Lur09, Chapter 2], Lurie proves that, for each simplicial set  $B$ , a morphism in  $\mathbf{sSet}/B$  is a covariant equivalence if and only if it is sent by the left cone functor  $C^\triangleleft: \mathbf{sSet}/B \rightarrow \mathbf{sSet}$  to a weak categorical equivalence.

**6.3. Remark.** While this characterisation of covariant equivalences does not appear to be stated explicitly in [Lur09, Chapter 2], it is nonetheless, by the following argument, a consequence of the results of that chapter.

By [Lur09, Proposition 2.1.4.7], there exists a model structure on  $\mathbf{sSet}/B$  whose cofibrations are the monomorphisms and whose weak equivalences are the morphisms in  $\mathbf{sSet}/B$  that are sent by the composite

$$\mathbf{sSet}/B \xrightarrow{C^\triangleleft} \mathbf{sSet} \xrightarrow{\mathfrak{e}} \mathbf{sSet}\text{-Cat}$$

of the left cone functor and the homotopy coherent realisation functor to Dwyer–Kan equivalences of simplicially enriched categories.

By [Lur09, Corollary 2.2.3.12] (proved as a corollary of the straightening theorem [Lur09, Theorem 2.2.1.2]), the fibrant objects of this model structure are the left fibrations with codomain  $B$ . Hence, by [Joy08b, Proposition E.1.10], this model structure on  $\mathbf{sSet}/B$  coincides with the covariant model structure (as defined and constructed in [Joy08b, Chapter 8]). In particular, a morphism in  $\mathbf{sSet}/B$  is a covariant equivalence if and only if it is sent by the composite functor displayed above to a Dwyer–Kan equivalence.

So the characterisation follows at last from [Lur09, Proposition 2.2.5.8] (whose proof depends on [Lur09, Theorem 2.4.6.1]), which states that a morphism of simplicial sets is a weak categorical equivalence if and only if it is sent by the homotopy coherent realisation functor to a Dwyer–Kan equivalence.

We now use the results of the preceding sections to give a direct proof of Lurie's characterisation of covariant equivalences. We note that this proof does not depend on the straightening theorem.

**6.4. Theorem.** *Let  $B$  be a simplicial set. The adjunction*

$$\mathbf{Cyl}(\Delta[0], B) \begin{array}{c} \xleftarrow{\Delta[0] \boxtimes (-)} \\ \perp \\ \xrightarrow{\Delta[0] \setminus (-)} \end{array} \mathbf{sSet}/B$$

*is a Quillen equivalence between the Joyal model structure on  $\mathbf{Cyl}(\Delta[0], B)$  and the covariant model structure on  $\mathbf{sSet}/B$ .*

*Proof.* Under the equivalence (3.4), this adjunction corresponds to the adjunction

$$[\Delta^{\mathrm{op}}, \mathbf{sSet}/B] \begin{array}{c} \xleftarrow{\mathrm{cst}} \\ \perp \\ \xrightarrow{\mathrm{ev}_0} \end{array} \mathbf{sSet}/B$$

whose left adjoint sends an object  $X$  of  $\mathbf{sSet}/B$  to the constant simplicial object in  $\mathbf{sSet}/B$  with value  $X$ , and whose right adjoint sends a simplicial object in  $\mathbf{sSet}/B$  to its value at  $[0] \in \Delta$ . Moreover, by Theorems 3.20 and 5.3, the model structure on  $[\Delta^{\mathrm{op}}, \mathbf{sSet}/B]$  which corresponds under this equivalence to the Joyal model structure on  $\mathbf{Cyl}(\Delta[0], B)$  is the Bousfield localisation of the Reedy model structure (with respect to the covariant model structure on  $\mathbf{sSet}/B$ ) whose local objects are the weakly constant simplicial objects. But this is precisely the “canonical model structure” (in the sense of [RSS01, Theorem 3.1]) on  $[\Delta^{\mathrm{op}}, \mathbf{sSet}/B]$  with respect to the covariant model structure on  $\mathbf{sSet}/B$ , so [RSS01, Theorem 3.9] implies that this adjunction is a Quillen equivalence.  $\square$

**6.5. Theorem (Lurie).** *Let  $B$  be a simplicial set. A morphism in  $\mathbf{sSet}/B$  is a covariant equivalence if and only if it is sent by the left cone functor  $C^{\triangleleft}: \mathbf{sSet}/B \rightarrow \mathbf{sSet}$  to a weak categorical equivalence.*

*Proof.* As observed in §6.1, the left cone functor is the composite

$$\mathbf{sSet}/B \xrightarrow{\Delta[0] \boxtimes (-)} \mathbf{Cyl}(\Delta[0], B) \longrightarrow \mathbf{sSet}$$

of the exterior product functor and the forgetful functor. Thus the result follows from Theorem 6.4 (since every object of  $\mathbf{sSet}/B$  is cofibrant in the covariant model structure).  $\square$

**6.6. Remark** (two definitions of covariant equivalences).

## APPENDIX A. THE PARAMETRISED JOYAL MODEL STRUCTURE

In this appendix, we introduce and study a new family of model structures on the slice categories  $\mathbf{sSet}/B$ , whose fibrant objects are the inner fibrations with codomain  $B$ , which we call the *parametrised Joyal model structure* on  $\mathbf{sSet}/B$  (see Theorem A.7). We use this family of model structures to define a new class of morphisms of simplicial sets, which we call the *absolute weak categorical equivalences* (see Definition A.21), using which we prove some new results about inner anodyne extensions and inner fibrations (see §A.20), building on [Cam20], and give a new proof of a theorem of Stevenson (see Proposition A.27).

**A.1. Remark.** The constructions and results of this section are inspired by, but not an instance of, the general theory presented in [Cis19, §2.5]. The existence of the parametrised Joyal model structures studied in this section is prefigured in [Cis19, Remark 5.1.22], where it is observed that the inner fibrations with codomain  $B$  are (among the) fibrant objects in the minimal Cisinski model structure on  $\mathbf{sSet}/B$ .

**A.2. Fibrewise isofibrations.** Recall that a morphism between quasi-categories  $p: X \rightarrow Y$  is an *isofibration* if it is an inner fibration and if the induced functor between homotopy categories  $\mathrm{ho}(p): \mathrm{ho}(X) \rightarrow \mathrm{ho}(Y)$  is an isofibration in the ordinary sense. Let  $J$  denote the nerve of the contractible groupoid with two objects 0 and 1. It follows from Joyal’s lifting theorem that an inner fibration between quasi-categories is an isofibration if and only if it has the right lifting property in  $\mathbf{sSet}$  with respect to the end-point inclusion  $\{0\} \rightarrow J$  (see [Joy02]).

A morphism between quasi-categories is a fibration in the *Joyal model structure* on  $\mathbf{sSet}$  (whose cofibrations are the monomorphisms, and whose fibrant objects are the quasi-categories) if and only if it is an isofibration. In Theorem A.7 below, we shall construct, for each simplicial set  $B$ , a model structure on  $\mathbf{sSet}/B$  (which we call the *parametrised Joyal model structure* on  $\mathbf{sSet}/B$ ) whose cofibrations are the monomorphisms and whose fibrant objects are the inner fibrations with codomain  $B$ . We shall prove that, in this model structure, a morphism between fibrant objects is a fibration if and only if it is a *fibrewise isofibration* in the following sense.

**A.3. Definition** (fibrewise isofibrations). Let  $B$  be a simplicial set, and let  $p: X \rightarrow B$  and  $q: Y \rightarrow B$  be inner fibrations. A morphism  $f: (X, p) \rightarrow (Y, q)$  in  $\mathbf{sSet}/B$  is said to be a *fibrewise isofibration* over  $B$  if it is an inner fibration and if, for each 0-simplex  $b \in B$ , the induced morphism between fibres  $f_b: X_b \rightarrow Y_b$  is an isofibration.

**A.4. A fundamental lemma of quasi-category theory.** We shall construct the aforementioned parametrised Joyal model structures by Cisinski's method for constructing model structures on presheaf categories presented in [Cis06, §1.3] and [Cis19, §2.4]. Our characterisation of the (fibrations between) fibrant objects in these model structures will depend on the following lemma.

**A.5. Lemma.** *Let  $j: S \rightarrow T$  be a bijective-on-0-simplices monomorphism of simplicial sets, and suppose that  $T$  is a quasi-category. Then the Leibniz product*

$$\partial_0 \hat{\times} j: (J \times S) \cup_{\{0\} \times S} (\{0\} \times T) \rightarrow J \times T$$

*is an inner anodyne extension.*

*Proof.* Since its codomain  $J \times T$  is a quasi-category, it suffices to show that the morphism  $\partial_0 \hat{\times} j$  has the left lifting property with respect to every inner fibration between quasi-categories  $p: X \rightarrow Y$ . Since  $j$  is a bijective-on-0-simplices monomorphism, it can be expressed as a countable composite of pushouts of coproducts of boundary inclusions  $b_n: \partial\Delta[n] \rightarrow \Delta[n]$  for  $n \geq 1$ , whence it suffices to show that the Leibniz product  $\partial_0 \hat{\times} b_n$  has the left lifting property with respect to  $p$  for every  $n \geq 1$ . So by adjointness (and the remarks above about isofibrations), it suffices to show that the Leibniz cotensor  $b_n \hat{\lrcorner} p: X^{\Delta[n]} \rightarrow X^{\partial\Delta[n]} \times_{Y^{\partial\Delta[n]}} Y^{\Delta[n]}$ , which is an inner fibration between quasi-categories, is an isofibration for every  $n \geq 1$ . This follows from the Leibniz product version of Joyal's lifting theorem (see e.g. the proof of [Cis19, Theorem 3.5.9]).  $\square$

**A.6. Recall** (the standard simplicial enrichment of  $\mathbf{sSet}/B$ ). For each simplicial set  $B$ , the slice category  $\mathbf{sSet}/B$  admits a standard tensored (and cotensored) enrichment over the cartesian closed category  $\mathbf{sSet}$ . The *tensor*  $S \otimes (X, p)$  of an object  $(X, p)$  in  $\mathbf{sSet}/B$  by a simplicial set  $S$  is the object  $(S \times X, p \circ \text{pr}_2)$  of  $\mathbf{sSet}/B$ . For each pair of objects  $(X, p), (Y, q)$  in  $\mathbf{sSet}/B$ , their *simplicial hom*  $\mathbf{Fun}_B((X, p), (Y, q))$  (which we shall often simply denote by  $\mathbf{Fun}_B(X, Y)$ ) is the simplicial set defined by the pullback square below.

$$\begin{array}{ccc} \mathbf{Fun}_B((X, p), (Y, q)) & \longrightarrow & Y^X \\ \downarrow \lrcorner & & \downarrow q^X \\ \Delta[0] & \xrightarrow{\lceil p \rceil} & B^X \end{array}$$

**A.7. Theorem** (the parametrised Joyal model structure). *Let  $B$  be a simplicial set. There exists a cofibrantly generated model structure on the category  $\mathbf{sSet}/B$  whose cofibrations are the monomorphisms and whose fibrant objects are the inner fibrations with codomain  $B$ . A morphism between fibrant objects is a fibration if and only if it is fibrewise isofibration over  $B$ .*

*Proof.* Since the category  $\mathbf{sSet}/B$  is equivalent to the presheaf category  $[(\Delta/B)^{\text{op}}, \mathbf{Set}]$ , we may apply the method of [Cis06, §1.3] and [Cis19, §2.4] for constructing cofibrantly generated model structures on presheaf categories whose cofibrations are the monomorphisms. Recall that one constructs a model structure by this method by specifying an *exact cylinder* (see [Cis06, Définition 1.3.6] or [Cis19, Definition 2.4.8]) and a class of *anodyne extensions* relative to that exact cylinder (see [Cis06, Définition 1.3.10] or [Cis19, Definition 2.4.11]).

We define our exact cylinder on  $\mathbf{sSet}/B$  to be the endofunctor  $J \otimes (-)$  on  $\mathbf{sSet}/B$  given by tensoring (see Recollection A.6) with the object  $J \in \mathbf{sSet}$ ; the required “boundary inclusion” and “projection” natural transformations are given by tensoring with the boundary inclusion  $\partial J \rightarrow J$  (where  $\partial J = \{0, 1\}$ ) and the morphism  $J \rightarrow \Delta[0]$ . The exact cylinder axioms DH1 and DH2 are easily verified (cf. [Cis06, Exemple 1.3.8] or [Cis19, Example 2.4.10]).

We define our class  $\mathbf{An}$  of anodyne extensions relative to this exact cylinder to be the (weak) saturation of the set of morphisms in  $\mathbf{sSet}/B$  consisting of the end-point inclusion  $\partial_0: (\Delta[0], b) \rightarrow J \otimes (\Delta[0], b)$  for every 0-simplex  $b$  in  $B$ , and the inner horn inclusion  $h_n^k: (\Lambda^k[n], \Lambda^k[\beta]) \rightarrow (\Delta[n], \beta)$  for every  $n \geq 2$ ,  $0 < k < n$ , and  $n$ -simplex  $\beta$  in  $B$ , as displayed below.

$$\begin{array}{ccc} \Delta[0] & \xrightarrow{\partial_0} & J \times \Delta[0] \\ & \searrow b & \swarrow b \circ \text{pr}_2 \\ & & B \end{array} \qquad \begin{array}{ccc} \Lambda^k[n] & \xrightarrow{h_n^k} & \Delta[n] \\ & \searrow \Lambda^k[\beta] & \swarrow \beta \\ & & B \end{array}$$

Note that every inner anodyne extension in  $\mathbf{sSet}/B$  belongs to the class  $\mathbf{An}$ .

We now show that this class  $\mathbf{An}$  satisfies the axioms for a class of anodyne extensions relative to the exact cylinder  $J \otimes (-)$ . Axiom An0 is immediate from the definition of the class  $\mathbf{An}$ . By Observation 3.8, to verify axiom An1 it suffices to show that the Leibniz tensor

$$\partial_0 \hat{\otimes} b_n: J \otimes (\partial\Delta[n], \partial\beta) \cup_{\{0\} \otimes (\partial\Delta[n], \partial\beta)} \{0\} \otimes (\Delta[n], \beta) \rightarrow J \otimes (\Delta[n], \beta)$$

belongs to  $\mathbf{An}$  for every  $n \geq 0$  and  $n$ -simplex  $\beta$  of  $B$ . If  $n = 0$ , this is immediate from the definition of  $\mathbf{An}$ ; if  $n \geq 1$ , this follows from Lemma A.5. To verify axiom An2, it suffices to show that the Leibniz tensor of every generating anodyne extension with the boundary inclusion  $\partial J \rightarrow J$  belongs to  $\mathbf{An}$ . But (the underlying morphism of simplicial sets of) every such Leibniz tensor is an inner anodyne extension, since the Leibniz product of any monomorphism and any inner horn inclusion is inner anodyne, and since the Leibniz product of  $\{0\} \rightarrow J$  and  $\partial J \rightarrow J$  is inner anodyne by Lemma A.5.

Hence it follows from [Cis06, Théorème 1.3.22, Proposition 1.3.36] or [Cis19, Theorem 2.4.19] that there exists a cofibrantly generated model structure on  $\mathbf{sSet}/B$  whose cofibrations are the monomorphisms, and in which an object is fibrant (resp. a morphism between fibrant objects is a fibration) if and only if it has the right lifting property with respect to the class of morphisms  $\mathbf{An}$ . It is straightforward to show that these (fibrations between) fibrant objects are precisely as described in the statement of the theorem.  $\square$

For each simplicial set  $B$ , we call the model structure of Theorem A.7 the *parametrised Joyal model structure* on  $\mathbf{sSet}/B$ .

**A.8. Remark.** In the case  $B = \Delta[0]$ , the parametrised model structure on  $\mathbf{sSet}/\Delta[0] \cong \mathbf{sSet}$  is precisely the Joyal model structure for quasi-categories. More generally, if  $B$  is a quasi-category whose homotopy category has no non-identity isomorphisms, then the parametrised Joyal model structure on  $\mathbf{sSet}/B$  coincides with the model structure induced from the Joyal model structure on  $\mathbf{sSet}$ , whose fibrant objects are the fibrations in  $\mathbf{sSet}$  with codomain  $B$ . This coincidence does not occur in general: for example, these two model structures do not coincide in the case  $B = J$ , since not every inner fibration with codomain  $J$  is an isofibration (e.g.  $\{0\} \rightarrow J$ ).

**A.9. Perspective** (parametrised categories). [*In which the parametrised Joyal model structures are related to the parametrised categories of [SS88].*]

The following proposition gives a recognition principle for left Quillen functors from the parametrised Joyal model structure on  $\mathbf{sSet}/B$ .

**A.10. Proposition.** *Let  $B$  be a simplicial set, and let  $F: \mathbf{sSet}/B \rightarrow \mathcal{C}$  be a cocontinuous functor to a model category  $\mathcal{C}$  that sends monomorphisms to cofibrations. Then  $F$  sends every weak equivalence in the parametrised Joyal model structure on  $\mathbf{sSet}/B$  to a weak equivalence in  $\mathcal{C}$  if and only if it sends the following morphisms to weak equivalences in  $\mathcal{C}$ :*

- (i) for each 0-simplex  $b \in B_0$ , the projection  $\text{pr}: J \otimes (\Delta[0], b) \longrightarrow (\Delta[0], b)$ , and
- (ii) for each  $n \geq 2$ ,  $0 < k < n$ , and  $n$ -simplex  $\beta \in B_n$ , the inner horn inclusion  $(\Lambda^k[n], \Lambda^k[\beta]) \longrightarrow (\Delta[n], \beta)$ .

*Proof.* This follows by a standard argument from Theorem A.7, [JT07, Proposition 7.14], and [JT07, Proposition 7.15]. Alternatively, it follows from the construction of the model structure given in the proof of Theorem A.7 by [Cis19, Proposition 2.4.40].  $\square$

**A.11. Proposition.** *For each morphism of simplicial sets  $u: B \longrightarrow B'$ , the adjunction*

$$\mathbf{sSet}/B \begin{array}{c} \xrightarrow{u_!} \\ \perp \\ \xleftarrow{u^*} \end{array} \mathbf{sSet}/B'$$

*is a Quillen adjunction between the parametrised Joyal model structures on  $\mathbf{sSet}/B$  and  $\mathbf{sSet}/B'$ . In particular, for each simplicial set  $B$ , every weak equivalence in the parametrised Joyal model structure on  $\mathbf{sSet}/B$  is a weak categorical equivalence.*

*Proof.* The first statement follows from Proposition A.10. The second statement follows from the first by considering the morphism  $B \longrightarrow \Delta[0]$ .  $\square$

**A.12. Remark.** By Remark A.8, it is not true in general that every weak categorical equivalence is a weak equivalence in the parametrised Joyal model structure on  $\mathbf{sSet}/B$ . For example, the end-point inclusion  $\partial_0: \{0\} \longrightarrow J$  is a weak categorical equivalence, but the morphism  $\partial_0: (\{0\}, \partial_0) \longrightarrow (J, 1_J)$  is not a weak equivalence in the parametrised Joyal model structure on  $\mathbf{sSet}/J$ , since it is a morphism between cofibrant-fibrant objects, but there exists no morphism  $(J, 1_J) \longrightarrow (\{0\}, \partial_0)$  in  $\mathbf{sSet}/J$ .

**A.13. Proposition.** *For each simplicial set  $B$ , the category  $\mathbf{sSet}/B$ , when equipped with the parametrised Joyal model structure and its standard simplicial enrichment, is enriched as a model category over the Joyal model structure on  $\mathbf{sSet}$ .*

*Proof.* This follows by two applications of Lemma A.10 from the facts that the Leibniz product in  $\mathbf{sSet}$  of a monomorphism with a monomorphism (resp. an inner anodyne extension) is again a monomorphism (resp. an inner anodyne extension), and that  $J$  is an injective object of  $\mathbf{sSet}$ .  $\square$

**A.14. Observation** (the  $\infty$ -cosmos of  $B$ -parametrised quasi-categories). It follows from [RV17, Lemma 2.2.1] applied to Proposition A.13 that the full simplicial subcategory of  $\mathbf{sSet}/B$  (with its standard simplicial enrichment, see Recollection A.6) whose objects are the inner fibrations with codomain  $B$  is an  $\infty$ -cosmos (in the sense of [RV17, Definition 2.1.1]), in which the “isofibrations” are the fibrewise isofibrations over  $B$  (see Definition A.3).

**A.15. Equivalences of parametrised quasi-categories.** Recall that a morphism of quasi-categories is a weak categorical equivalence if and only if it is essentially surjective on objects and fully faithful (see e.g. [Cis19, Theorem 3.9.7]). The goal for the next part of this appendix is to give an analogous characterisation of the weak equivalences between fibrant objects in the parametrised Joyal model structures on the categories  $\mathbf{sSet}/B$ .

Let us first fix some notation. Let  $p: X \longrightarrow B$  be an inner fibration. For each  $n \geq 0$  and each  $n$ -simplex  $\beta$  of  $B$ , let us denote by  $X_\beta$  the fibre of the inner fibration  $p$  over  $\beta$ , i.e. the simplicial set defined by the pullback square below.

$$\begin{array}{ccc} X_\beta & \longrightarrow & X \\ \downarrow & \lrcorner & \downarrow p \\ \Delta[n] & \xrightarrow{\beta} & B \end{array}$$

**A.16. Definition.** Let  $B$  be a simplicial set, and let  $p: X \longrightarrow B$  and  $q: Y \longrightarrow B$  be inner fibrations. A morphism  $u: (X, p) \longrightarrow (Y, q)$  in  $\mathbf{sSet}/B$  is said to be:

- (i) *fibrewise essentially surjective on objects* if, for every 0-simplex  $b$  of  $B$ , the induced morphism  $u_b: X_b \rightarrow Y_b$  is essentially surjective on objects (as a morphism of quasi-categories);
- (ii) *parametrised fully faithful* if, for every pair of 0-simplices  $x, y$  in  $X$ , and every 1-simplex  $f: p(x) \rightarrow p(y)$  in  $B$ , the induced morphism on hom-spaces  $u: \text{Hom}_{X_f}(x, y) \rightarrow \text{Hom}_{Y_f}(u(x), u(y))$  is an equivalence of Kan complexes.

The following lemma makes precise the sense in which the category  $\mathbf{sSet}/B$ , equipped with the parametrised Joyal model structure, is generated by the 1-simplices of  $B$  under homotopy colimits.

**A.17. Lemma.** *Let  $B$  be a simplicial set and let  $\mathcal{D}$  be a class of objects of  $\mathbf{sSet}/B$  with the following properties:*

- (i)  $\mathcal{D}$  is saturated by monomorphisms in  $\mathbf{sSet}/B$  (see [Cis06, Définition 1.1.12] or [Cis19, Definition 1.3.9]),
- (ii)  $\mathcal{D}$  is replete with respect to weak equivalences in the parametrised model structure on  $\mathbf{sSet}/B$ , and
- (iii) for every 1-simplex  $f$  of  $B$ , the object  $f: \Delta[1] \rightarrow B$  belongs to  $\mathcal{D}$ .

*Then every object of  $\mathbf{sSet}/B$  belongs to  $\mathcal{D}$ .*

*Proof.* It suffices to show that every representable  $\beta: \Delta[n] \rightarrow B$  belongs to  $\mathcal{D}$ . Every 1-simplex of  $B$  belongs to  $\mathcal{D}$  by assumption (iii). Since any 0-simplex of  $B$  is a retract of its degenerate 1-simplex, every 0-simplex of  $B$  belongs to  $\mathcal{D}$  by (i). By (i), any object  $I[n] \rightarrow B$  (where  $I[n]$  denotes the  $n$ -spine) belongs to  $\mathcal{D}$  for each  $n \geq 2$ . For each  $n \geq 2$ , the spine inclusion  $I[n] \rightarrow \Delta[n]$  is inner anodyne; hence by (ii), every simplex of  $B$  belongs to  $\mathcal{D}$ .  $\square$

**A.18. Proposition** (equivalences of parametrised quasi-categories). *Let  $B$  be a simplicial set,  $p: X \rightarrow B$  and  $q: Y \rightarrow B$  inner fibrations, and  $u: X \rightarrow Y$  a morphism such that  $qu = p$ . Then the following are equivalent.*

- (i)  $u: (X, p) \rightarrow (Y, q)$  is a weak equivalence in the parametrised Joyal model structure on  $\mathbf{sSet}/B$ .
- (ii) For every 1-simplex  $f$  of  $B$ , the induced morphism on fibres  $u: X_f \rightarrow Y_f$  is an equivalence of quasi-categories.
- (iii) For every 1-simplex  $f$  of  $B$ , the morphism

$$\mathbf{Fun}_B(1, u): \mathbf{Fun}_B((\Delta[1], f), X) \rightarrow \mathbf{Fun}_B((\Delta[1], f), Y)$$

*is an equivalence of quasi-categories.*

- (iv)  $u: (X, p) \rightarrow (Y, q)$  is fibrewise essentially surjective on objects and parametrised fully faithful.

*Proof.* (i)  $\implies$  (ii) since pullback along  $f: \Delta[1] \rightarrow B$  is right Quillen by Proposition A.11, and so preserve weak equivalences between fibrant objects, together with the observation that the parametrised Joyal model structure on  $\mathbf{sSet}/\Delta[1]$  coincides with the usual slice Joyal model structure.

(ii)  $\implies$  (iii) since  $\mathbf{Fun}_B((\Delta[1], f), X) \cong \mathbf{Fun}_{\Delta[1]}(\Delta[1], X_f)$  and since  $\mathbf{Fun}_{\Delta[1]}(\Delta[1], -)$  is right Quillen by Proposition A.13.

(iii)  $\implies$  (i) from Lemma A.17 by considering the class of objects  $A$  of  $\mathbf{sSet}/B$  for which  $\mathbf{Fun}_B(A, u)$  is an equivalence of quasi-categories.

(iv) is an evident rephrasing of (ii), using the standard theorem that a morphism of quasi-categories is an equivalence if and only if it is essentially surjective on objects and fully faithful.  $\square$

The following proposition is a corollary of the preceding proposition; note however that it can also be easily proved directly.

**A.19. Proposition.** *Let  $p: X \rightarrow B$  be an inner fibration. The following are equivalent.*

- (i)  $p: X \rightarrow B$  is a trivial fibration.
- (ii) For every 1-simplex  $f$  of  $B$ , the fibre  $X_f$  is categorically equivalent to  $\Delta[1]$ .

- (iii) For every 1-simplex  $f$  of  $B$ , the pullback  $X_f \rightarrow \Delta[1]$  of  $p$  along  $f: \Delta[1] \rightarrow B$  is a trivial fibration.
- (iv)  $p$  is surjective on objects and, for each pair of 0-simplices  $x, y \in X_0$  and each 1-simplex  $f: p(x) \rightarrow p(y)$  in  $B$ , the hom-space  $\text{Hom}_{X_f}(x, y)$  is a contractible Kan complex.

*Proof.* This follows from the previous proposition applied to the fibration between fibrant objects  $p: (X, p) \rightarrow (B, 1_B)$  in  $\mathbf{sSet}/B$ . (ii) is intended to be a snappy rephrasing of (iii).  $\square$

**A.20. Absolute weak categorical equivalences.** For many years, the following questions (here suggestively posed) were open:

- (1) Is a monomorphism inner anodyne if and only if it is a surjective-on-0-simplices weak categorical equivalence?
- (2) Is an inner fibration a trivial fibration if and only if it is a surjective-on-0-simplices weak categorical equivalence?

In [Cam20], I showed that the answer to both of these questions is no (amusingly, a single morphism was a counterexample to both questions); note that both statements are true if the codomains of the morphisms in question are assumed to be quasi-categories. But that paper still left unresolved the general question of how to “intrinsically” characterise the inner anodyne extensions among the monomorphisms, and the trivial fibrations among the inner fibrations.

In this section I will show that these statements may be corrected by replacing the property “surjective-on-0-simplices weak categorical equivalence” by the new property *absolute weak categorical equivalence* (which we introduce in Definition A.21); that is, we prove (see Proposition A.24):

- (1) A monomorphism is inner anodyne if and only if it is an absolute weak categorical equivalence.
- (2) An inner fibration is a trivial fibration if and only if it is an absolute weak categorical equivalence.

Note that we have in fact already proved another characterisation of the trivial fibrations among the inner fibrations (see Proposition A.19):

- (3) An inner fibration  $X \rightarrow B$  is a trivial fibration if and only if the fibre  $X_f$  over each 1-simplex  $f$  of  $B$  is categorically equivalent to  $\Delta[1]$ .

This triple of statements should be thought of as analogous to the following triple:

- (1) A monomorphism is right anodyne if and only if it is final.
- (2) A right fibration is a trivial fibration if and only if it is final.
- (3) A right fibration  $X \rightarrow B$  is a trivial fibration if and only if the fibre  $X_b$  over each 0-simplex  $b$  of  $B$  is homotopy equivalent to  $\Delta[0]$ .

**A.21. Definition** (absolute weak categorical equivalence). A morphism of simplicial sets  $u: A \rightarrow B$  is said to be an *absolute weak categorical equivalence* if, for every simplicial set  $C$  and every morphism  $g: B \rightarrow C$ , the morphism  $u: (A, gu) \rightarrow (B, g)$  is a weak equivalence in the parametrised Joyal model structure on  $\mathbf{sSet}/C$ .

**A.22. Example.** Any inner anodyne extension and any trivial fibration is an absolute weak categorical equivalence.

**A.23. Proposition.** The class of absolute weak categorical equivalences is closed under composition and satisfies the right cancellation property.

*Proof.* This is immediate from the definition.  $\square$

**A.24. Proposition.**

- (1) A morphism of simplicial sets  $u: A \rightarrow B$  is an absolute weak categorical equivalence if and only if the morphism  $u: (A, u) \rightarrow (B, 1_B)$  is a weak equivalence in the parametrised Joyal model structure on  $\mathbf{sSet}/B$ .
- (2) An inner fibration is a trivial fibration if and only if it is an absolute weak categorical equivalence.

- (3) *A morphism of simplicial sets is an absolute weak categorical equivalence if and only if it factors as an inner anodyne extension followed by a trivial fibration.*
- (4) *A morphism of simplicial sets is inner anodyne if and only if it is a monic absolute weak categorical equivalence.*

*Proof.* (1) Necessity is immediate from the definition, and sufficiency follows from Proposition A.11.

(2) Let  $p: X \rightarrow B$  be an inner fibration. Then  $p: (X, p) \rightarrow (B, 1_B)$  is a fibration between fibrant objects in the parametrised Joyal model structure on  $\mathbf{sSet}/B$ , and is therefore a weak equivalence therein if and only if it is a trivial fibration.

(3) Any inner anodyne extension and any trivial fibration is an absolute weak categorical equivalence, and the class of absolute weak categorical equivalences is closed under composition. Conversely, let  $u$  be an absolute weak categorical equivalence, and let  $u = pi$  be a factorisation of  $u$  as an inner anodyne extension  $i$  followed by an inner fibration  $p$ . It follows from the right cancellation property of the class of absolute weak categorical equivalences that  $p$  is an absolute weak categorical equivalence, and hence a trivial fibration by (2).

(4) Every inner anodyne extension is both monic and an absolute weak categorical equivalence. The converse follows from part (3).  $\square$

**A.25. Proposition.** *Every absolute weak categorical equivalence is a surjective-on-0-simplices weak categorical equivalence. Any surjective-on-0-simplices weak categorical equivalence whose codomain is a quasi-category is an absolute weak categorical equivalence.*

*Proof.* The first statement follows from part (3) of Proposition A.24. For the partial converse, let  $u: A \rightarrow B$  be a surjective-on-0-simplices weak categorical equivalence whose codomain is a quasi-category. Let  $u = pi$  be a factorisation of  $u$  into an inner anodyne extension  $i$  followed by an inner fibration  $p$ . Then  $p$  is an inner fibration between quasi-categories which is also a surjective-on-0-simplices weak categorical equivalence. Hence  $\mathrm{ho}(p)$  is a surjective equivalence, and in particular an isofibration, and so  $p$  is an isofibration, and hence a trivial fibration. Hence  $u$  is an absolute weak categorical equivalence.  $\square$

**A.26. Example.** Not every surjective-on-0-simplices weak categorical equivalence is an absolute weak categorical equivalence. The morphism  $f: \Delta[1] \rightarrow S$  defined in [Cam20] is a counterexample.

Finally, we recover Stevenson’s result [Ste18b, Theorem 1.5] that the class of inner anodyne extensions satisfies the right cancellation property among monos.

**A.27. Proposition** (Stevenson). *Let  $u: A \rightarrow B$  and  $v: B \rightarrow C$  be a composable pair of monomorphisms of simplicial sets. If  $u$  and  $vu$  are inner anodyne, then so is  $v$ .*

*Proof.* This follows from part (4) of Proposition A.24 and from Proposition A.23.  $\square$

## REFERENCES

- [Ara14] Dimitri Ara. Higher quasi-categories vs higher Rezk spaces. *J. K-Theory*, 14(3):701–749, 2014.
- [Bén72] Jean Bénabou. 2-Dimensional limits and colimits of distributors. Mathematisches Forschungsinstitut Oberwolfach, Tagungsbericht 30, pp. 6–7, 1972.
- [Bén73] Jean Bénabou. Les distributeurs. Univ. Catholique de Louvain, Seminaires de Math. Pure, Rapport 33, 1973.
- [Cam20] Alexander Campbell. A counterexample in quasi-category theory. *Proc. Amer. Math. Soc.*, 148(1):37–40, 2020.
- [Cis06] Denis-Charles Cisinski. Les préfaisceaux comme modèles des types d’homotopie. *Astérisque*, (308):xxiv+390, 2006.
- [Cis19] Denis-Charles Cisinski. *Higher categories and homotopical algebra*, volume 180 of *Cambridge Studies in Advanced Mathematics*. Cambridge University Press, Cambridge, 2019.
- [DCH19] Gabriel C. Drummond-Cole and Philip Hackney. A criterion for existence of right-induced model structures. *Bull. Lond. Math. Soc.*, 51(2):309–326, 2019.
- [GKR18] Richard Garner, Magdalena Kędziorek, and Emily Riehl. Lifting accessible model structures. arXiv:1802.09889v1, 2018.



- [Hir03] Philip S. Hirschhorn. *Model categories and their localizations*, volume 99 of *Mathematical Surveys and Monographs*. American Mathematical Society, Providence, RI, 2003.
- [Hir05] Philip S. Hirschhorn. Overcategories and undercategories of model categories. <http://www-math.mit.edu/~psh/undercat.pdf>, 2005.
- [Hir18] Philip S. Hirschhorn. Errata for *Model categories and their localizations*. <http://www-math.mit.edu/~psh/MCATL-errata-2018-08-01.pdf>, 2018.
- [Joy02] A. Joyal. Quasi-categories and Kan complexes. *J. Pure Appl. Algebra*, 175(1-3):207–222, 2002. Special volume celebrating the 70th birthday of Professor Max Kelly.
- [Joy08a] André Joyal. Notes on quasi-categories. Preprint. <https://www.math.uchicago.edu/~may/IMA/Joyal.pdf>, 2008.
- [Joy08b] André Joyal. The theory of quasi-categories and its applications. Lecture notes. <http://mat.uab.cat/~kock/crm/hocat/advanced-course/Quadern45-2.pdf>, 2008.
- [JT07] André Joyal and Myles Tierney. Quasi-categories vs Segal spaces. In *Categories in algebra, geometry and mathematical physics*, volume 431 of *Contemp. Math.*, pages 277–326. Amer. Math. Soc., Providence, RI, 2007.
- [Lur09] Jacob Lurie. *Higher topos theory*, volume 170 of *Annals of Mathematics Studies*. Princeton University Press, Princeton, NJ, 2009.
- [Ree74] C. L. Reedy. Homotopy theory of model categories. <http://www-math.mit.edu/~psh/reedy.pdf>, 1974.
- [Ros17] J. Rosický. Accessible model categories. *Appl. Categ. Structures*, 25(2):187–196, 2017.
- [RSS01] Charles Rezk, Stefan Schwede, and Brooke Shipley. Simplicial structures on model categories and functors. *Amer. J. Math.*, 123(3):551–575, 2001.
- [RV14] Emily Riehl and Dominic Verity. The theory and practice of Reedy categories. *Theory Appl. Categ.*, 29:256–301, 2014.
- [RV17] Emily Riehl and Dominic Verity. Fibrations and Yoneda’s lemma in an  $\infty$ -cosmos. *J. Pure Appl. Algebra*, 221(3):499–564, 2017.
- [RV19] Emily Riehl and Dominic Verity. Elements of  $\infty$ -category theory. Book in progress, draft available at <http://www.math.jhu.edu/~eriehl/elements.pdf>, 2019.
- [SS88] Dietmar Schumacher and Ross Street. Some parametrized categorical concepts. *Comm. Algebra*, 16(11):2313–2347, 1988.
- [Ste17] Danny Stevenson. Covariant model structures and simplicial localization. *North-West. Eur. J. Math.*, 3:141–203, 2017.
- [Ste18a] Danny Stevenson. Model structures for correspondences and bifibrations. arXiv:1807.08226v1, 2018.
- [Ste18b] Danny Stevenson. Stability for inner fibrations revisited. *Theory Appl. Categ.*, 33:Paper No. 19, 523–536, 2018.
- [Str01] Ross Street. Powerful functors. <http://web.science.mq.edu.au/~street/Pow.fun.pdf>, 2001.
- [Str04] Ross Street. Cauchy characterization of enriched categories. *Repr. Theory Appl. Categ.*, (4):1–16, 2004.

CENTRE OF AUSTRALIAN CATEGORY THEORY, MACQUARIE UNIVERSITY, NSW 2109, AUSTRALIA  
 URL: <http://web.science.mq.edu.au/~alexc/>