



UNIVERSITY OF STRATHCLYDE

DEPARTMENT OF CHEMICAL & PROCESS ENGINEERING

M.ENG CHEMICAL & PROCESS ENGINEERING 18530

Computer-aided Design of Bio-inspired Nanoporous Silica Materials

Author:
André CRESCENZO

Supervisor:
Miguel JORGE

March 30, 2015

1 Summary

Contents

	Page
1 Summary	i
2 Acknowledgements	iii
3 Introduction	1
4 Experimental Methods	2
4.1 Theoretical Basis	2
4.2 Methods Description	3
4.3 Experiment 1	4
4.4 Experiment 2	5
4.5 Experiment 3	5
5 Results	5
6 Discussion	5
7 Conclusion	6
8 Nomenclature	6
9 References	6
10 Appendix	1

2 Acknowledgements

3 Introduction

Molecular Dynamics (MD) and Monte Carlo (MC) have been powerful tools to simulate molecular interactions of surfactants in solvent systems, allowing a deeper understanding of self-assembly process of this sort of material (Need & Someone, 2025). Therefore, these structural conformations are specially useful to design bio-inspired silica materials (Patwardhan, 2011) since with the addition of silica they remain behaving as scaffolds to mesoporous or nanoporous structures that are maintained even after surfactant removal, silica oligomer polymerization and calcination (Tolbert, Firouzi, Stucky, & Chmelka, 1997). A vast range of silica materials are examples of this phenomena, such as MCM-41 as reported by Kresge and Roth (2013), HR (Jin et al., 2008), MSU-V (Tanev, Liang, & Pinnavaia, 1997) and many others, in which the self-assembly structure depend on the type of surfactant and concentration of the substances involved. Moreover, most of the experimental methods used to obtain data are based on observation and interpretation of final silica structure by using X-ray diffraction (XRD) and transmission electron microscopy (TEM), thus initial self-assembled conformations are predicted as a reflex of the final results and little is known about the mechanistic of this process. However, by using MD simulations it is possible to observe and analyse these initial steps of self-assembly and further predict, with more accuracy, properties and framework provided by surfactant (Tresset, 2009) to generate silica structures.

A major concern is that even though MD uses sophisticated software prepared to simulate enormous systems with thousands of atoms, using all capacities of hardware available, such as high-speed multi-core processors in conjunction with GPUs designed specifically to process data from arrays (Pronk et al., 2013), they hardly can achieve long time horizons and are commonly limited to a few μs depending on the size of the system. For this reason, several techniques have been develop to optimize the performance of the simulations such as coarse-grain models (Need & Someone, 2025). Briefly, the idea behind using this technique is to fit parameters of heavy atoms groups with similar properties in a bigger "bead", which includes atomic masses and electrostatic charges lumped in approximated values. For example, by merging carbon heavy atoms, that means carbon atom and its hydrogen atoms, at a lipid tail in groups of three or four since they have similar hydrophobic properties.

In order to provide topologies to GROMACS simulations several force-fields, for exemple MARTINI (?, ?), use Lennard-Jones potentials fitted to a range of pre-defined bead types to describe coarse-grain models. Furthermore, not only intramolecular beads are possible, but also intermolecular beads can be specified, such as multiple solvent molecules merged in a single bead or ions surround by water molecules. Previous works conducted by (Pérez-Sánchez, Gomes, & Jorge, 2013) with this method recreated a model of surfactant in presence of silica with explicit water that was successful in describing rod-like self-assembly structures detected on MCM-41 materials, demonstrating the capacities of up-scaling this type of systems. Nevertheless, solvent presence demands most of the computational resources, hence implicit solvent scheme has been focus of many studies (Pronk et al., 2013).

Different concepts has been applied to develop a suitable model for implicit solvents, studies conducted by Arnarez et al. (2015) developed the Dry MARTINI force-field by modifying parameters of its predecessor, in such manner that solvent interactions became incorporated in these values and then solvent beads are no longer necessary. Another method, described by (Mirzoev & Lyubartsev, 2013) is able to recreate a implicit solvent system from interaction potentials generated from a bottom-up approach, that means by using an all-atoms simulation to generate parameters for the coarse-grain model. It is supposed that thermodynamic changes on the system, originated from solvent interaction with amphiphilic molecules, are incorporated on the approximated potentials. Therefore, changes in parameters as concentration may not affect coarse-grained model performance on recreating self-assembly structures (Mirzoev & Lyubartsev, 2014).

The work presented on following experiments are an attempt to create a cross-link method to upscale silica crystal-liquid phase interactions (Tolbert et al., 1997) from the atomistic model to a mesoscale model with the advent of this later coarse-grain technique. The methodology applied to reach the desired model is based on the MagiC software package (Mirzoev & Lyubartsev, 2013) that in conjunction MD simulation software, in this case GROMACS (Pronk et al., 2013), will provide a suitable approximation to self-assembly of amphiphilic molecules in presence of silica. Further explanations of the process are described on Experimental Methods section. For the scope of this project, a bolaamphiphilic molecule 1,12-diaminododecane (DMDD) has been chosen as surfactant because that as seen in previous research by Tanev et al. (1997) it self-assembly in multilamellar vesicles that in presence of a silica precursor is capable of generating a mesoporous structure with remarkable properties. In order to validate this structure formation and further framework formation for silica oligomers, a coarse-grain approximation is suitable option since amphiphilic molecules interaction with solvents can be described efficiently with tabulated Lennard-Jones potentials interactions. As a final

objective at the end of this project, a implicit water coarse-grain model for DMDD will be generated and properly validated based on MD simulations and thermodynamic properties, in order to provide a good approximation to interactions with silica oligomers in a mesoscale model.

4 Experimental Methods

4.1 Theoretical Basis

This work is based on two molecular simulation techniques the former, called Molecular Dynamics, accounts for integration of Newton forces in order to describe positions and velocities of particles in the system. The latter, called Monte Carlo method, is a powerful statistical tool based on stochastic inputs to measure properties of many systems, in this case thermodynamic equilibrium of molecular systems. Following sections will describe the mechanistic behind each technique and further explain how they are used in simulation software.

Molecular Dynamics simulation

Molecular Dynamics is a simulation method that originates from dynamic nature of atomic interactions. Generally, a system of atoms can be treated as a multi-particle system ruled by Newton's Law. Considering a system with \mathcal{N} atoms for each i th particle of the system the follow differential equation:

$$m_i \frac{d^2 \vec{r}_i(t)}{dt^2} = \vec{F}_i(t) \quad (1)$$

Where $\vec{r} = (x, y, z)$ is the position vector and $F = (F_x, F_y, F_z)$ are the force components. Therefore, for this \mathcal{N} particle system, it is necessary to solve $3\mathcal{N}$ differential equations in order to fully describe it analytically at any t . Since any molecular system involves an enormous number of molecules, solving these equations analytically is impracticable and simulation methods are necessary. The derivatives terms need to be numerically calculated and as soon as simulation efficiency is a major concern, a suitable approximation to the second derivative of position is the Taylor's expansion, taking the x coordinate as an example, for a given Δt is:

$$x(t + \Delta t) = x(t) + \Delta t \frac{dx(t)}{dt} + \frac{1}{2!} \Delta t^2 \frac{d^2 x(t)}{dt^2} + \frac{1}{3!} \Delta t^3 \frac{d^3 x(t)}{dt^3} + \dots \quad (2)$$

$$x(t - \Delta t) = x(t) - \Delta t \frac{dx(t)}{dt} + \frac{1}{2!} \Delta t^2 \frac{d^2 x(t)}{dt^2} - \frac{1}{3!} \Delta t^3 \frac{d^3 x(t)}{dt^3} + \dots \quad (3)$$

$$\frac{d^2 x(t)}{dt^2} = \frac{x(t + \Delta t) - 2x(t) + x(t - \Delta t)}{\Delta t^2} + \mathcal{O}(\Delta t^4) \quad (4)$$

Hence, by neglecting the error of order $\mathcal{O}(\Delta t^4)$, it is possible to use equations Eqn.(1) and Eqn.(4) to derive the called "Verlet method" where position and velocity vectors for each i th atom on next step ($t + \Delta t$) are calculated using previous and current step:

$$\vec{r}_i(t + \Delta t) = 2\vec{r}_i(t) - \vec{r}_i(t - \Delta t) + \frac{\Delta t^2}{m_i} \vec{F}_i(t) \quad (5)$$

$$= \frac{2\vec{r}_i(t)\vec{r}_i(t + \Delta t) - \vec{r}_i(t - \Delta t)}{\Delta t^2} + \frac{\Delta t^2}{m_i} \vec{F}_i(t) \quad (6)$$

Monte Carlo simulation

Monte Carlo is a simulation method based on statistical probability of existence of a system at thermodynamic equilibrium. This happens when free energy reaches a minimum by calculating the energy for each particle's microscopic state. Given a system of \mathcal{N} particles, temperature T and volume V , thus it can be denoted as a canonical ensemble, it is possible to assume that the free Helmholtz energy of the system is:

$$A = U - TS \quad (7)$$

Where S is the entropy and U is the internal energy, meaning that not only a minimum in free energy can arise from a minimum in total energy, but also an increase in entropy of the system can contribute to this energy minimization. A system can be described by a group of coordinates, in this case to give

a simplified idea assume a group of distance vectors $\lambda = (\vec{r}_1, \vec{r}_2, \dots, \vec{r}_N)$ from system origin. For an arbitrary λ the probability of a single particle of the system to statistically occupy that position is described by using a probability distribution function (Sato, 2011):

$$\rho(\lambda) = \frac{\exp\left(-\frac{U(\lambda)}{kT}\right)}{\int_V \dots \int_V \exp\left(-\frac{U(\lambda)}{kT}\right) d\vec{r}_1 d\vec{r}_2 \dots d\vec{r}_N} \quad (8)$$

Whether the system configuration is generated with considerable number of microstates that satisfies this probability, the final configuration would have a real physical meaning. But, as soon as it is almost impossible to define an analytical solution to this equation if N is too large another method became necessary to use Monte Carlo simulations in molecular dynamics. The Metropolis method (Metropolis, Rosenbluth, Rosenbluth, Teller, & Teller, 1953) allowed the use of Monte Carlo technique by introducing the following concept: given 2 different microstates, the probability to change from state 1 to 2 is defined by:

$$P_{1 \rightarrow 2} = \begin{cases} 1 & \text{for } \frac{\rho(\lambda_2)}{\rho(\lambda_1)} \geq 1 \\ \frac{\rho(\lambda_2)}{\rho(\lambda_1)} & \text{for } \frac{\rho(\lambda_2)}{\rho(\lambda_1)} < 1 \end{cases} \quad (9)$$

Therefore, the integral term in eq.(8) vanishes and it is clear to observe that when $U(\lambda_1) \geq U(\lambda_2)$ the system will certainly change to this new microstate since it has lower energy, however in the case of $U(\lambda_1) < U(\lambda_2)$ the system has a certain probability of changing to this new microstate indicating an increase of entropy in the system. So, by applying stochastic inputs for particle displacements and acceptance values for eq.(9), with a large number of Monte Carlo steps the system will eventually reach a minimum in free energy. Moreover, when this system reaches equilibrium it is possible to calculate accurately microstate dependent properties via ensemble averages. That means with n samples of Monte Carlo steps, the average of a ξ property can be obtained by:

$$\langle \xi \rangle = \sum_{i=1}^n \frac{\xi_i}{n} \quad (10)$$

It is important to notice that Monte Carlo simulations do not consider dynamic properties of the system such as kinetic energy, and therefore only Molecular Dynamics can account such things. Finally, now that both molecular simulation methods had been properly introduced it is possible to proceed to an overview of both simulation software used on this project. Together they can unite the benefits of both simulation methods, in order to recreate a suitable coarse-grain model for Molecular Dynamics.

- gromacs(van der Spoel et al., 2010)
- MPI and GPU

About MagiC

- General idea
- IBI + IMC
- Parallelism and sample size
- Convergence

4.2 Methods Description

The experimental process of this project is divided in 3 main parts: Firstly, develop a efficient coarse-grain model and validate its use. Secondly, analyse the relationship between concentration of all-atoms simulation and concentration of coarse-grained simulation. Then finally, create and reproduce a Silica surfactant mesoscale system.

In order to maintain standard parameters for all GROMACS simulations, all-atoms simulations that were used for coarse-grained reference were run in NPT ensemble. All simulation boxes were created using genbox (GROMACS package utility), first by inserting the desired number of surfactant molecules and then adding the solvent at the adequate ratio. The molecules were designed based on OPLS-AA forcefield (?, ?) parameters, therefore the surfactant DMDD was created with aid of co-workers and water type was TIP4P (?, ?). Moreover, no initial configuration was determined, such as a pre self-assembled system, to ensure stochastic nature of the system.

Initially, all boxes received a energy minimization step in order to avoid system blow-up deal to possible extreme potential energy spots created during box generation, because due to random displacement of molecules some of them could overlap others and then cause a destabilization of the system. After, an equilibration step was necessary to reach desired initial temperature and pressure conditions for the MD simulation. During equilibration temperature coupling was kept as 298 *K* using a v-rescaling thermostat (?, ?) with time constant of 0.01 *ps* and pressure coupling was kept at 1 *bar* using a Berendsen barostat (?, ?) with time constant of 0.5 *ps*.

4.3 Experiment 1

–Objective




Figure 1: A) Schematic of M1 molecule split plan, Carbon atoms represented in black, Nitrogen in green and Hydrogen in gray. B) Final coarse-grain model for M1. This model aggregated the three most polar heavy atoms on the tips and four apolar heavy atoms at the center. In order to avoid electrostatic interactions on the IMC process and further simulation the heavy atoms groups (heavy atom + hydrogen) were kept thus final charge remain unchanged in each bead




Figure 2: A) Schematic of M2 molecule split plan, Carbon atoms represented in black, Nitrogen in green and Hydrogen in gray. B) Final coarse-grain model for M2. Heavy atom were grouped pairwise in such a manner that amphiphilic parts were kept as distinct as possible. It is important to notice the necessity of two carbon bead types CE and CI

–Model 1 vs Model 2

–gromacs Simulation Set-up

–MagiC CG M1 and M2 topology generation+Ref RDFs

–IMC process

–gromacs Set-up Reproduction Tests

4.4 Experiment 2

- Objective
- gromacs Set-up Up-scaling
- gromacs Set-up Self-assembly Model X(best)
- AA Low, CG Low
 - 1k(?) up-scale @ low
 - 1k(?) up-scale @ high
- AA Ideal, CG ideal
 - 1k(?) up-scale @ low
 - 1k(?) up-scale @ ideal
 - 1k(?) up-scale @ high
- AA High, CG High
 - 1k(?) up-scale @ high
 - 1k(?) up-scale @ high

4.5 Experiment 3

- Silica addition
- MagiC CG silica topology
- gromacs Set-up Self-assembly Model X + Si
 - 1000, 10000 (Maybe more?)

5 Results

- Experiment 1
 - Model 1 vs Model 2
 - gromacs Simulation
 - IMC process Convergence
 - gromacs Reproduction Tests RDFs and Properties
- Experiment 2
 - Self-assembly vs concentration Model X
 - Ideal con validation: Low, Ideal, High con comparison
 - Ideal @ Low -> validate
 - Ideal @ Ideal -> evaluate
 - Ideal @ High -> validate
 - Extremes evaluation:
 - Low @ Low vs High @ Low -> compare
 - High @ High vs Low @ High -> compare
- Experiment 3
 - Silica addition
 - CG silica Properties
 - Self-assembly Model X + Si
 - 1000, 10000 (Maybe more?)

6 Discussion

- Experiment 1
 - Which is the best Model?
 - Why Model X is better than Model Y?
 - To what extent Model X represent well the AA model?
 - What are the limitations and advantages of Model X?
- Experiment 2
 - What is the relation between the entropy "error" and concentration?
 - Is the concentration really an problem?
 - Do you need to run an AA for each con or the Ideal can represent any?
 - Even extreme cases can be approximated? In this case how much is the "error"?
- Experiment 3

- Is The silica CG model suitable?
- Can you the proprieties validate it?
- The self-assembly behaviour change with silica addition?

7 Conclusion

What did you really see from the results?
 There was any bad assumption that is contestable?
 What will be the next step to your research?

8 Nomenclature

H	Hamiltonian hdsdgfag (Mirzoev & Lyubartsev, 2013)a asdfasdfas dfadsfasdfasdfsdfa gsdflhg as djfg ajdf jasdfj adjhg asdjhgafajksdfaj,jahdskjfas
C_n	Molar concentration of species n (mM)

9 References

- Arnarez, C., Uusitalo, J. J., Masman, M. F., Ingólfsson, H. I., de Jong, D. H., Melo, M. N., ... Marrink, S. J. (2015). Dry martini, a coarse-grained force field for lipid membrane simulations with implicit solvent. *Journal of Chemical Theory and Computation*, 11(1), 260-275. Retrieved from <http://dx.doi.org/10.1021/ct500477k> doi: 10.1021/ct500477k
- Jin, H., Qiu, H., Sakamoto, Y., Shu, P., Terasaki, O., & Che, S. (2008). Mesoporous silicas by self-assembly of lipid molecules: Ribbon, hollow sphere, and chiral materials. *Chemistry – A European Journal*, 14(21), 6413–6420. Retrieved from <http://dx.doi.org/10.1002/chem.200701988> doi: 10.1002/chem.200701988
- Kresge, C. T., & Roth, W. J. (2013). The discovery of mesoporous molecular sieves from the twenty year perspective. *Chem. Soc. Rev.*, 42, 3663-3670. Retrieved from <http://dx.doi.org/10.1039/C3CS60016E> doi: 10.1039/C3CS60016E
- Metropolis, N., Rosenbluth, A. W., Rosenbluth, M. N., Teller, A. H., & Teller, E. (1953). Equation of state calculations by fast computing machines. *The Journal of Chemical Physics*, 21(6), 1087-1092. Retrieved from <http://scitation.aip.org/content/aip/journal/jcp/21/6/10.1063/1.1699114> doi: <http://dx.doi.org/10.1063/1.1699114>
- Mirzoev, A., & Lyubartsev, A. P. (2013). Magic: Software package for multiscale modeling. *Journal of Chemical Theory and Computation*, 9(3), 1512-1520. Retrieved from <http://dx.doi.org/10.1021/ct301019v> doi: 10.1021/ct301019v
- Mirzoev, A., & Lyubartsev, A. P. (2014). Systematic implicit solvent coarse graining of dimyristoylphosphatidylcholine lipids. *Journal of Computational Chemistry*, 35(16), 1208–1218. Retrieved from <http://dx.doi.org/10.1002/jcc.23610> doi: 10.1002/jcc.23610
- Need, A., & Someone, T. C. (2025). Mock article: You need to read more. *Journal of Void Theory*, 1(1), 151-1510.
- Patwardhan, S. V. (2011). Biomimetic and bioinspired silica: recent developments and applications. *Chem. Commun.*, 47, 7567-7582. Retrieved from <http://dx.doi.org/10.1039/C0CC05648K> doi: 10.1039/C0CC05648K
- Pronk, S., Páll, S., Schulz, R., Larsson, P., Bjelkmar, P., Apostolov, R., ... Lindahl, E. (2013). Gromacs 4.5: a high-throughput and highly parallel open source molecular simulation toolkit. *Bioinformatics*, 29(7), 845-854. Retrieved from <http://bioinformatics.oxfordjournals.org/content/29/7/845.abstract> doi: 10.1093/bioinformatics/btt055
- Pérez-Sánchez, G., Gomes, J. R. B., & Jorge, M. (2013). Modeling self-assembly of silica/surfactant mesostructures in the templated synthesis of nanoporous solids. *Langmuir*, 29(7), 2387-2396. Retrieved from <http://dx.doi.org/10.1021/la3046274> (PMID: 23343439) doi: 10.1021/la3046274
- Satoh, A. (2011). 1 - outline of molecular simulation and microsimulation methods. In A. Satoh (Ed.), *Introduction to practice of molecular simulation* (p. 1 - 27). London: Elsevier. Retrieved from <http://www.sciencedirect.com/science/article/pii/B978012385148200001X> doi: <http://dx.doi.org/10.1016/B978-0-12-385148-2.00001-X>

- Tanev, P. T., Liang, Y., & Pinnavaia, T. J. (1997). Assembly of mesoporous lamellar silicas with hierarchical particle architectures. *Journal of the American Chemical Society*, 119(37), 8616-8624. Retrieved from <http://dx.doi.org/10.1021/ja970228v> doi: 10.1021/ja970228v
- Tolbert, S. H., Firouzi, A., Stucky, G. D., & Chmelka, B. F. (1997). Magnetic field alignment of ordered silicate-surfactant composites and mesoporous silica. *Science*, 278(5336), 264-268. Retrieved from <http://www.jstor.org/stable/2894713> doi: 10.2307/2894713
- Tresset, G. (2009). The multiple faces of self-assembled lipidic systems. *PMC Biophysics*, 2(1). Retrieved from <http://dx.doi.org/10.1186/1757-5036-2-3> doi: 10.1186/1757-5036-2-3
- van der Spoel, D., Lindahl, E., Hess, B., van Buuren, A. R., Apol, E., Meulenhoff, P. J., . . . Berendsen, H. J. C. (2010). Gromacs user manual version 4.5.6 [Computer software manual].

10 Appendix