

Computer-aided Design of Bio-inspired Nanoporous Silica Materials

By André Crescenzo

Miguel Jorge

Supervisor

Department of Chemical
and Process Engineering

Alessia Centi


Supervisor

Department of Chemical
and Process Engineering

Carlos F. Rangel

Supervisor


Department of Chemical
and Process Engineering



Magic



GROMACS



Silica-surfactant materials

Highly Structured Materials

High Porosity
High Surface area

Generated by


Synthesis



MCM-50




MCM-48




MCM-41

Nature




MSU-V

1,2-diaminododecane



Self-Assembly structure




Molecular simulation

Molecular Dynamics

Newton's Law

$$m_i \frac{d^2 \vec{r}_i(t)}{dt^2} = \vec{F}_i(t)$$



“Leap-frog” Algorithm


$$\vec{r}_i(t + \Delta t) = \vec{r}_i(t) + \vec{v}_i(t + \Delta t/2) \Delta t$$

$$\vec{v}_i(t + \Delta t/2) = \vec{v}_i(t - \Delta t/2) + \frac{\vec{F}_i(t)}{m_i} \Delta t$$

Monte Carlo Simulations

Probability distribution function

$$\rho(\lambda) = \frac{\exp\left(-\frac{U(\lambda)}{kT}\right)}{\int_V \dots \int_V \exp\left(-\frac{U(\lambda)}{kT}\right) d\vec{r}_1 d\vec{r}_2 \dots d\vec{r}_N}$$




Metropolis Method


$$P_{1 \leftrightarrow 2} = \begin{cases} 1 & \text{for } \frac{\rho(\lambda_2)}{\rho(\lambda_1)} \geq 1 \\ \frac{\rho(\lambda_2)}{\rho(\lambda_1)} & \text{for } \frac{\rho(\lambda_2)}{\rho(\lambda_1)} < 1 \end{cases}$$

Coarse-graining


What is
coarse-grain?



Simulation Time
x
Computational limits




Time = ~1,5 days




Time = ~10 minutes

Tailored Potentials




MagiC: a systematic method

- Boltzmann inversion
- Inverse Monte Carlo



Atomistic Model



- Concentration
- pH
- Equilibrium Structure
- Standardized simulations
 - NPT
 - Over 100 ns

CG Modelling: Bead Size and Concentration

- Multiple Models

- Model 1:



- Model 2:




- Multiple Concentrations

- Low

- Medium

- High

CG Modelling: Charged Systems

- New Techniques for process efficiency
 - Simplified model
 - 
 - Multi-states reference
 - Changes on MagiC inputs
- Influence of Electrostatic forces
 - Integrated Electrostatics
 - Short range Electrostatics
 - Long range Electrostatics

CG Modelling: Silica introduction

- Ionic silica with neutral surfactant
- Ionic silica with singly charged surfactant

Conclusion

- (general conclusion)
- (still need to think)
- Ideas for the future
 - Artificial CG model
 - Multi-state IBI
 - More focus in pH value