RELAXING THE COLOUR-REFINEMENT CONDITION IN OPENAI'S SIDORENKO COUNTEREXAMPLE: A CONDITIONAL 17-VERTEX, 30-EDGE INSTANCE

[AUTHOR NAME]

Abstract. OpenAI's preprint *A counterexample to Sidorenko's conjecture* uses the point–face incidence graph (35 vertices, 66 edges) of a 13-point triangulated sphere satisfying four conditions (1)–(4). Condition (4), discrete colour refinement, is used in exactly one place: the pinning lemma. We show that pinning still works when the stable colouring is not discrete, provided every colour-preserving homomorphism is an automorphism. The reason is that the coefficient C is invariant under automorphisms. A 7-point, 10-face triangulation satisfies (1)–(3) and this weaker condition. Its incidence graph H_7 has 17 vertices and 30 edges. **Conditionally** on the paper's rank-layer lemma (and its proof in Sections 4–7) holding for this complex with the same argument, H_7 is a counterexample to Sidorenko's conjecture. This conditional transfer is made at the level of the TEX argument only and has no machine certificate. It is therefore **less certain than the original**, whose 35-vertex statement is formalised in Lean.

Attribution. The construction and its proof are OpenAI's work (family 161 of `https://github.com/openai/math`, commit `fd4aeeb2ee4fc729c18d98444fed42fd0529eeeb`, [1]). This note changes only the choice of the finite complex and one lemma (pinning). It inherits every other step. Developed with AI assistance (Anthropic Claude, Opus 5.5) under [AUTHOR NAME]'s direction; independent adversarial reviews were run by separate AI sessions and are not human peer review.

Premise and confidence.

(P) Lemma 3.1 of [1] (rank-layer limit) and its proof in Sections 4–7 hold for the complex K_7 below, with the complex-dependent constants re-instantiated (Section 4). This is *assumed*, not re-proved line by line.

- The official Lean development certifies only the 13-point/22-face complex (index types are fixed as `Fin 13`, `Fin 22`, `Fin 33`). Our instance has no machine certificate. Its reliability is lower than that of the original theorem.
- Unconditional and exactly checked: (1)–(3) for K_7; the colour-preserving homomorphisms of H_7; the coefficient values; the generalised pinning lemma (a short proof, Section 3).

1. Statement

Let K_7 be the complex on points $\{0,\dots,6\}$ with the ten faces in Table 1, and let $H_7 = (I \sqcup J, \mathscr{C})$ be its incidence graph: $|I| = 7$, $|J| = 10$, $|\mathscr{C}| = 30$.

Theorem 1.1. *Assume (P). There is a finite simple graph G with $t(H_7, G) < p(G)^{30}$, where $p(G)$ is the edge density.*

The original graph has 35 vertices and 66 edges [1, Thm. 1.1].

2. Conditions (1)–(3) hold; (4) fails

Proposition 2.1. *K_7 satisfies conditions (1)–(3) of* [1, Prop. 2.1]*, with* $11, 13$ *replaced by* $5, 7$*:*

Date: October 9, 2026.

face j	points	opposite faces on $\{i_1i_2\},\{i_1i_3\},\{i_2i_3\}$	bits
0	0 1 2	1, 2, 6	000
1	0 1 6	0, 5, 7	001
2	0 2 3	0, 3, 8	001
3	0 3 4	2, 4, 8	011
4	0 4 5	3, 5, 9	111
5	0 5 6	4, 1, 7	011
6	1 2 5	0, 7, 9	100
7	1 5 6	6, 1, 5	110
8	2 3 4	2, 9, 3	100
9	2 4 5	8, 6, 4	110

TABLE 1. The complex K_7. The edges of H_7 are the 30 pairs (point, face containing it).

(1) *each of the* $|E| = 15$ *point pairs lies in exactly two faces, and the face-neighbour graph is connected (Euler characteristic* $7 - 15 + 10 = 2$*);*
(2) *for each bit position, each class has* 5 *faces covering all* 7 *points, and it has an exposure order in which every later face shares a pair with an earlier face and adds exactly one new point:*

bit $1 = 0$:	$0,1,2,3,5$	(*new* $6,3,4,5$)	*bit* $1 = 1$:	$4,9,6,7,8$	$(2,1,6,3)$
bit $2 = 0$:	$0,1,2,6,8$	$(6,3,5,4)$	*bit* $2 = 1$:	$3,4,5,7,9$	$(5,6,1,2)$
bit $3 = 0$:	$0,6,7,9,8$	$(5,6,4,3)$	*bit* $3 = 1$:	$1,5,4,3,2$	$(5,4,3,2)$;

(3) *every* $a_e \geq 1/3$*, and* $a_{\{1,6\}} = a_{\{3,4\}} = 1$*,* $a_{\{2,3\}} = a_{\{5,6\}} = 2/3$ *(all other* $a_e = 1/3$*).*

Condition (4) fails. The stable colouring has point classes $\{0\},\{1,4\},\{2,5\},\{3,6\}$ *and face classes* $\{012,045\},\{016,034\},\{023,056\},\{125,245\},\{156,234\}$*. Moreover* $\mathrm{Aut}(H_7) = \{\mathrm{id},\varphi\}$*, where* φ *acts on points by* $(1\,4)(2\,5)(3\,6)$ *and on faces by* $j \mapsto (4,3,5,1,0,2,9,8,7,6)_j$*.*

Proof. Direct verification from Table 1. Two independent scripts check it exactly (`certificate/check_v7.py`, `construction/verify3.py`). □

3. Pinning without discreteness

We use the notation of [1, Section 3]: base laws A^{base} with corner signs σ_κ (exactly one negative), coefficients $C(a,b)$ of [1, (3.9)], and the scores $\mathcal{Q}$ of [1, Lemma 3.4].

Lemma 3.1 (Pinning with ties)**.** *Let* $F = (P \sqcup Q, L)$ *be a finite bipartite graph, and run colour refinement from the two part colours until it is stable (round* R*). Define the scores and* $\mathcal{Q}$ *exactly as in the proof of* [1, Lemma 3.4]*. Then for every part-preserving map* f *we have* $\mathcal{Q}(f) \leq \mathcal{Q}(\mathrm{id})$*, with equality if and only if* f *preserves the stable colouring.*

Proof. The proof of [1, Lemma 3.4] shows two things. If f preserves the colours of round $t-1$, then $F_t(\mathrm{id}) - F_t(f) = \sum_u \|h_t(f(u)) - h_t(u)\|^2 \geq 0$, with equality if and only if f also preserves the round-t colours. And at the first round where a colour is not preserved, the loss outweighs all later gains. Discreteness was used only to conclude that every $f \neq \mathrm{id}$ has such a round. Without it, the maps with no such round are exactly the maps preserving the stable colouring, and they tie with the identity in every round. □

Lemma 3.2. *If a part-preserving map* $f = (a,b)$ *is not a homomorphism* $H \to H$*, then* $C(a,b) = 0$*.*

Proof. Some corner (i,j) is sent to a non-incident type pair $(a(i),b(j))$. There the activation law is $S = \varnothing$ [1, Section 3.3], so $A_{a(i)b(j)}(\alpha,\beta) = 0$ unless $\alpha = \beta = \varnothing$. Every term in [1, (3.9)] has all labels nonempty. □

Lemma 3.3 (Automorphism invariance)**.** *For every automorphism φ of H (preserving the parts), $C(\varphi) = \big(\prod_\kappa \sigma_\kappa\big)(\theta/2)^{|\mathscr{C}|} = C(\mathrm{id})$.*

Proof. An automorphism permutes corners, pairs and faces compatibly, so $\varphi(U_e) = U_{\varphi(e)}$, $\varphi(M_\kappa) = M_{\varphi(\kappa)}$ and $\varphi(\Gamma_\kappa) = \Gamma_{\varphi(\kappa)}$. Relabelling the coordinate set by φ^{-1} gives $A^{\mathrm{base},\sigma}_{\varphi(\kappa)}(\alpha,\beta) = A^{\mathrm{base},\sigma\circ\varphi}_{\kappa}(\varphi^{-1}\alpha, \varphi^{-1}\beta)$, and hence $C_\sigma(\varphi) = C_{\sigma\circ\varphi}(\mathrm{id})$. The support-forcing Lemma 3.3 of [1] holds for every sign vector (its proof uses only (1) and the structure of M_κ, Γ_κ), with value $\prod_\kappa (\sigma\circ\varphi)_\kappa\,(\theta/2)^{|\mathscr{C}|}$. The product of the signs is unchanged by the permutation. □

Proposition 3.4. *The colour-preserving part-preserving maps of H_7 number $2^6 \cdot 2^{10} = 65536$. Exactly two of them are homomorphisms, namely* id *and φ.*

Proof. Exhaustive enumeration. Three implementations agree: `construction/verify3.py` (all 65536 maps), `construction/pinexact.py` (variable elimination over all part-preserving maps and over all homomorphisms), and `certificate/check_v7.py` (written independently from the TEX: 32033 point maps extend to homomorphisms; exactly two maximise $\mathcal{Q}$, with exact $C = -(\theta/2)^{30}$ each). □

Proof of Theorem 1.1. In [1, (3.21)], replace $13, 22, 66$ by $7, 10, 30$. By Lemmas 3.1–3.3 and Proposition 3.4, the maximal-score part of the sum is $C(\mathrm{id}) + C(\varphi) = -2(\theta/2)^{30}$. Every other map has score at most $\mathcal{Q}_* - \delta_*$ with $\delta_* > 0$. Choosing ρ with $B_* e^{-\rho\delta_*} < 2(\theta/2)^{30}$ gives $t_0 < 0$. Dilution gives $t(W_\varepsilon) = 1 + \varepsilon^7 t_0 + o(1)$ and $t(W_\varepsilon^\top) = 1 + \varepsilon^{10} t_1 + o(1)$, so the bracket of [1, (3.25)] becomes $t_0 + \varepsilon^3 t_1 + \varepsilon^{10} t_0 t_1$. The crossed product and the finite-host transfer [1, Prop. 3.5, Lemma 3.6] apply unchanged. All of these steps use Lemma 3.1, hence (P). □

4. What the premise covers

In [1], condition (4) is cited only in the application of the pinning lemma (`kernel.tex` l. 455). Sections 4–7 use (1)–(3) and numbers derived from the complex. Re-instantiated for K_7, these are: $33 \to 15$ pairs; $22 \to 10$ faces in the slack bound [1, §7] ($22C_* \to 10C_*$, so $D > M_* + 20C_*$ suffices there); point degrees $(6,4,5,3,4,5,3)$, so $\sum_i \binom{2\deg i}{2} = 242$ (was 618); and the profile exponent bound $m^2 - 4m + 242 \le 407$ for $m \le 15$ (was 1575). A separate review session checked these substitutions. We did *not* re-derive Sections 4–7 for K_7; this is the content of (P).

5. Controls: smaller complexes

Removing (4), every sphere triangulation with $V \le 6$ satisfying (2)–(3) fails for a definite reason.

- $V = 4, 5$ and the octahedron ($V = 6$, #2): H is already a Sidorenko graph. A reflection (an involution of H splitting the faces into two halves glued along fixed points) and the Cauchy–Schwarz inequality reduce H to a half with a part of size ≤ 4 (or a dominating vertex), which is Sidorenko [2, 3] (`construction/survey.py`). The tetrahedron's H has a part of size 4.
- $V = 6$, #1 (faces $012, 015, 023, 034, 045, 124, 145, 234$): the pinning step fails. Eight colour-preserving *folding* homomorphisms have $C = +\theta^{22}/2^{24}$, which dominates $C(\mathrm{aut}) = -\theta^{24}/2^{24}$ for small θ. They form pairs with $\mathrm{vec}(f) + \mathrm{vec}(f') = 2\,\mathrm{vec}(\mathrm{id})$ (counts of point, face and corner images), so by AM–GM no choice of priors and thinning weights the identity above both (`construction/fold1.py`). Whether this H is Sidorenko is not decided here.

At $V = 7$ the other two complexes behave differently. #4 is Sidorenko by reflection. For #5, generalised pinning is inconclusive (270 maximising homomorphisms against $|\mathrm{Aut}| = 6$). So K_7 (#3) is the smallest instance in the relaxed template for which this argument goes through. The same argument also works for $V = 8$ (#6) and $V = 9$ (#10).

6. H_7 is not in a known Sidorenko class

class (positive theorem)	why H_7 is not covered
trees, even cycles, complete bipartite; one part of size ≤ 4 [2]	parts $7, 10$; cycles; min degree 3
vertex complete to the other part [3]	max degrees $6 < 10$ and $3 < 7$
weakly norming (needs regular parts, [4])	point degrees $3, 3, 4, 4, 5, 5, 6$
Coregliano's degree-count criterion [7]	$10 < \binom{7}{3} = 35$; reverse orientation also fails
blow-ups [6]	no twin vertices
Cartesian products	$\lvert V(H_7)\rvert = 17$ is prime
subdivisions / even-path substitutions	min degree 3
reflection (symmetric gluing)	Aut $= \mathbb{Z}/2$; φ fixes only point 0, and $H_7 - 0$ is connected
strongly tree-decomposable [5], reflection trees, tree-arrangeable	first-step leaf test: 56 candidate leaf sets, none passes (`classes/stdcheck.py`)
Szegedy's "thick" class	**not decided**

TABLE 2. Location of H_7; invariants in `classes/invariants.py`. Conlon–Lee's further sufficient conditions (divisibility, $|B| \geq \binom{|A|}{r}$) were checked by the review session to fail as well.

Also, a numerical search over 69 step-kernel starts (`wsearch`) always converged to the constant kernel. The review session reports the same behaviour for the original graph: the counterexample kernel is a high-dimensional rank-layer kernel, not a small step function.

7. Relation to other work

The public repository `rohanarun/sidorenko-template-refinements` (2026-10-09, commit `4e1b9b6`) keeps condition (4) and proves that, within that template, the 13-point/35-vertex complex is minimal. For $V \leq 12$ they give exact cut-obstruction certificates. Their crossing-sum identity $\sum_{e \ni v} a_e = 2$ forces point degree ≤ 6, and K_7 attains this ($a_e = 1/3$ on all six pairs at point 0). Relaxing (4) is exactly what takes us past their template minimum. Appendix A is an independent recomputation of their $V \leq 10$ range.

Appendix A. Independent LP certificates with (4) kept

Lemma A.1. *For a sphere triangulation, the admissible bit splits (both classes have spanning exposure orders) correspond bijectively, up to swapping the classes, to Hamiltonian cycles of the point graph. The cycle is the set of pairs separated by the split.*

Proof. An exposure order glues each new triangle along a boundary edge of a disc and adds a new vertex. Each class is therefore a triangulated disc with all V points on its boundary, and the common boundary is a Hamiltonian cycle. Conversely, a Hamiltonian cycle bounds two discs with no interior vertices (Jordan). Reversing an ear decomposition of each disc gives exposure orders. □

Fractional (2)+(3) asks for a probability vector w on Hamiltonian cycles covering every pair with mass $\geq 1/3$. A rational certificate $y \geq 0$ with $\sum_e y_e = 1$ and $y(C) < 1/3$ for every Hamiltonian cycle C rules this out. `appendix_lp/r6_verify.py` enumerates all triangulations with $V \leq 10$ (plantri 5.8) and finds the discrete ones: $0, 0, 0, 0, 2, 16, 137$ for $V = 4, \ldots, 10$. It certifies all 155 in `fractions.Fraction` against every Hamiltonian cycle; the worst certified bounds are $1/5, 2/7, 2/7$ at $V = 8, 9, 10$. As a positive control, the original complex is feasible with $t^* = 1/3$. Restricting to point degree ≤ 6 gives $0, 1, 4$ at $V = 8, 9, 10$, the same as in `rohanarun`. An earlier run (`primary/`,

plantri 5.5) covered $V \leq 12$ (7881 discrete complexes, all infeasible with exact certificates). Its $V = 11, 12$ part has not been re-run independently.

References

[1] OpenAI, *A counterexample to Sidorenko's conjecture*, OpenAI Math Release preprint, September 23, 2026, https://github.com/openai/math/tree/main/preprints/A-counterexample-to-Sidorenkos-conjecture-September-23-2026.

[2] A. Sidorenko, Inequalities for functionals generated by bipartite graphs, Diskret. Mat. 3 (1991); and A correlation inequality for bipartite graphs, Graphs Combin. 9 (1993).

[3] D. Conlon, J. Fox, B. Sudakov, An approximate version of Sidorenko's conjecture, GAFA 20 (2010).

[4] H. Hatami, Graph norms and Sidorenko's conjecture, Israel J. Math. 175 (2010).

[5] D. Conlon, J. H. Kim, C. Lee, J. Lee, Some advances on Sidorenko's conjecture, J. London Math. Soc. 98 (2018).

[6] D. Conlon, J. Lee, Sidorenko's conjecture for blow-ups, Discrete Anal. (2021).

[7] L. N. Coregliano, as cited in [1, §1] (Theorem 3.6 there).