

# Probabilistic perspective of Machine Learning

András Attila Sulyok

Pázmány Péter Catholic University  
Faculty of Information Technology and Bionics


2024-04-09



- 1 Probabilistic perspective
- 2 Maximum Likelihood Estimation
- 3 Maximum A Posteriori Estimation

# Probabilistic model

Supervised learning:



Likelihood (of the parameters):  $P(\mathcal{D} \mid w) = P(Y \mid X, w)P(X)$

- 1 Probabilistic perspective
- 2 Maximum Likelihood Estimation
- 3 Maximum A Posteriori Estimation

# Maximum Likelihood Estimation

$$\arg \max_w P(\mathcal{D} \mid w) = \arg \max_w \{P(Y \mid X, w)P(X)\}$$

Assumption: samples are independent and identically distributed.  
Assumption: distribution of  $x$  is uniform

# Maximum Likelihood Estimation

$$\begin{aligned}\arg \max_w P(\mathcal{D} \mid w) &= \arg \max_w \{P(Y \mid X, w)P(X)\} \\ &= \arg \max_w P(Y \mid X, w)\end{aligned}$$

Assumption: samples are independent and identically distributed.

Assumption: distribution of  $x$  is uniform

# Maximum Likelihood Estimation

$$\begin{aligned}\arg \max_w P(\mathcal{D} \mid w) &= \arg \max_w \{P(Y \mid X, w)P(X)\} \\ &= \arg \max_w P(Y \mid X, w) \\ &= \arg \max_w P(y_1, \dots, y_N \mid x_1, \dots, x_N, w)\end{aligned}$$

Assumption: samples are independent and identically distributed.

Assumption: distribution of  $x$  is uniform

# Maximum Likelihood Estimation

$$\begin{aligned}\arg \max_w P(\mathcal{D} \mid w) &= \arg \max_w \{P(Y \mid X, w)P(X)\} \\&= \arg \max_w P(Y \mid X, w) \\&= \arg \max_w P(y_1, \dots, y_N \mid x_1, \dots, x_N, w) \\&= \arg \max_w \prod_{i=1}^N P(y_i \mid x_i, w)\end{aligned}\quad \text{ys are i.i.d}$$

Assumption: samples are independent and identically distributed.

Assumption: distribution of  $x$  is uniform



# Maximum Likelihood Estimation

$$\begin{aligned}\arg \max_w P(\mathcal{D} \mid w) &= \arg \max_w \{P(Y \mid X, w)P(X)\} \\&= \arg \max_w P(Y \mid X, w) \\&= \arg \max_w P(y_1, \dots, y_N \mid x_1, \dots, x_N, w) \\&= \arg \max_w \prod_{i=1}^N P(y_i \mid x_i, w) \quad \text{ys are i.i.d} \\&\text{(some algebraic magic follows)} \\&= \arg \max_w \log \prod_{i=1}^N P(y_i \mid x_i, w)\end{aligned}$$

Assumption: samples are **independent and identically distributed**.

Assumption: distribution of  $x$  is uniform

# Maximum Likelihood Estimation

$$\arg \max_w P(\mathcal{D} \mid w) = \arg \max_w \log \prod_{i=1}^N P(y_i \mid x_i, w)$$

Assumption: samples are independent and identically distributed.

Assumption: distribution of  $x$  is uniform

# Maximum Likelihood Estimation

$$\begin{aligned}\arg \max_w P(\mathcal{D} \mid w) &= \arg \max_w \log \prod_{i=1}^N P(y_i \mid x_i, w) \\ &= \arg \max_w \sum_{i=1}^N \log P(y_i \mid x_i, w)\end{aligned}$$

Assumption: samples are independent and identically distributed.

Assumption: distribution of  $x$  is uniform

# Maximum Likelihood Estimation

$$\begin{aligned}\arg \max_w P(\mathcal{D} \mid w) &= \arg \max_w \log \prod_{i=1}^N P(y_i \mid x_i, w) \\ &= \arg \max_w \sum_{i=1}^N \log P(y_i \mid x_i, w) \\ &= \arg \min_w \sum_{i=1}^N -\log P(y_i \mid x_i, w)\end{aligned}$$

Assumption: samples are independent and identically distributed.

Assumption: distribution of  $x$  is uniform

# Maximum Likelihood Estimation

$$\begin{aligned}\arg \max_w P(\mathcal{D} \mid w) &= \arg \max_w \log \prod_{i=1}^N P(y_i \mid x_i, w) \\&= \arg \max_w \sum_{i=1}^N \log P(y_i \mid x_i, w) \\&= \arg \min_w \sum_{i=1}^N -\log P(y_i \mid x_i, w) \\&= \arg \min_w \mathbb{E}_{(x,y) \sim \mathcal{D}} -\log P(y \mid x, w)\end{aligned}$$

Assumption: samples are independent and identically distributed.

Assumption: distribution of  $x$  is uniform

# Example: logistic regression

Remember the likelihood for the logistic regression:

$$P(y = + \mid x, w) = \theta(w^T x)$$

( $\theta$  is the logistic sigmoid)

# Example: logistic regression

Remember the likelihood for the logistic regression:

$$P(y = + \mid x, w) = \theta(w^T x)$$

( $\theta$  is the logistic sigmoid)

Binary cross entropy error (for one sample point):

$$\begin{cases} -\log(1 - \theta(w^T x)) & \text{if } y = - \\ -\log \theta(w^T x) & \text{if } y = + \end{cases}$$

# Example: linear regression

Likelihood:

$$P(y \mid x, w) = \mathcal{N}(y; w^T x, \sigma^2) = \frac{1}{\sqrt{2\pi\sigma^2}} \exp\left(-\frac{(y - w^T x)^2}{2\sigma^2}\right)$$



# Example: linear regression

Likelihood:

$$P(y \mid x, w) = \mathcal{N}(y; w^T x, \sigma^2) = \frac{1}{\sqrt{2\pi\sigma^2}} \exp \left( -\frac{(y - w^T x)^2}{2\sigma^2} \right)$$

Cross entropy error for linear regression:

$$\begin{aligned} \arg \min_w -\log P(y \mid x, w) &= \arg \min_w \frac{1}{2} \log (2\pi\sigma^2) + \left( \frac{(y - w^T x)^2}{2\sigma^2} \right) \\ &= \arg \min_w (y - w^T x)^2 \end{aligned}$$

Looks familiar?

- 1 Probabilistic perspective
- 2 Maximum Likelihood Estimation
- 3 Maximum A Posteriori Estimation

# Maximum A Posteriori Estimation

ML objective was:

$$\arg \max_w P(\mathcal{D} \mid w)$$

# Maximum A Posteriori Estimation

ML objective was:

$$\arg \max_w P(\mathcal{D} \mid w)$$

Alternative objective:

$$\arg \max_w P(w \mid \mathcal{D})$$

# Maximum A Posteriori Estimation

ML objective was:

$$\arg \max_w P(\mathcal{D} \mid w)$$

Alternative objective:

$$\arg \max_w P(w \mid \mathcal{D}) = \arg \max_w \frac{P(\mathcal{D} \mid w)P(w)}{P(\mathcal{D})}$$

# Maximum A Posteriori Estimation

ML objective was:

$$\arg \max_w P(\mathcal{D} \mid w)$$

Alternative objective:

$$\arg \max_w P(w \mid \mathcal{D}) = \arg \max_w \frac{P(\mathcal{D} \mid w)P(w)}{P(\mathcal{D})} = \arg \max_w P(\mathcal{D} \mid w)P(w)$$

Similar to before, but there is a prior:  
incorporates prior knowledge

## Example: linear regression

Exercise 1 :)

## Aside: Bayesian inference

During training, calculate:

$$P(w \mid \mathcal{D}) = \frac{P(\mathcal{D} \mid w)P(w)}{P(\mathcal{D})} = \frac{P(\mathcal{D} \mid w)P(w)}{\int_{\mathbb{R}^d} P(\mathcal{D} \mid w')P(w')dw}$$

The evidence is usually hard to calculate



## Aside: Bayesian inference

During training, calculate:

$$P(w \mid \mathcal{D}) = \frac{P(\mathcal{D} \mid w)P(w)}{P(\mathcal{D})} = \frac{P(\mathcal{D} \mid w)P(w)}{\int_{\mathbb{R}^d} P(\mathcal{D} \mid w')P(w')dw}$$

The **evidence** is usually hard to calculate

Inference (prediction) for a new sample point  $x'$ :

$$P(y \mid x') = \int_{\mathbb{R}^d} P(y \mid x', w)P(w \mid \mathcal{D})dw$$

It is not a point estimate:

it gives a distribution over possible parameter/prediction values.