

1 **Using Electronic Health Record Audit Logs to Study Clinical Activity: A Systematic**
2 **Review of Aims, Measures, and Methods**

3
4 Adam Rule, PhD¹, Michael F. Chiang, MD^{1,2}, Michelle R. Hribar, PhD^{1,2}
5

6 ¹ Department of Medical Informatics and Clinical Epidemiology
7 Oregon Health & Science University
8 Portland, Oregon, USA
9

10 ² Department of Ophthalmology, Casey Eye Institute
11 Oregon Health & Science University
12 Portland, Oregon, USA
13

14 *Corresponding Author:*

15 Adam Rule, PhD
16 Oregon Health & Science University
17 Mail Code: BICC
18 3181 SW Sam Jackson Park Road
19 Portland, OR 97239
20 rulea@ohsu.edu
21 (206) 291-2533
22

23 Keywords: Electronic Health Records; Audit Logs
24

25 Word Count: 3,927 / 4,000 words
26

27 **Note: This is the Author's Original Version as it was submitted for peer review. The final**
28 **Accepted Manuscript and Version of Record will differ slightly. This article has been**
29 **accepted for publication in the Journal of the American Medical Informatics Association.**

ABSTRACT

Objective: To systematically review published literature and identify consistency and variation in the aims, measures, and methods of studies using electronic health record (EHR) audit logs to observe clinical activities.

Materials and Methods: In July 2019, we searched PubMed for articles using EHR audit logs to study clinical activities. We coded and clustered the aims, measures, and methods of each article into recurring categories. We likewise extracted and summarized the methods used to validate measures derived from audit logs and limitations discussed of using audit logs for research.

Results: Eighty-six articles met inclusion criteria. Study aims included examining EHR use, care team dynamics, and clinical workflows. Studies employed six key audit log measures: counts of actions captured by audit logs (e.g., problem list viewed), counts of higher-level activities imputed by researchers (e.g., chart review), activity durations, activity sequences, activity clusters, and EHR user networks. Methods used to preprocess audit logs varied, including how authors filtered extraneous actions, mapped actions to higher-level activities, and interpreted repeated actions or gaps in activity. Twenty studies validated results (23%), but only nine (10%) through direct observation, demonstrating varying levels of measure accuracy.

Discussion: While originally designed to aid access control, EHR audit logs have been used to observe diverse clinical activities. However, most studies lack sufficient discussion of measure definition, calculation, and validation to support replication, comparison, and cross-study synthesis.

Conclusion: EHR audit logs have potential to scale observational research but the complexity of audit log measures necessitates greater methodological transparency and validated standards.

BACKGROUND AND SIGNIFICANCE

Recently mandated logging of electronic health record (EHR) access in audit logs provides a promising resource for researchers to observe clinical activities at scale. Informaticians currently use diverse methods to study both clinical activities and use of health information technology (HIT) including surveys, interviews, and time-motion studies.[1–5] Time-motion studies in particular have seen wide adoption as they avoid many of the biases and inconsistencies of self-report through surveys and interviews. However, the most common form of time-motion study – continuous observation by an external observer – is time-consuming, expensive, and difficult to scale in terms of the diversity, duration, and detail of activity that can be recorded.[1–3] Researchers can scale certain aspects of observational studies with sensors such as Bluetooth beacons and video recorders, but this equipment can be difficult to set up and may provide, depending on the sensor, either a limited stream of data or multi-faceted recordings that require extensive ethnographic analysis.[6,7] Despite the many methods at their disposal, informaticians struggle to observe clinical activity and HIT use accurately, efficiently, and at scale.

Starting in 2005, the Security Rule of the Health Insurance Portability and Accountability Act (HIPAA) required all healthcare organizations to “*implement hardware, software, and/or procedural mechanisms that record and examine activity in information systems that contain or use electronic protected health information.*”[8] As part of the second stage of the Meaningful Use regulations in 2014,[9] the ONC further clarified that certified EHRs would need to maintain audit logs adhering to the ASTM E2147 standard for tracking HIT use.[10] Due to these regulations, virtually all EHRs in the United States now track at least four pieces of information about every episode of patient record access including *who* accessed *which* patient record at *what*

time and the *action* they performed in that record such as adding, deleting, or copying information (**Table 1**). Depending on the vendor, EHR audit logs may track additional information about the computer, user, or record involved in each action, track those actions at different levels of granularity, and give them different names.

Table 1: Example EHR audit log

TIME	USER	RECORD	ACTION	COMPUTER
05/12/2019 13:04:35	SMITHJANE	104738297	Edit Note Section	MED2938
05/12/2019 13:04:37	SMITHJANE	104738297	Pend Note	MED2938
05/12/2019 13:04:42	SMITHJANE	104738297	Sign Note	MED2938
05/12/2019 13:04:52	DOEJOHN	105837489	View Problem List	MED1238
05/12/2019 13:05:02	DOEJOHN	105837489	View Note	MED1238
05/12/2019 13:05:04	DOEJOHN	105837489	View Note	MED1238
05/12/2019 13:05:32	SMITHJANE	107483726	View Patient Summary	MED2938
05/12/2019 13:13:32	SMITHJANE	107483726	View Patient Summary	MED2938
...

While originally designed to monitor record access, due to widespread use of EHRs in healthcare EHR audit logs present a unique opportunity to study clinical activities at a scale unachievable with direct observation and with less setup than external sensors. However, like other forms of time-motion study, audit log research is subject to challenges and limitations. Since audit logs are not purpose-built to track workflows they may lack vital contextual information and logged actions may be difficult to map to clinical activities such as chart review or patient exams. Nor do all clinical activities involve EHR use. While EHR audit logs have been used to study diverse clinical activities, there has been little synthesis of the aims of this research, or examination of the variation and validity of measures and methods employed. This lack of synthesis hampers efforts to replicate, generalize, and compare research in areas that may benefit from audit log analysis such as EHR usability and provider burnout.[11–17]

Objective

With this systematic review we identify consistency and variation in the aims, measures, and methods of audit log research. Moreover, we consolidate evidence for the validity of

measures derived from audit logs and limitations of using audit logs to observe clinical activities. With this review we aim to improve the quality and generalizability of audit log research and provide literature-driven recommendations for the design of future studies to ultimately foster knowledge discovery in areas of critical informatics research.

MATERIALS AND METHODS

We identified articles for review by searching PubMed. Since the terms used to describe audit logs vary, we first hand-selected twenty-one audit log articles familiar to us and identified the terms each used to describe audit logs (e.g., access log, usage log, EHR timestamps). Using these synonyms for “audit log” and descriptors of EHRs used in prior systematic reviews,[18,19] we searched PubMed in July 2019 for all literature referencing EHR audit logs (see Appendix for full query). The PubMed query and hand-selection together returned 1775 unique articles, with only one of the hand-selected articles not included in the PubMed results. Through manual title, abstract, and text review, one author (AR) identified 74 of these articles which met inclusion criteria summarized in **Figure 1**. Scanning the references of included articles, we identified 12 additional articles which met inclusion criteria, yielding a total of 86 articles for review (**Figure 2**).

One author (AR) iteratively reviewed and extracted features of each article. These included *study features*: the terms used to describe audit logs, EHR vendor, users studied (e.g., physicians, nurses), duration of study, and reported sample sizes (e.g., number of users, patient records, or encounters studied). This author also extracted each article’s research questions, measures, and data preprocessing methods and together with a second author (MRH) iteratively coded these into a concise set of *aims*, *measures*, and preprocessing *methods* used in audit log research. Lastly, one author (AR) extracted and summarized the methods and results of

validation studies and sensitivity analyses reported in reviewed articles as well as *limitations* discussed of using audit logs for research.

RESULTS

Features of Audit Log Research

The 86 articles included in this review used a variety of terms to describe audit logs in their titles and abstracts (**Table 2**). Only 31 used terms including the words “audit” or “access” while the remainder referenced more ambiguous EHR data, metadata, timestamps, and logs. Articles also varied in the EHRs, features, and users studied (**Table 2**). Just over half analyzed audit-logs from commercial EHRs (28 from one vendor, Epic (Verona, WI)). Most articles (66) examined all EHR activity while a minority (20) measured interactions with specific features or data-types such as info buttons, handoff reports, or CT scans. Just over half (47 articles) examined EHR activity in individual departments such as internal medicine, outpatient primary-care, and ophthalmology, while the remainder spanned departments. Only six articles examined EHR use across multiple institutions: four of which were conducted outside the United States and two of which examined interactions with a web-based EHR. Most articles (52) studied all EHR users while the remainder largely studied physician and resident use (31) with only a small number focused on nurses and medical students (3). Most articles (74) reported the length of time studied with the median duration being one year. Articles were less consistent in reporting the number of users, actions, patient records, and encounters studied (**Table 3**). Just over half of articles were published in 2016 or later (44 articles). See **Table 4** for features by article.

Table 2: Features of studies using EHR audit logs to study clinical activity

<i>Study Attribute</i>	#	%
Audit Log Term		
Audit (e.g., audit log, access log)	31	36
Generic (e.g., log file, EHR log)	18	21

	Usage (e.g., usage log, usage patterns)	10	12
	Time (e.g., timestamp, time data)	7	8
	Data (e.g., EHR data, EHR metadata)	8	9
	Event (e.g., event file, event sequence)	6	7
	Other (e.g., system log, user log)	6	7
EHR Type	Vendor	45	52
	Homegrown	25	29
	Unstated	16	19
Scope	Whole EHR	66	77
	Specific Feature	20	23
Department	Multiple	39	45
	Ophthalmology	10	12
	Primary Care	9	10
	General Internal Medicine	7	8
	Emergency	6	7
	Other	15	17
Institution	Single	80	93
	Multi	6	7
Users	All	52	60
	Physicians	20	23
	Residents/Fellows	11	13
	Nurses	2	2
	Medical Students	1	1

139

140 Table 3: Amount of time, actions, users, patients, and encounters studied varied across articles

	<i>Time</i> <i>(Months)</i>	<i>Users</i>	<i>Actions</i>	<i>Encounters</i>	<i>Patients</i>
<i>Studies Reporting</i>	74	50	24	18	19
<i>Minimum</i>	0.25	15	20,249	249	100
<i>Median</i>	12	155	1,930,620	38,628	3,450
<i>Maximum</i>	120	10,659	118,000,000	3,219,910	815,114

141

142 Table 4: Select features of the 86 articles included in this systematic review

Article				Terms	Who			What			Aims					Measures						Methods			Validation		
Ref.	PMID	First Author	Year	Audit Log Term	Users Studied	Department	Multi-Institute	EHR Type	Scope	Months Logged	EHR Use	Workflow	Care Team	Model EHR Use	Model Outcome	Action	Activity	Duration	Sequence	Cluster	Network	Filter Activity	Map Activity	Gaps / Repeat	Validate Mapping Method	Validate Time Method	Sensitivity
20	8130443	Michael PA	1993	audit trail	All	All		Homegrown	Feature	9	X					X											
21	14728157	Cimino JJ	2003	log file	All	All		Homegrown	Feature	6	X					X										Survey	
22	17102263	Chen ES	2006	log file	All	All		Homegrown	Feature	18	X					X											
23	17238321	Cimino JJ	2006	log file	All	All		Homegrown	Feature	24	X					X											
24	18693813	Cimino JJ	2007	log file	All	All		Homegrown	Feature	14	X					X		X									
25	18308208	McLean TR	2008	metadata	Trainee	Surgery		Homegrown	Feature	5	X					X											
26	20180439	Bernstein JA	2010	log data	All	All		Vendor	Feature	40	X					X											
27	22874273	Ries M	2012	system log	All	All		Vendor	Feature	8	X					X											
28	25024755	Hum RS	2014	user log	All	NICU		Vendor	Feature	22	X					X											
29	25954381	Jiang SY	2014	usage log	All	All		Vendor	Feature	1	X					X				X							
30	26958202	Jiang SY	2015	audit log	All	All		Vendor	Feature	1	X					X				X		X					
31	28808942	Cutrona SL	2017	access/audit log	MD	Primary		Vendor	Feature	12	X		X			X		X									
32	29696473	Mongan J	2018	audit log	MD	Rad		Vendor	Feature	14	X					X					X						
33	30879188	Epstein RH	2019	access log	MD	Anes		Vendor	Feature	19	X					X						X					
34	-	Asaro P	2001	access log	All	All		Homegrown	EHR	12	X					X			X								
35	14728151	Chen ES	2003	log file	All	All		Homegrown	EHR	12	X					X		X	X		X	X					
36	15360766	Chen ES	2004	log file	All	All		Homegrown	EHR	12	X					X		X	X			X					
37	16779018	Clayton PD	2005	audit trail	All	All		Homegrown	EHR	60	X					X											
38	17213496	Hripcsak G	2007	audit log	All	ED		Homegrown	EHR	7	X					X											
39	18999307	Wilcox A	2008	usage statistics	MD	All		Homegrown	EHR	1	X					X											
40	20442152	Zheng K	2010	audit log	MD	Primary		Homegrown	EHR	14	X		X			X					X						
41	20841655	Bowes WA III	2010	audit log	All	All		Homegrown	EHR		X					X										Interview	
42	21292704	Sykes TA	2011	system log	All	All		EHR			X		X	X		X											
43	23909863	Park JY	2014	log data	All	All		Homegrown	EHR	24	X					X											
44	24914013	Ancker JS	2014	EHR data	All	Primary		Vendor	EHR	42	X					X											
45	26618036	Choi W	2015	log file	All	All		Homegrown	EHR	12	X					X											
46	26831123	Kim S	2016	log file	All	All		Homegrown	EHR	24	X		X			X											
47	27332378	Kajimura A	2016	access log	Nurses	IM		EHR	<1	X	X					X											
48	29046269	Kim J	2017	usage log	MD	All		Homegrown	EHR	10	X		X			X			X		X						
49	29237579	Lee Y	2017	usage log	All	All		Homegrown	EHR	54	X					X											
50	29295318	Kim J	2017	usage patterns	MD	All		EHR	10	X	X					X											
51	30240357	Shenvi EC	2018	access log	Trainee	IM		Vendor	EHR	6	X		X			X											
52	30274967	Graham TA	2018	audit log	MD	ED	X	Homegrown	EHR	18	X					X						X					
53	31183688	Cohen GR	2019	log	All	Primary	X	EHR	1	X	X		X			X						X			Vendor		
54	24907594	Chi J	2014	audit data	MD Stu	All		Vendor	EHR	7	X			X		X	X				X				Survey		
55	26642261	Ouyang D	2016	electronic audit	Trainee	IM		Vendor	EHR	12	X					X	X	X					X				
56	26913101	Chen L	2016	audit log	Trainee	IM		Vendor	EHR	4	X					X	X				X	X		Vendor			
57	30522828	Cox ML	2018	time data	Trainee	Surgery		Vendor	EHR	11	X		X			X	X	X				X					X
58	30815089	Goldstein IH	2018	audit log	MD	Ophth		Vendor	EHR	12	X					X	X					X			Survey		
59	30726208	Wang JK	2019	event log	Trainee	IM		Vendor	EHR	41	X					X	X				X	X					X

Ref = Reference; PMID = PubMed ID; MD = Physicians; Med Stu = Medical Students
Anes = Anesthesiology; Card = Cardiology; ED = Emergency Department; Heme = Hematology; IM = Internal Medicine; NICU = Neonatal Intensive Care Unit; Ophth = Ophthalmology;
Peds = Pediatrics; Rad = Radiology

Aims of Audit Log Research

Most articles used audit logs to study *EHR use* directly (63 articles, see **Table 4** for details by article).[20–82] This included how often providers accessed individual pieces of information,[20–33] patterns of EHR use across features,[34–53,67,74–78], and the total time providers spent using EHRs.[54–60,68–73] More recently, studies began to use audit logs to examine *clinical workflows* that extend beyond the EHR, using audit log timestamps to mark the boundaries of clinical events (30 articles).[69–98] For example, a few articles calculated resident duty hours using EHR login and logout timestamps, assuming residents log into and out of the EHR near shift boundaries.[79,85–87] Other studies used timestamps to identify the start and end of clinical exams and calculate exam length or patient wait time.[91–98]. Still other workflow studies focused instead on sequences of actions providers took after specific events occurred (e.g. receiving an alert) or their typical workflow when caring for certain patient groups, such as those with complex cardiac conditions.[80–84] A third common use of audit logs was to study *care team structure and dynamics* (17 articles).[63–69,77,97–105] While a few studies used EHR access to identify care teams for individual patients,[66,99] more used co-access of the same records to identify which providers or departments consistently worked together across patients.[65,100–105]

In addition to these three core aims, many studies collected additional demographic, contextual, or outcome data to model the effect of EHR use on clinical outcomes (12 articles),[42,54,63,64,67,73–76,79,86,104] or the effect of patient, provider, and context on EHR use (15 articles).[31,40,42,46,48,53,57,62,69–71,81,82,96,105] For example, two studies modeled EHR adoption as a function of providers’ demographics and professional

networks.[40,46] Several other studies assessed whether accessing a patient’s historical record decreased their length of stay or chances of being admitted to the hospital.[64,73–76]

Measures of Audit Log Research

Reviewed articles derived a variety of measures derived from audit logs to study these topics including 1) counts of actions captured by audit logs, 2) counts of higher-level activities imputed by researchers, 3) activity durations, 4) activity sequences, 5) activity clusters, and 6) networks of EHR users (summarized in **Figure 3**, see **Table 4** for details by article).

Counts of actions captured directly by audit logs (63 articles) [20–53,55,61–67,74–78,80,82–84,86,88,89,97–105] such as “problem list viewed” were often used to quantify use of specific features such as info buttons and radiology reports. Alternatively, these actions were sometimes aggregated to identify peak periods of EHR use throughout the day or week. *Counts of higher-level activities* (27 articles) [54–60,68–73,79–81,85–88,90–96] typically involved first mapping low-level actions to higher-level activities such as chart review and documentation. Alternatively, it might involve looking for significant gaps between actions to identify entire sessions of EHR use or work shifts. These activity boundaries could then be used to compute counts or rates, such as the number of unique EHR sessions across all users in the past month or the percent of encounters where providers reviewed the patient’s historical record. Other studies grouped actions into higher-level activities not to report counts but to compute the *duration of those activities* including total time devoted to EHR use (30 articles). [24,31,54–60,68–73,77,79,81,83,85–87,90–96,105]

These first three measures were used to create more complex measures and models of activity, three of which were employed in multiple studies. Eight studies constructed *event sequences* to identify routine patterns of care and deviations from

them.[35,36,78,83,84,89,96,104] Twelve studies clustered *patterns of activity* to identify recurring patterns of EHR use, such as which sections of the record providers routinely accessed.[29,30,34–36,48,64,78,83,84,89,104] Finally, eleven articles studying care teams used co-access of a patient’s record to develop *networks of users or departments* that typically work together.[64–66,68,69,100–105] Across all six measures, there was one significant change in use over time: 47% of articles published since 2016 reported a time duration, whereas only 21% of the articles published before 2016 did so ($\chi^2 = 6.54$, $p = 0.01$) (**Figure 4**).

Preprocessing Methods of Audit Log Research

Computing even seemingly simple measures from audit logs such as duration of EHR use is not necessarily straightforward. Yet, less than half of articles (32) discussed how raw audit logs were preprocessed before analysis (see **Table 4** for details by article). Fewer still discussed this data wrangling in enough detail to support replication. When reported, common practices included 1) filtering actions, 2) mapping actions to higher-level clinical activities, and 3) selecting criteria to define time-periods. *Filtering actions* included removing actions that were considered incidental or irrelevant to the study.[30,32,35,40,48,54,56,57,59,75,81,86] For example, one study of medical student EHR use removed short bursts of activity on off-service days, labeling this incidental use.[54] Other studies considered all activity within 24 hours of a patient’s visit as relevant,[40] or only activity in periods with “more than 3 mouse clicks (or 15 keystrokes) or 1700 mouse miles (pixels) per minute”.[56] Another common preprocessing practice was *mapping individual actions to higher-level activities* such as chart review or documentation.[33,35,36,52,53,64,72,73,82,84,88] While no study reported the actual action-activity mappings, some reported the process used to develop these mappings, which varied as discussed in the next section. A final recurring practice was *selecting actions and criteria to*

define time-periods. [55,56,58–61,70,71,81,85,94,96,104] This involved defining which actions constituted the start and end of clinical events (such as the first non-login action) and how gaps in activity would be handled. Depending on the research question, meaningful gaps ranged from 5 minutes, which could indicate the user was no longer actively using the EHR, [59] to six hours, which could indicate the end of a shift. [85] Another study identified shifts using a three-step process of 1) identifying distinct shifts based on 4-hour gaps, 2) merging shifts that were less than 7 hours apart which would result in a combined shift length of less than 30 hours, and 3) merging shifts that were less than 2 hours long and would result in a combined shift of less than 20 hours. [87]

Validating Audit Log Measures

Using EHR audit logs to study clinical activity assumes audit logs consistently and accurately track clinical activities and that the methods used to process them into more complex measures are sound. However, a minority of studies reported checking these assumptions through validation or sensitivity analyses. Validation studies, which compared measures derived from audit logs with those obtained through other methods, checked both the mapping of audit log actions to higher-level activities as well as the accuracy of activity patterns or durations derived from audit logs. Of the twenty studies that reported validation analyses, six validated activity mappings and sixteen validated patterns or durations (see **Table 4** for details by article).

The six studies that reported validating action-activity mappings used a variety of methods including consensus among two or more researchers, [88] consulting the EHR vendor, [53,56] and direct observation of clinical activities. [72,73,82] Only one of these studies reported the accuracy of mappings, noting that 6.9% of the audit log actions were originally misclassified as representing the wrong activity when compared to direct observation. [72] Of the

sixteen studies that reported validating activity patterns or durations, nine compared them to data self-reported by EHR users, administrators, or the authors themselves.[21,33,41,54,69,85,87,88,102] Only seven compared timing data to values obtained through direct observation.[70–72,90,91,94,95] Of these, only five reported measure accuracy. Accuracy for EHR time per encounter ranged from overestimating by 43% (4.3 vs 3.0 minutes)[70] to underestimating by 33% (2.4 ± 1.7 vs. 1.6 ± 1.2 min).[91] Measures of appointment lengths which tend to be longer were more accurate, overestimated by just 4% in one study (13.8 ± 8.2 vs. 13.3 ± 7.3 min),[95] underestimated by 14% in another (19.4 vs 22.5 min),[71] and overestimated by 29% in a third (24.4 ± 13.0 vs. 18.9 ± 11.0 min).[91]

Computing duration data in particular requires making a number of assumptions about what constitutes the start and end of certain activities and how to handle gaps in activity. Four studies reported sensitivity analyses in this vein,[57,59,62,86] such as varying the gap in actions considered idle activity from 5-10 minutes[59] or seeing what impact discarding the first and last 5% of actions in a given work shift had on calculated shift length.[86] None reported a significant change in results due to changing these parameters.

Challenges and Limitations of Audit Log Research

Finally, reviewed articles mentioned a few limitations of using audit logs to study clinical activity. First, 20 articles mentioned that audit logs do not provide a full picture of clinical activity as they only capture interactions with the EHR.[20,30,38,51,52,57,64,68,69,72,77,78,85,87,91,92,96,100,104,105] Audit logs do not track phone, pager, or face-to-face interactions nor do they track interaction with paper records and printouts. This may lead to systematic underestimation of interaction or workload. Second, 15 articles noted that gaps between timestamps and multiple concurrent timestamps can be difficult

to interpret.[41,54,58,59,68,70,72,81,82,86,88,90,93–95] For example, does a long gap mean the provider was meaningfully engaged with the EHR that entire time, or had they turned their attention away? Do sequential identical audit log entries (except for the timestamp) represent a repeated or continued activity? Third, seven articles mentioned audit log data were often either too coarse or too detailed for clear interpretation.[20,49,54,69,70,100,103] Logs might capture who accessed a patient record, but not which exact note or result they were viewing. Alternatively, more detailed logs might use different names to track accessing the same piece of information through different screens. It can take researchers substantial time to map these isometric actions to higher-level activities. Lastly, 6 articles noted that audit logs may capture what a user did in the EHR, but data from more qualitative methods such as interviews or observations are needed to understand why.[21,32,38,61,68,81]

DISCUSSION

With this systematic review, we surveyed articles using EHR audit logs to study clinical activities. We found a diverse body of literature employing a range of measures to study EHR use directly, clinical workflows extending beyond the EHR, and care team dynamics. This body of research is growing with over half of reviewed articles published in the last three-and-a-half years. Moreover the increased measurement of duration of EHR use may reflect growing concern over the association between EHR use and provider burnout.[14–17]

Whereas some measures employed in this literature were relatively simple counts of actions tracked explicitly by audit logs, others required researchers to manipulate audit logs in sophisticated ways, generating durations, sequences, clusters, and networks. Many studies glossed over the details of how raw audit logs were preprocessed to compute these measures and even when methods were reported there was significant variation. This variation reflects the

difficulty of interpreting audit logs which requires professional judgement and domain knowledge, such as understanding resident duty hour restrictions.

Recommendations

The variability of measures and methods in reviewed articles echoes the variability observed in prior systematic reviews of the time-motion studies in healthcare.[2] It also highlights areas where research using EHR audit logs might improve. We focus our recommendations on four areas: sample size reporting, reporting of methods used to pre-process audit logs, validation and sensitivity analyses, and methodological transparency leading to validated standards.

First, we recommend standard reporting of the time period, number of users, and patient records studied. While most studies report the duration of time studied, not all did. Just over half reported the number of users studied, and far fewer reported the number of patients or encounters analyzed. This use of time to report sample sizes likely reflects the fact that audit log data are routinely queried by time period rather than number of patient records or users desired for analysis. We suggest other reported sample size measures be clinically relevant, such as the number of patient encounters, rather than dataset measures such as number audit log rows, as these are harder to compare across vendors and institutions with different logging practices.

Second, we recommend detailed reporting of steps used to wrangle raw audit log data into measures. Given the variable accuracy of time durations reported in studies that validated them (e.g., from 33% underestimation to 43% overestimation of provider EHR time per appointment), there is still a need to develop more accurate and consistent methods of tracking activities with audit logs. Methods reporting should include any criteria used to filter logs and at least the *process* used to map granular actions into higher-level activities such as documentation

or chart review. Ideally researchers would also report the exact mapping of actions to activities; however, this may not be feasible given the large number of actions that may map to a single activity or the potential for EHR vendors to consider audit log action names proprietary information. For time durations, we recommend authors report how they handle repeated actions and gaps in activity, as well as how they identify the boundaries of activities, especially if data are missing (e.g., “if a log-out action was missing, we considered the last action before a gap of 2 or more hours the end of the provider’s shift”). We recommend the audit log research community develop standards for reporting more complex measures such as activity sequences, activity clusters, and user networks.

Third, we recommend researchers take more steps to validate and test the sensitivity of their results. Ultimately, the validity of audit log research rests on assumptions that audit logs consistently and accurately track EHR use and clinical activities more broadly. While some methods seem to be approaching parity with direct observation for measuring the duration of longer activities such as patient exams, measures of shorter events such as EHR time per encounter are more varied. Validation may occur in a number of ways including surveys and member-checks, but the gold-standard should remain comparing measures derived from audit logs with those obtained through direct observation. More sensitivity analyses are also warranted as the parameters of methods used to preprocess audit logs may significantly affect results.

Finally, there is a need for greater methodological transparency and validated standards to support replication and synthesis. This includes clear documentation and sharing of data schemas, action-activity mappings, and preprocessing scripts between institutions. We recommend that vendors, institutions, and the audit-log research community work together to share methods and develop validated standards for tracking, querying, and analyzing audit logs

to compute the diverse measures of clinical activity uncovered in this review. These standards could in turn support replication and comparison across departments and institutions to identify consistency and variation in EHR use and clinical workflows between them.

Limitations

This review has a few limitations. First, it does not survey use of all health-information technology logs, nor even all uses of EHR audit logs. EHR related technologies such as Personal Health Records, Health Information Exchanges, and mobile health apps often track user activity with logs similar to EHR audit logs,[106–109] and workflow researchers may use timing data directly from patient records in their studies (such as admit time or time of placing an order). EHR audit logs are also routinely used for their primary purpose of access control and several publications have explored how to use them more effectively for that purpose.[110–114] While the measures and methods used in these related domains may be similar to those reported in this review, we scoped our analysis to use of EHR audit logs to study clinical activity to provide targeting insights for this growing research community. Second, we limited our search to articles on PubMed which may exclude articles published in computer science or engineering venues not routinely indexed there. We mitigated this risk by searching the citations of included articles for relevant references, regardless of venue. Third, our coding process was largely subjective and performed by a single author. While the authors of each article may not agree with our classification, we aimed to develop a consistent coding scheme that captured the breadth of the literature by iteratively defining and applying each category label. Finally, this review likely reflects a publication bias in which some types of audit log research are more readily published than others (e.g. workflow studies vs. studies of IT infrastructure needs)

CONCLUSION

EHR audit logs have been used to study a wide range of clinical activities, extending beyond their original purpose of monitoring patient record access. The 86 articles included in this review demonstrate a diverse and growing literature, reflecting researchers' desire to gather precise data on HIT use and clinical activities at scale. However, the process of turning raw audit logs into insights is complex, requires professional judgement, and varies from study to study, when it is even reported. Moreover, there are relatively few articles in the literature that report testing the validity and sensitivity of audit log measures. This lack of rigor and reporting prevents synthesis and comparison across studies, as well as efforts to improve the accuracy of using audit logs for clinical event measurement. EHR audit logs have untapped potential to support quality improvement and research, but the continued growth of the field will require greater methodological transparency and validated standards to support replication and cross-study knowledge discovery.

COMPETING INTERESTS

The authors have no commercial, proprietary, or financial interest in any of the products or companies described in this article. MFC is an unpaid member of the Scientific Advisory Board for Clarity Medical Systems (Pleasanton, CA), a Consultant for Novartis (Basel, Switzerland), and an initial member of Intelereitina, LLC (Honolulu, HI).

FUNDING

Supported by grants R00LM12238, P30EY10572, and T15LM007088 from the National Institutes of Health (Bethesda, MD), and by unrestricted departmental funding from Research to Prevent Blindness (New York, NY). The funding organizations had no role in the design or conduct of this research.

ACKNOWLEDGEMENTS

Thank you to Julia Adler-Milstein and members of the National Research Network for EHR Audit-Logs and Metadata for help hand-selecting articles to seed this systematic review. Thank you to Julia Adler-Milstein, Genna Cohen, and Nicole Weiskopf for feedback on early versions of this article.

APPENDIX

PubMed Search Query

(
("audit"[All Fields] AND (log[All Fields] OR file[All Fields] OR data[All Fields]))
OR (log[All Fields] AND (file[All Fields] OR "event"[All Fields] OR "access"[All Fields] OR
"system"[All Fields]

384 OR "usage"[All Fields] OR "activity"[All Fields]))
 385 OR "timestamp"[All Fields]
 386 OR "interaction patterns"[All Fields]
 387 OR "utilization patterns"[All Fields]
 388)
 389 AND
 390 (
 391 ("electronic"[All Fields] AND "health"[All Fields] AND record[All Fields])
 392 OR ("electronic"[All Fields] AND "medical"[All Fields] AND record[All Fields])
 393 OR (("computerised"[All Fields] OR "computerized"[All Fields]) AND "medical"[All Fields]
 394 AND record[All Fields])
 395 OR ("electronic health records"[MeSH Terms])
 396 OR ("medical records systems, computerized"[MeSH Terms])
 397)
 398 AND
 399 English[lang]

REFERENCES

- 1 Unertl KM, Novak LL, Johnson KB, *et al.* Traversing the many paths of workflow research: developing a conceptual framework of workflow terminology through a systematic literature review. *J Am Med Inform Assoc* 2010;17:265–73.
- 2 Zheng K, Guo MH, Hanauer DA. Using the time and motion method to study clinical work processes and workflow: methodological inconsistencies and a call for standardized research. *J Am Med Inform Assoc* 2011;18:704–10.
- 3 Lopetegui M, Yen P-Y, Lai A, *et al.* Time motion studies in healthcare: what are we talking about? *J Biomed Inform* 2014;49:292–9.
- 4 Friedman CP, Wyatt J. *Evaluation methods in biomedical informatics*. New York; London: : Springer 2011.
- 5 Kannampallil TG, Abraham J. Evaluation of health information technology: methods, frameworks and challenges. In: *Cognitive Informatics for Biomedicine*. Springer 2015. 81–109.
- 6 Zheng K, Hanauer DA, Weibel N, *et al.* Computational Ethnography: Automated and Unobtrusive Means for Collecting Data In Situ for Human–Computer Interaction Evaluation Studies. In: Patel VL, Kannampallil TG, Kaufman DR, eds. *Cognitive Informatics for Biomedicine*. Cham: : Springer International Publishing 2015. 111–40.
- 7 Weibel N, Rick S, Emmenegger C, *et al.* LAB-IN-A-BOX: semi-automatic tracking of activity in the medical office. *Pers Ubiquit Comput* 2015;19:317–34.
- 8 Health Insurance Portability and Accountability Act, Technical Safeguards, 45 C.F.R § 164.312. 2003.
- 9 Standards for health information technology to protect electronic health information created, maintained, and exchanged, 45 C.F.R. § 170.210. 2015.
- 10 ASTM E2147 - Standard Specification for Audit and Disclosure Logs for Use in Health Information Systems. <https://www.astm.org/Standards/E2147.htm>
- 11 Middleton B, Bloomrosen M, Dente MA, *et al.* Enhancing patient safety and quality of care by improving the usability of electronic health record systems: recommendations from AMIA. *J Am Med Inform Assoc* 2013;20:e2-8.
- 12 Zhang J, Walji MF. TURF: toward a unified framework of EHR usability. *J Biomed Inform* 2011;44:1056–67.
- 13 Ratwani RM, Fairbanks RJ, Hettinger AZ, *et al.* Electronic health record usability: analysis of the user-centered design processes of eleven electronic health record vendors. *J Am Med Inform Assoc* 2015;22:1179–82.

- 434 14 Bodenheimer T, Sinsky C. From triple to quadruple aim: care of the patient requires care of
435 the provider. *Ann Fam Med* 2014;12:573–6.
- 436 15 Shanafelt TD, Hasan O, Dyrbye LN, *et al.* Changes in Burnout and Satisfaction With Work-
437 Life Balance in Physicians and the General US Working Population Between 2011 and 2014.
438 *Mayo Clin Proc* 2015;90:1600–13.
- 439 16 Shanafelt TD, Dyrbye LN, Sinsky C, *et al.* Relationship Between Clerical Burden and
440 Characteristics of the Electronic Environment With Physician Burnout and Professional
441 Satisfaction. *Mayo Clin Proc* 2016;91:836–48.
- 442 17 Gardner RL, Cooper E, Haskell J, *et al.* Physician stress and burnout: the impact of health
443 information technology. *J Am Med Inform Assoc* 2019;26:106–14.
- 444 18 Hogan WR, Wagner MM. Accuracy of data in computer-based patient records. *J Am Med*
445 *Inform Assoc* 1997;4:342–55.
- 446 19 Weiskopf NG, Weng C. Methods and dimensions of electronic health record data quality
447 assessment: enabling reuse for clinical research. *J Am Med Inform Assoc* 2013;20:144–51.
- 448 20 Michael PA. Physician-directed software design: the role of utilization statistics and user
449 input in enhancing HELP results review capabilities. *Proc Annu Symp Comput Appl Med*
450 *Care* 1993;107–11.
- 451 21 Cimino JJ, Li J, Graham M, *et al.* Use of online resources while using a clinical information
452 system. *AMIA Annu Symp Proc* 2003;175–9.
- 453 22 Chen ES, Bakken S, Currie LM, *et al.* An automated approach to studying health resource
454 and infobutton use. *Stud Health Technol Inform* 2006;122:273–8.
- 455 23 Cimino JJ. Use, usability, usefulness, and impact of an infobutton manager. *AMIA Annu*
456 *Symp Proc* 2006;151–5.
- 457 24 Cimino JJ, Friedmann BE, Jackson KM, *et al.* Redesign of the Columbia University
458 Infobutton Manager. *AMIA Annu Symp Proc* 2007;135–9.
- 459 25 McLean TR, Burton L, Haller CC, *et al.* Electronic medical record metadata: uses and
460 liability. *J Am Coll Surg* 2008;206:405–11.
- 461 26 Bernstein JA, Imler DL, Sharek P, *et al.* Improved physician work flow after integrating
462 sign-out notes into the electronic medical record. *Jt Comm J Qual Patient Saf* 2010;36:72–8.
- 463 27 Ries M, Golcher H, Prokosch H-U, *et al.* An EMR based cancer diary - utilisation and initial
464 usability evaluation of a new cancer data visualization tool. *Stud Health Technol Inform*
465 2012;180:656–60.

- 466 28 Hum RS, Cato K, Sheehan B, *et al.* Developing clinical decision support within a
467 commercial electronic health record system to improve antimicrobial prescribing in the
468 neonatal ICU. *Appl Clin Inform* 2014;5:368–87.
- 469 29 Jiang SY, Murphy A, Vawdrey D, *et al.* Characterization of a handoff documentation tool
470 through usage log data. *AMIA Annu Symp Proc* 2014;749–56.
- 471 30 Jiang SY, Hum RS, Vawdrey D, *et al.* In Search of Social Translucence: An Audit Log
472 Analysis of Handoff Documentation Views and Updates. *AMIA Annu Symp Proc* 2015;669–
473 76.
- 474 31 Cutrona SL, Fouayzi H, Burns L, *et al.* Primary Care Providers’ Opening of Time-Sensitive
475 Alerts Sent to Commercial Electronic Health Record InBaskets. *J Gen Intern Med*
476 2017;32:1210–9.
- 477 32 Mongan J, Avrin D. Impact of PACS-EMR Integration on Radiologist Usage of the EMR. *J*
478 *Digit Imaging* 2018;31:611–4.
- 479 33 Epstein RH, Dexter F, Schwenk ES. Provider Access to Legacy Electronic Anesthesia
480 Records Following Implementation of an Electronic Health Record System. *J Med Syst*
481 2019;43:105.
- 482 34 Asaro PV, Ries JE. Data mining in medical record access logs. In: *AMIA Annu Symp Proc.*
483 American Medical Informatics Association 2001;855.
- 484 35 Chen ES, Cimino JJ. Automated discovery of patient-specific clinician information needs
485 using clinical information system log files. *AMIA Annu Symp Proc* 2003;145–9.
- 486 36 Chen ES, Cimino JJ. Patterns of usage for a Web-based clinical information system. *Stud*
487 *Health Technol Inform* 2004;107:18–22.
- 488 37 Clayton PD, Naus SP, Bowes WA, *et al.* Physician use of electronic medical records: issues
489 and successes with direct data entry and physician productivity. *AMIA Annu Symp Proc*
490 2005;141–5.
- 491 38 Hripcsak G, Sengupta S, Wilcox A, *et al.* Emergency department access to a longitudinal
492 medical record. *J Am Med Inform Assoc* 2007;14:235–8.
- 493 39 Wilcox A, Bowes WA, Thornton SN, *et al.* Physician use of outpatient electronic health
494 records to improve care. *AMIA Annu Symp Proc* 2008;809–13.
- 495 40 Zheng K, Padman R, Krackhardt D, *et al.* Social networks and physician adoption of
496 electronic health records: insights from an empirical study. *J Am Med Inform Assoc*
497 2010;17:328–36.
- 498 41 Bowes WA. Measuring use of electronic health record functionality using system audit
499 information. *Stud Health Technol Inform* 2010;160:86–90.

500 42 Sykes TA, Venkatesh V, Rai A. Explaining physicians' use of EMR systems and
501 performance in the shakedown phase. *J Am Med Inform Assoc* 2011;18:125–30.

502 43 Park J-Y, Lee G, Shin S-Y, *et al.* Lessons learned from the development of health
503 applications in a tertiary hospital. *Telemed J E Health* 2014;20:215–22.

504 44 Ancker JS, Kern LM, Edwards A, *et al.* How is the electronic health record being used? Use
505 of EHR data to assess physician-level variability in technology use. *J Am Med Inform Assoc*
506 2014;21:1001–8.

507 45 Choi W, Park M, Hong E, *et al.* Early Experiences with Mobile Electronic Health Records
508 Application in a Tertiary Hospital in Korea. *Healthc Inform Res* 2015;21:292–8.

509 46 Kim S, Lee K-H, Hwang H, *et al.* Analysis of the factors influencing healthcare
510 professionals' adoption of mobile electronic medical record (EMR) using the unified theory
511 of acceptance and use of technology (UTAUT) in a tertiary hospital. *BMC Med Inform Decis*
512 *Mak* 2016;16:12.

513 47 Kajimura A, Takemura T, Hikita T, *et al.* Nurses' Actual Usage of EMRs: An Access Log-
514 Based Analysis. *Stud Health Technol Inform* 2016;225:858–9.

515 48 Kim J, Lee Y, Lim S, *et al.* What Clinical Information Is Valuable to Doctors Using Mobile
516 Electronic Medical Records and When? *J Med Internet Res* 2017;19:e340.

517 49 Lee Y, Park YR, Kim J, *et al.* Usage Pattern Differences and Similarities of Mobile
518 Electronic Medical Records Among Health Care Providers. *JMIR Mhealth Uhealth*
519 2017;5:e178.

520 50 Kim J, Lee Y, Lim S, *et al.* How Are Doctors Using Mobile Electronic Medical Records? An
521 In-Depth Analysis of the Usage Pattern. *Stud Health Technol Inform* 2017;245:1231.

522 51 Shenvi EC, Feupe SF, Yang H, *et al.* "Closing the loop": a mixed-methods study about
523 resident learning from outcome feedback after patient handoffs. *Diagnosis (Berl)*
524 2018;5:235–42.

525 52 Graham TA, Ballermann M, Lang E, *et al.* Emergency Physician Use of the Alberta Netcare
526 Portal, a Province-Wide Interoperable Electronic Health Record: Multi-Method
527 Observational Study. *JMIR Med Inform* 2018;6:e10184.

528 53 Cohen GR, Friedman CP, Ryan AM, *et al.* Variation in Physicians' Electronic Health Record
529 Documentation and Potential Patient Harm from That Variation. *J Gen Intern Med* Published
530 Online First: 10 June 2019.

531 54 Chi J, Kugler J, Chu IM, *et al.* Medical students and the electronic health record: "an epic
532 use of time." *Am J Med* 2014;127:891–5.

533 55 Ouyang D, Chen JH, Hom J, *et al.* Internal Medicine Resident Computer Usage: An
534 Electronic Audit of an Inpatient Service. *JAMA Intern Med* 2016;176:252–4.

535 56 Chen L, Guo U, Illipparambil LC, *et al.* Racing Against the Clock: Internal Medicine
536 Residents' Time Spent On Electronic Health Records. *J Grad Med Educ* 2016;8:39–44.

537 57 Cox ML, Farjat AE, Risoli TJ, *et al.* Documenting or Operating: Where Is Time Spent in
538 General Surgery Residency? *J Surg Educ* 2018;75:e97–106.

539 58 Goldstein IH, Hribar MR, Reznick LG, *et al.* Analysis of Total Time Requirements of
540 Electronic Health Record Use by Ophthalmologists Using Secondary EHR Data. *AMIA Annu*
541 *Symp Proc* 2018;490–7.

542 59 Wang JK, Ouyang D, Hom J, *et al.* Characterizing electronic health record usage patterns of
543 inpatient medicine residents using event log data. *PLoS ONE* 2019;14:e0205379.

544 60 Goldstein IH, Hwang T, Gowrisankaran S, *et al.* Changes in Electronic Health Record Use
545 Time and Documentation over the Course of a Decade. *Ophthalmology* 2019;126:783–91.

546 61 Senathirajah Y, Kaufman D, Bakken S. User-composable Electronic Health Record
547 Improves Efficiency of Clinician Data Viewing for Patient Case Appraisal: A Mixed-
548 Methods Study. *EGEMS (Wash DC)* 2016;4:1176.

549 62 Orenstein EW, Rasooly IR, Mai MV, *et al.* Influence of simulation on electronic health
550 record use patterns among pediatric residents. *J Am Med Inform Assoc* 2018;25:1501–6.

551 63 Zhang W, Gunter CA, Liebovitz D, *et al.* Role prediction using Electronic Medical Record
552 system audits. *AMIA Annu Symp Proc* 2011;858–67.

553 64 Chen Y, Patel MB, McNaughton CD, *et al.* Interaction patterns of trauma providers are
554 associated with length of stay. *J Am Med Inform Assoc* 2018;25:790–9.

555 65 Malin B, Nyemba S, Paulett J. Learning relational policies from electronic health record
556 access logs. *J Biomed Inform* 2011;44:333–42.

557 66 Gray JE, Feldman H, Reti S, *et al.* Using Digital Crumbs from an Electronic Health Record
558 to identify, study and improve health care teams. *AMIA Annu Symp Proc* 2011;491–500.

559 67 Adler-Milstein J, Huckman RS. The impact of electronic health record use on physician
560 productivity. *Am J Manag Care* 2013;19:SP345-352.

561 68 Hripcsak G, Vawdrey DK, Fred MR, *et al.* Use of electronic clinical documentation: time
562 spent and team interactions. *J Am Med Inform Assoc* 2011;18:112–7.

563 69 Grando A, Groat D, Furniss SK, *et al.* Using Process Mining Techniques to Study
564 Workflows in a Pre-operative Setting. *AMIA Annu Symp Proc* 2017;790–9.

565 70 Read-Brown S, Hribar MR, Reznick LG, *et al.* Time Requirements for Electronic Health
566 Record Use in an Academic Ophthalmology Center. *JAMA Ophthalmol* 2017;135:1250–7.

567 71 Tai-Seale M, Olson CW, Li J, *et al.* Electronic Health Record Logs Indicate That Physicians
568 Split Time Evenly Between Seeing Patients And Desktop Medicine. *Health Aff (Millwood)*
569 2017;36:655–62.

570 72 Arndt BG, Beasley JW, Watkinson MD, *et al.* Tethered to the EHR: Primary Care Physician
571 Workload Assessment Using EHR Event Log Data and Time-Motion Observations. *Ann*
572 *Fam Med* 2017;15:419–26.

573 73 Kannampallil TG, Denton CA, Shapiro JS, *et al.* Efficiency of Emergency Physicians:
574 Insights from an Observational Study using EHR Log Files. *Appl Clin Inform* 2018;9:99–
575 104.

576 74 Ben-Assuli O, Leshno M, Shabtai I. Using electronic medical record systems for admission
577 decisions in emergency departments: examining the crowdedness effect. *J Med Syst*
578 2012;36:3795–803.

579 75 Ben-Assuli O, Shabtai I, Leshno M. The impact of EHR and HIE on reducing avoidable
580 admissions: controlling main differential diagnoses. *BMC Med Inform Decis Mak*
581 2013;13:49.

582 76 Ben-Assuli O, Shabtai I, Leshno M. Using electronic health record systems to optimize
583 admission decisions: the Creatinine case study. *Health Informatics J* 2015;21:73–88.

584 77 Wanderer JP, Gruss CL, Ehrenfeld JM. Using Visual Analytics to Determine the Utilization
585 of Preoperative Anesthesia Assessments. *Appl Clin Inform* 2015;6:629–37.

586 78 Soh JY, Jung S-H, Cha WC, *et al.* Variability in Doctors' Usage Paths of Mobile Electronic
587 Health Records Across Specialties: Comprehensive Analysis of Log Data. *JMIR Mhealth*
588 *Uhealth* 2019;7:e12041.

589 79 Gilleland M, Komis K, Chawla S, *et al.* Resident duty hours in the outpatient electronic
590 health record era: inaccuracies and implications. *J Grad Med Educ* 2014;6:151–4.

591 80 Hanauer DA, Zheng K, Commiskey EL, *et al.* Computerized prescriber order entry
592 implementation in a physician assistant-managed hematology and oncology inpatient
593 service: effects on workflow and task switching. *J Oncol Pract* 2013;9:e103-114.

594 81 Coleman JJ, Hodson J, Thomas SK, *et al.* Temporal and other factors that influence the time
595 doctors take to prescribe using an electronic prescribing system. *J Am Med Inform Assoc*
596 2015;22:206–12.

597 82 Amroze A, Field TS, Fouayzi H, *et al.* Use of Electronic Health Record Access and Audit
598 Logs to Identify Physician Actions Following Noninterruptive Alert Opening: Descriptive
599 Study. *JMIR Med Inform* 2019;7:e12650.

600 83 Chen Y, Xie W, Gunter CA, *et al.* Inferring Clinical Workflow Efficiency via Electronic
601 Medical Record Utilization. *AMIA Annu Symp Proc* 2015;416–25.

602 84 Yan C, Chen Y, Li B, *et al.* Learning Clinical Workflows to Identify Subgroups of Heart
603 Failure Patients. *AMIA Annu Symp Proc* 2016;1248–57.

604 85 Shine D, Pearlman E, Watkins B. Measuring resident hours by tracking interactions with the
605 computerized record. *Am J Med* 2010;123:286–90.

606 86 Ouyang D, Chen JH, Krishnan G, *et al.* Patient Outcomes when Housestaff Exceed 80 Hours
607 per Week. *Am J Med* 2016;129:993-999

608 87 Dziorny AC, Orenstein EW, Lindell RB, *et al.* Automatic Detection of Front-Line Clinician
609 Hospital Shifts: A Novel Use of Electronic Health Record Timestamp Data. *Appl Clin*
610 *Inform* 2019;10:28–37.

611 88 Wu DTY, Smart N, Ciemins EL, *et al.* Using EHR audit trail logs to analyze clinical
612 workflow: A case study from community-based ambulatory clinics. *AMIA Annu Symp Proc*
613 2017;1820–7.

614 89 Chen Y, Kho AN, Liebovitz D, *et al.* Learning bundled care opportunities from electronic
615 medical records. *J Biomed Inform* 2018;77:1–10.

616 90 Karp EL, Freeman R, Simpson KN, *et al.* Changes in Efficiency and Quality of Nursing
617 Electronic Health Record Documentation After Implementation of an Admission Patient
618 History Essential Data Set. *Comput Inform Nurs* 2019;37:260–5.

619 91 Hribar MR, Read-Brown S, Reznick L, *et al.* Secondary Use of EHR Timestamp data:
620 Validation and Application for Workflow Optimization. *AMIA Annu Symp Proc* 2015;1909–
621 17.

622 92 Hribar MR, Biermann D, Read-Brown S, *et al.* Clinic Workflow Simulations using
623 Secondary EHR Data. *AMIA Annu Symp Proc* 2016;647–56.

624 93 Hribar MR, Read-Brown S, Reznick L, *et al.* Evaluating and Improving an Outpatient Clinic
625 Scheduling Template Using Secondary Electronic Health Record Data. *AMIA Annu Symp*
626 *Proc* 2017;921–9.

627 94 Hribar MR, Read-Brown S, Goldstein IH, *et al.* Secondary use of electronic health record
628 data for clinical workflow analysis. *J Am Med Inform Assoc* 2018;25:40–6.

629 95 Hribar MR, Huang AE, Goldstein IH, *et al.* Data-Driven Scheduling for Improving Patient
630 Efficiency in Ophthalmology Clinics. *Ophthalmology* 2019;126:347–54.

631 96 Hirsch AG, Jones JB, Lerch VR, *et al.* The electronic health record audit file: the patient is
632 waiting. *J Am Med Inform Assoc* 2017;24:e28–34.

633 97 Goldstein IH, Hribar MR, Sarah R-B, *et al.* Quantifying the Impact of Trainee Providers on
634 Outpatient Clinic Workflow using Secondary EHR Data. *AMIA Annu Symp Proc* 2017;760–
635 9.

- 636 98 Goldstein IH, Hribar MR, Read-Brown S, *et al.* Association of the Presence of Trainees With
637 Outpatient Appointment Times in an Ophthalmology Clinic. *JAMA Ophthalmol*
638 2018;136:20–6.
- 639 99 Vawdrey DK, Wilcox LG, Collins S, *et al.* Awareness of the Care Team in Electronic Health
640 Records. *Appl Clin Inform* 2011;2:395–405.
- 641 100 Chen Y, Lorenzi N, Nyemba S, *et al.* We work with them? Healthcare workers
642 interpretation of organizational relations mined from electronic health records. *Int J Med*
643 *Inform* 2014;83:495–506.
- 644 101 Soulakakis ND, Carson MB, Lee YJ, *et al.* Visualizing collaborative electronic health record
645 usage for hospitalized patients with heart failure. *J Am Med Inform Assoc* 2015;22:299–311.
- 646 102 Chen Y, Lorenzi NM, Sandberg WS, *et al.* Identifying collaborative care teams through
647 electronic medical record utilization patterns. *J Am Med Inform Assoc* 2017;24:e111–20.
- 648 103 Yao N, Zhu X, Dow A, *et al.* An exploratory study of networks constructed using access
649 data from an electronic health record. *J Interprof Care* 2018;32:1–8.
- 650 104 Durojaiye AB, Levin S, Toerper M, *et al.* Evaluation of multidisciplinary collaboration in
651 pediatric trauma care using EHR data. *J Am Med Inform Assoc* 2019;26:506–15.
- 652 105 Zhu X, Tu S-P, Sewell D, *et al.* Measuring electronic communication networks in virtual
653 care teams using electronic health records access-log data. *Int J Med Inform* 2019;128:46–
654 52.
- 655 106 Kim E-H, Stolyar A, Lober WB, *et al.* Challenges to using an electronic personal health
656 record by a low-income elderly population. *J Med Internet Res* 2009;11:e44.
- 657 107 Cimino JJ, Patel VL, Kushniruk AW. What do patients do with access to their medical
658 records? *Stud Health Technol Inform* 2001;84:1440–4.
- 659 108 Weingart SN, Rind D, Tofias Z, *et al.* Who uses the patient internet portal? The PatientSite
660 experience. *J Am Med Inform Assoc* 2006;13:91–5.
- 661 109 Yamin CK, Emani S, Williams DH, *et al.* The digital divide in adoption and use of a
662 personal health record. *Arch Intern Med* 2011;171:568–74.
- 663 110 Bakker A. Access to EHR and access control at a moment in the past: a discussion of the
664 need and an exploration of the consequences. *Int J Med Inform* 2004;73:267–70.
- 665 111 Boxwala AA, Kim J, Grillo JM, *et al.* Using statistical and machine learning to help
666 institutions detect suspicious access to electronic health records. *J Am Med Inform Assoc*
667 2011;18:498–505.
- 668 112 Chen Y, Malin B. Detection of Anomalous Insiders in Collaborative Environments via
669 Relational Analysis of Access Logs. *CODASPY* 2011;63–74.

- 670 113 Chen Y, Nyemba S, Malin B. Detecting Anomalous Insiders in Collaborative Information
671 Systems. *IEEE Trans Dependable Secure Comput* 2012;9:332–44.
- 672 114 Menon AK, Jiang X, Kim J, *et al.* Detecting Inappropriate Access to Electronic Health
673 Records Using Collaborative Filtering. *Mach Learn* 2014;95:87–101.
- 674

675 **FIGURE LEGENDS**

676 Figure 1: Article inclusion criteria


The article...

1. is **peer reviewed**
2. reports **original research**
3. is a **full paper** rather than an abstract
4. **studies use of an EHR** rather than related technologies such as Health Information Exchanges, Personal Health Records, and mobile health apps
5. **uses EHR audit log data** rather than other logs or EHR data such as admission and discharge timestamps from the patient record
6. involves **secondary use** of EHR audit logs rather than using audit logs for their original purpose of access control

677

678

679 Figure 2: Article review process



680

681




Figure 3: A) Audit logs track actions EHR users perform in patient records. Here we show a simplified example of an audit log for one provider performing actions (e.g., “View Problem List”) in three different patient records. We have already mapped these actions to three higher-level clinical activities (record review, orders, documentation). B) Audit logs can be used to compute a variety of measures including simple measures such as 1) action counts, 2) higher-level activity counts and 3) activity durations. These base measures may be used to create more complex models and measures such as 4) sequences of activities, 5) clusters of similar activity patterns, and 6) networks of providers based on their access of the same patient records.




Figure 4: Audit log publications over time with publications reporting a time duration highlighted