



Order for severe sepsis/septic shock

Order for one day	Order for continuation
- Admit ward	- Diet.....
Laboratory:	- Record vital signs, I/O, O ₂ saturation
- H/C X 2 ๒๖๓, CBC, Bun, Cr, Electrolyte, LFT, PT, PTT, INR, DTX	<input type="checkbox"/> Retain Foley's catheter
- Serum lactate at 0,4 and 12 hour	Medication:
<input type="checkbox"/> U/A, Urine C/S	- Empirical antibiotic with stat
<input type="checkbox"/> Sputum G/S, C/S
<input type="checkbox"/> Other.....
.....
- Chest X-ray	- Paracetamol (500)1 tab oral prn for T ≥ 38°c q 6 hrs
- EKG 12 leads	-
Management:
- Load NSS (total as least 30 ml/kg in 2 hrs)
.....
.....
.....
.....
<input type="checkbox"/> Norepinephrine 4 mg + 5%DW 250 ml IV 10 μdrop/min titrate เพิ่ม/ลด ครั้งละ 5 μdrop/min q 5 min keep BP ≥ 90/60 mmHg, MAP ≥ 65 mmHg
- On oxygen keep O2sat ≥ 95%
<input type="checkbox"/> On ET tube No.....depth.....cm
- DTX q 4 hrs. keep 80-180 mg%
- Monitor vital sign q 15 minutes keep BP ≥ 90/60 mmHg, MAP ≥ 65 mmHg
- Keep urine output ≥ 0.5 ml/kg/hr
- Record MEWS at q 4 hr in 24 hour