

Boring but important disclaimers:

- ▶ If you are not getting this from the GitHub repository or the associated Canvas page (e.g. CourseHero, Chegg etc.), you are probably getting the substandard version of these slides Don't pay money for those, because you can get the most updated version for free at

<https://github.com/julianmak/academic-notes>

The repository principally contains the compiled products rather than the source for size reasons.

- ▶ Associated Python code (as Jupyter notebooks mostly) will be held on the same repository. The source data however might be big, so I am going to be naughty and possibly just refer you to where you might get the data if that is the case (e.g. JRA-55 data). I know I should make properly reproducible binders etc., but I didn't...
- ▶ I do not claim the compiled products and/or code are completely mistake free (e.g. I know I don't write Pythonic code). Use the material however you like, but use it at your own risk.
- ▶ As said on the repository, I have tried to honestly use content that is self made, open source or explicitly open for fair use, and citations should be there. If however you are the copyright holder and you want the material taken down, please flag up the issue accordingly and I will happily try and swap out the relevant material.

OCES 2003 : Descriptive Physical Oceanography

(a.k.a. physical oceanography by drawing pictures)

Lecture 8: Mechanical forcing 2 (rotation/Coriolis)

Outline

- ▶ rotation of Earth, **Coriolis “force”** (recall from OCES 2001)
 - rotation axis
 - consequences for flow
 - **Rossby number** + **geostrophic balance**
- ▶ **thermal wind balance**
 - hydrostatic (vertical) + geostrophic (horizontal) balance
 - SSH anomaly example revisited (see Lec. 6 + 7)

Key terms: Coriolis “force”, Rossby number, geostrophic balance/flow, thermal wind (balance)

Recap: equations of motion

Denoting $\mathbf{u} = (u, v)$ and $\mathbf{u}_3 = (u, v, w)$, to numerous approximations (!!!) (see OCES 3203) ocean dynamics is governed by

$$\rho_0 \left(\frac{\partial \mathbf{u}}{\partial t} + \mathbf{u} \cdot \nabla \mathbf{u} + 2\boldsymbol{\Omega} \times \mathbf{u} \right) = -\nabla p + F_u + D_u \quad (1)$$

$$\frac{\partial p}{\partial z} = -\rho g \quad (2)$$

$$\nabla \cdot \mathbf{u}_3 = 0 \quad (3)$$

$$\left(\frac{\partial T}{\partial t} + \mathbf{u}_3 \cdot \nabla T \right) = F_T + D_T \quad (4)$$

$$\left(\frac{\partial S}{\partial t} + \mathbf{u}_3 \cdot \nabla S \right) = F_S + D_S \quad (5)$$

$$\rho = \rho(T, S, p) \quad (6)$$

Respectively, (1) momentum equation, (2) hydrostatic balance, (3) incompressibility, (4) temperature equation, (5) salinity equation, and (6) equation of state (EOS)

Recap: hydrostatic pressure




Figure: Schematic of hydrostatic pressure

- **hydrostatic approximation:**
pressure **approximately equal** to weight above when static
→ **weight** is $F = mg$ so for force balance,


$$F = mg = g \int_{-z}^{z_{\text{atm}}} \rho \, dz = p ,$$

with $g \approx 9.81 \, \text{m s}^{-2}$

→ if $\rho = \text{const}$ then $p = \rho g z + p_{\text{atm}}$

$$\frac{\partial p}{\partial z} = -\rho g$$


Recap: pressure gradients and flows



- ▶ assuming hydrostatic balance, water moves from $+\Delta p$ to $-\Delta p$ because there is a **net force** (negative pressure gradient $-\nabla p$)
→ important for **geostrophic flows**

Figure: Horizontal effect because of hydrostatic pressure.

Geostrophic flows: atmosphere



Winds do not go direct from high to low p ? (more on wind patterns next Lec.)

Geostrophic flows: atmosphere

日期/Date: 14.10.2020 香港時間/HK Time: 14:00 香港天文台 Hong Kong Observatory




Figure: Atmospheric weather chart with isobars (in units of hPa = 100 Pa = 1 mbar) and wind directions. From HKO.

- note that flow doesn't go in the direction of $-\nabla p!$
→ **along** rather than **across** isobars (Coriolis effect, see next Lec.)

Geostrophic flows: ocean




Figure: Time-mean global SSH (also called **mean dynamic topography**, with time-mean currents drawn on (notice the orientation around high/low SSH regions). Modified from Rio *et al* (2011), J. Geophys. Res: Oceans.

- contours of SSH related to isobars via **hydrostatic balance**
 - flow is **along** rather than **across** isobars (**Coriolis effect**, see next Lec.)

Coriolis effect

The Earth rotates around the **rotation axis**

- ▶ the **geographical North** (as opposed to magnetic north)
- ▶ **rate** of rotation is the **angular frequency** Ω (units: s^{-1}), with

$$\Omega = \frac{2\pi}{T}$$

→ T is the **period** (see again in Lec. 15 - 18), time needed to do one rotation (2π radians or 360°)

→ for Earth (units!)

$$\Omega = \frac{2\pi}{3600 \times 24} \approx 7.29 \times 10^{-5} \text{ s}^{-1}$$




Figure: Rotation axis and angular frequency Ω .

Coriolis effect: co-ordinates




Figure: Mis-alignment of Ω and e_z used locally for depth.

- ▶ for a spherical Earth we take rotation axis to be z-axis, i.e. $\Omega = \Omega e_z$ (this a vector), but locally, z is depth...
- ▶ introduce the latitudinally varying **Coriolis parameter**

$$f = 2\Omega \sin(\text{latitude})$$

- ▶ to take into account of mis-alignments between Ω and the local e_z for depth
 - Coriolis = $-2\Omega \times u$ (global case, z is North)
 - Coriolis $-f e_z \times u$ (local case, z is depth) (mostly going to use this one)

Frame of reference

(sort of) a demonstration of a **frame of reference**

- ▶ one could work in global picture (with Ω and z being North) or local picture (with f and z being depth)
→ change in point-of-view, **co-ordinate system**

Frame of reference

(sort of) a demonstration of a **frame of reference**

- ▶ one could work in global picture (with Ω and z being North) or local picture (with f and z being depth)
→ change in point-of-view, **co-ordinate system**
- ▶ one point-of-view might be “easier” than other
→ e.g. tracking change in meaning of z vs. remembering f varies in latitude

Frame of reference

(sort of) a demonstration of a **frame of reference**

- ▶ one could work in global picture (with Ω and z being North) or local picture (with f and z being depth)
→ change in point-of-view, **co-ordinate system**
- ▶ one point-of-view might be “easier” than other
→ e.g. tracking change in meaning of z vs. remembering f varies in latitude
- ▶ underlying physics should still be the **same**, although description might **look** different (e.g. Ω vs. $f e_z$)

Frame of reference

(sort of) a demonstration of a **frame of reference**

- ▶ one could work in global picture (with Ω and z being North) or local picture (with f and z being depth)
→ change in point-of-view, **co-ordinate system**
- ▶ one point-of-view might be “easier” than other
→ e.g. tracking change in meaning of z vs. remembering f varies in latitude
- ▶ underlying physics should still be the **same**, although description might **look** different (e.g. Ω vs. $f\mathbf{e}_z$)
- ▶ **freedom in choice of frame!**
→ e.g. frame rotating with the planet, others...
→ fine as long as we keep **consistency**

Coriolis effect

You can think of it as a “hack”

- ▶ Newton's laws are formulated for **inertial frames**
→ non-accelerating frames

Coriolis effect

You can think of it as a “hack”

- ▶ Newton's laws are formulated for **inertial frames**
→ non-accelerating frames
- ▶ a rotating frame *is* an accelerating frame (so non-inertial)
→ change in the flow vector \Leftrightarrow acceleration

can't use Newton's laws then?

Coriolis effect

You can think of it as a “hack”

- ▶ Newton's laws are formulated for **inertial frames**
→ non-accelerating frames
- ▶ a rotating frame *is* an accelerating frame (so non-inertial)
→ change in the flow vector \Leftrightarrow acceleration

can't use Newton's laws then?

- ▶ “Hack”: put in a **fictitious forces** to compensate the fact we are not in an inertial frame
→ can then treat non-inertial frames **as if it were inertial**
→ proceed as normal with that caveat

Coriolis effect

Earth's **spin** has a major influence on large-scale winds

- ▶ extra “force” from the spinning: **Coriolis** “force” $2\boldsymbol{\Omega} \times \boldsymbol{u}$

(Gaspard-Gustave de Coriolis, 1792-1843, French mathematician)

→ apparent deflection \Leftrightarrow a net force on it

- ▶ apparent “force” because not being in **inertial frame**
 - can think of it arising only because of perspective...
 - note Coriolis “force” does **no work** (see assignment)

- ▶ try it yourself! (seriously this really helped me...)

→ on a piece of paper, try drawing a straight line while rotating the paper underneath

→ something similar but on e.g. a basketball

Coriolis effect




Figure: Schematic of apparent deflection from Coriolis "force". From the drawer perspective, the drawer is doing a straight line and sees a straight line. By from observer's perspective, there is a deflection. The action is the same, but to describe it from the observer's point of view, we need to additionally describe this apparently deflection arising from the system's rotation.

- on a piece of paper, try drawing a straight line while rotating the paper underneath

Coriolis effect




Figure: Schematic of apparent deflection from Coriolis “force”, from Vallis (2011).

- ▶ apparent deflection **to the right** in NH
→ looking down from North Pole, rotating **anti-clockwise**
- ▶ apparent deflection **to the left** in SH
→ looking down from South Pole, now rotating **clockwise**

Rossby number

Note there is a competition between **fluid velocity** (intended path) and **rotation**

Rossby number

Note there is a competition between **fluid velocity** (intended path) and **rotation**

- ▶ matter of **time-scales**

- if fluid moves quickly relative to system spinning (**inertial period**), then Coriolis influence small

- if fluid moves relatively slowly, more time for Coriolis effect to act

Rossby number

Note there is a competition between **fluid velocity** (intended path) and **rotation**

► matter of **time-scales**

→ if fluid moves quickly relative to system spinning (**inertial period**), then Coriolis influence small

→ if fluid moves relatively slowly, more time for Coriolis effect to act

► measured by the **Rossby number**: for $f = 2\Omega \sin(\text{latitude})$,
(Carl-Gustav Rossby, 1898-1957, Swedish meteorologist)

$$\text{Ro} = \frac{U/L}{f} = \frac{1/T}{f} = \frac{\text{advection}}{\text{rotation}}$$

→ note f decreases to zero at EQ (mis-alignment of rotation axis, recall $-2\mathbf{\Omega} \times \mathbf{u}$)

Rossby number

$$\text{Ro} = \frac{U/L}{f} = \frac{1/T}{f} = \frac{\text{advection}}{\text{rotation}}$$

- For large-scale motion in Earth's atmosphere in mid-lats (say 50°N) and $\Omega = 2\pi/\text{day}$,

$$\text{Ro} = \frac{10 \text{ m s}^{-1}/1000 \text{ km}}{2 \times 2\pi \text{ day}^{-1} \times \sin(50^\circ)} \approx \frac{10^1 \times 10^{-6}}{10^{-4}} = 0.1,$$

so rotational effects dominant

Rossby number

$$\text{Ro} = \frac{U/L}{f} = \frac{1/T}{f} = \frac{\text{advection}}{\text{rotation}}$$

- For large-scale motion in Earth's atmosphere in mid-lats (say 50°N) and $\Omega = 2\pi/\text{day}$,

$$\text{Ro} = \frac{10 \text{ m s}^{-1}/1000 \text{ km}}{2 \times 2\pi \text{ day}^{-1} \times \sin(50^\circ)} \approx \frac{10^1 \times 10^{-6}}{10^{-4}} = 0.1,$$

so rotational effects dominant

Rossby number

$$Ro = \frac{U/L}{f} = \frac{1/T}{f} = \frac{\text{advection}}{\text{rotation}}$$

- For large-scale motion in Earth's atmosphere in mid-lats (say 50°N) and $\Omega = 2\pi/\text{day}$,

$$Ro = \frac{10 \text{ m s}^{-1}/1000 \text{ km}}{2 \times 2\pi \text{ day}^{-1} \times \sin(50^\circ)} \approx \frac{10^1 \times 10^{-6}}{10^{-4}} = 0.1,$$

so rotational effects dominant

- $Ro \sim O(0.01)$ for Jupiter <https://www.youtube.com/watch?v=exkeE6aIlpc>
→ very fast flows but also very large length scales

Rossby number

$$\text{Ro} = \frac{U/L}{f} = \frac{1/T}{f} = \frac{\text{advection}}{\text{rotation}}$$

- ▶ For large-scale motion in Earth's atmosphere in mid-lats (say 50°N) and $\Omega = 2\pi/\text{day}$,

$$\text{Ro} = \frac{10 \text{ m s}^{-1}/1000 \text{ km}}{2 \times 2\pi \text{ day}^{-1} \times \sin(50^\circ)} \approx \frac{10^1 \times 10^{-6}}{10^{-4}} = 0.1,$$

so rotational effects dominant

- ▶ $\text{Ro} \sim O(0.01)$ for Jupiter <https://www.youtube.com/watch?v=exkeE6aIlpc>
→ very fast flows but also very large length scales
- ▶ flows in the interior of the Sun, $\text{Ro} = O(1)$
→ Sun not spinning too fast, rotationally influenced

Rossby number

$$Ro = \frac{U/L}{f} = \frac{1/T}{f} = \frac{\text{advection}}{\text{rotation}}$$

- For large-scale motion in Earth's atmosphere in mid-lats (say 50°N) and $\Omega = 2\pi/\text{day}$,

$$Ro = \frac{10 \text{ m s}^{-1}/1000 \text{ km}}{2 \times 2\pi \text{ day}^{-1} \times \sin(50^\circ)} \approx \frac{10^1 \times 10^{-6}}{10^{-4}} = 0.1,$$

so rotational effects dominant

- $Ro \sim O(0.01)$ for Jupiter <https://www.youtube.com/watch?v=exkeE6aIlpc>
→ very fast flows but also very large length scales
- flows in the interior of the Sun, $Ro = O(1)$
→ Sun not spinning too fast, rotationally influenced
- ? what about in a toilet bowl? (see assignment)

Geostrophic flow

After **non-dimensionalisation** (see OCES 3301 or ask me), momentum equation is (other numbers appear for forcing + dissipation, assume small; see Lec 9 + 10)

$$\text{Ro}\rho \left(\frac{\partial \mathbf{u}}{\partial t} + \mathbf{u} \cdot \nabla \mathbf{u} \right) + \rho 2\boldsymbol{\Omega} \times \mathbf{u} = -\nabla p + \dots$$

If $\text{Ro} \ll 1$, dominant force balance is:

$$2\boldsymbol{\Omega} \times \mathbf{u}_g = -\frac{1}{\rho} \nabla p$$

► geostrophic balance

→ given p and $\boldsymbol{\Omega}$, \mathbf{u}_g (the **geostrophic flow**) has to be **something** so resulting forces balance

what is the implied velocity \mathbf{u} then?

Geostrophic flow: rationalisation




Figure: Geostrophic balance and resulting geostrophic flow u_g in Northern Hemisphere. Note u_g is **along** (rather than **across**) isobars.

- in NH top-down view, rotation is **anti**-clockwise

Geostrophic flow: rationalisation




Figure: Geostrophic balance and resulting geostrophic flow u_g in Northern Hemisphere. Note u_g is **along** (rather than **across**) isobars.

- in NH, deflection to the **right** of $-\nabla p$

Geostrophic flow: rationalisation





Figure: Geostrophic balance and resulting geostrophic flow u_g in Northern Hemisphere. Note u_g is **along** (rather than **across**) isobars.

- for **force balance**, Coriolis has to be opposite of $-\nabla p$

Geostrophic flow: rationalisation



- ▶ or, really, **because** of force balance, u_g has to be to the right of $-\nabla p$ in NH
- ▶ $\Omega \sim +e_z$ (Earth NH), $-\nabla p \sim e_y$, so

$$-2\Omega \times u = -\frac{1}{\rho} \nabla p \quad \Rightarrow \quad -e_z \times u_g \sim e_y \quad \text{or} \quad u_g \times e_z \sim e_y,$$

so $u \sim +e_x$ only possibility, i.e. to the E (right of N is E)

Geostrophic flow: rationalisation




Figure: Geostrophic balance and resulting geostrophic flow u_g in Southern Hemisphere. Note u_g is **along** (rather than **across**) isobars.

- in SH top-down view, rotation is **clockwise**

Geostrophic flow: rationalisation




Figure: Geostrophic balance and resulting geostrophic flow u_g in Southern Hemisphere. Note u_g is **along** (rather than **across**) isobars.

- u_g to the **left** of $-\nabla p$ (same arguments as but $\underline{\Omega} \rightarrow -\underline{\Omega}$)

Geostrophic flow

Suppose hydrostatic as well as geostrophic balance:




Figure: Schematic for an anti-cyclonic (warm core) eddy.

- ▶ bulge ($+\Delta h$) so $+\Delta p$ in the centre
 $\rightarrow -\nabla p$ points **away** from region
- ▶ **geostrophic current** u_g to the **right** of $-\nabla p$ (since NH)
 \Rightarrow **clockwise** around bulge
- ▶ **opposite** sense to planet rotation f (in NH),
anti-cyclonic eddy (in NH)
 \rightarrow other direction in SH (since $f < 0$)

Geostrophic flow




Figure: Schematic for a cyclonic (cold core) eddy.

- ▶ depression ($-\Delta h$) so $-\Delta p$ in the centre
→ $-\nabla p$ points **into** region
- ▶ **geostrophic current** u_g to the **right** of $-\nabla p$ (since NH)
⇒ **anti-clockwise** around depression
- ▶ **same** sense as rotation of planet f (in NH), **cyclonic eddy** (in NH)
→ other direction in SH (since $f < 0$)

Geostrophic flow

Above was for NH, for SH:

- ▶ pressure and $-\nabla p$ still related to hydrostatic balance
- ▶ deflection is to the **left** (cf. opposite rotation sense)
- ▶ u_g is now
 - anti-clockwise around bulge
 - clockwise around depression
- ▶ BUT!
 - bulges are still **anti-cyclonic**
 - depressions are still **cyclonic**

Geostrophic flow

Above was for NH, for SH:

- ▶ pressure and $-\nabla p$ still related to hydrostatic balance
- ▶ deflection is to the **left** (cf. opposite rotation sense)
- ▶ u_g is now
 - anti-clockwise around bulge
 - clockwise around depression
- ▶ BUT!
 - bulges are still **anti-cyclonic**
 - depressions are still **cyclonic**
- ▶ cf. atmosphere, low pressures \leftrightarrow depressions \leftrightarrow cyclonic
 - in atmosphere low pressures are **convergence zones**, related to unsettled weather (and vice-versa in high pressures) (relation to Ekman up/downwelling next Lec.)

Geostrophic flow from SSH




Figure: Time-mean global SSH (also called **mean dynamic topography**, with time-mean currents drawn on (notice the orientation around high/low SSH regions). Modified from Rio *et al* (2011), J. Geophys. Res: Oceans.

- contours of SSH related to isobars via **hydrostatic balance**
 - flow is **along** rather than **across** isobars (**Coriolis effect**, see next Lec.)

Geostrophic flow from SSH




Figure: SSH and inferred currents from AVISO satellite altimeter data. Figure 4 of Son *et al.* (2014), *Biogeosciences*.

- ▶ observed SSH from **AVISO** (see Lec. 20) + inferred geostrophic velocities near Japan
→ anti-cyclonic around positive SSH anomalies (anti-clockwise in NH)
- ▶ note current is strongest where gradients are large (the arrow lengths)

Summary

geostrophic flow goes to the right of pressure gradient in NH

- ▶ flip this in SH (because rotation is “reversed”)
 - again, **Coriolis effect is frame dependent**, i.e. depend on your point of view, a **pseudo-force**
 - Coriolis force does **no work** (cf. Lec 5 + 6; see assignment)
- ▶ **anti-cyclonic** eddies \sim bulges in SSH (in SH and NH)
 - rotation same sense as planet, $\nabla \times \mathbf{u} \sim f$
 - clockwise in NH (because $f > 0$), reverse in SH
- ▶ **cyclonic** eddies \sim depressions in SSH (in SH and NH)
 - rotation opposite sense as planet, $\nabla \times \mathbf{u} \sim -f$
 - anti-clockwise in NH (because $f > 0$), reverse in SH

Summary

Turns out the deflection aspect is also (largely) true for wind forced flows (see next Lec.):

Ekman transport is to the right of the wind in NH (flip in SH)

→ wind drives flow at surface in direction of wind...

→ but flow needs to turn 90° (or $\pm\pi/2$) at depth?

► Ekman spirals