
Schrodinger

Release 0.0.1

Group Schrodinger

Feb 02, 2023

CONTENTS

SYNOPSIS:

Impact craters are the most ubiquitous surface feature on rocky planetary bodies. Crater number density can be used to estimate the age of the surface: the more densely cratered the terrain, the older the surface.

When independent absolute ages for a surface are available for calibration of crater counts, as is the case for some lava flows and regions of the Moon, crater density can be used to estimate an absolute age of the surface.

Crater detection and counting has traditionally been done by laborious manual interrogation of images of a planetary surface taken by orbiting spacecraft (Robbins and Hynek, 2012; Robbins, 2019). However, the size frequency distribution of impact craters is a steep negative power-law, implying that there are many small craters for each larger one. For example, for each 1-km crater on Mars, there are more than a thousand 100-m craters. With the increased fidelity of cameras on orbiting spacecraft, the number of craters visible in images of remote surfaces has become so large that manual counting is unfeasible. Furthermore, manual counting can be time consuming and subjective (Robbins et al., 2014). This motivates the need for automated crater detection and counting algorithms (DeLatta et al., 2019). Recent work has shown that widely used object detection algorithms from computer vision, such as the YOLO (You Only Look Once) object detection algorithm (Redmon et al., 2016; Jocher et al., 2021), can be effective for crater detection on Mars (Benedix et al., 2020; Lagain et al., 2021) and the Moon (Fairweather et al., 2022).

This tool provides the yolov5 models used for detecting a wide range of sizes of crater both on Mars and on Moon dataset.

MODEL DESCRIPTION

2.1 Inputs

Our tool takes input as one or loads of images of the surface of a planet, as well as optional inputs of the planet, planet radius, location and physical size of the image. Our tool has been tested for Mars and the Moon dataset.

2.2 Outputs

Our tool output a list of all the bounding boxes for craters detected in each image.

The additional requirements are the following:

Generate a visualisation of the original image with annotated bounding boxes.

When a real-world image size and location is provided for the image, our tool can also provide physical locations and size for each crater.

When ground truth labels are provided, our tool should also determine the number of True Positive, False Positive and False Negative detections and return these values for each image.

Finally, the tool can also plot a comparison of the ground truth bounding boxes and the model detection bounding boxes.

2.3 Visualisation

Users can visualize the following:

The original input image without annotations.

The original input image with bounding boxes for craters detected by the yolov5.

The original input image with bounding boxes for both the results of the yolov5 and for the ground truth bounding boxes.

A separate plot of the cumulative crater size-frequency distribution of detected craters.

Performance statistics including the number of True Positive, False Negative and False Positive detections.

2.4 Additional sections

You should expand this documentation to include explanatory text for all components of your tool.