

The temporal assembly of plant-pollinator networks following restoration

Lauren C. Ponisio¹, Marilia P. Gaiarsa², Claire Kremen¹

1. Department of Environmental Science, Policy, and Management
University of California, Berkeley
130 Mulford Hall
Berkeley, California, USA
94720
2. Departamento de Ecologia
Universidade de Sao Paulo
Sao Paulo, SP, Brazil
05508-900

Abstract

The structure of networks is related to ability of communities to maintain function in the face of species extinction. Understanding network structure and how it relates to network disassembly, therefore, is a priority for system-level conservation biology. We explore the assembly of plant-pollinator communities on native plant restorations in the Central Valley of California. The assembling communities are paired with un-restored field margins (controls) and mature (non-assembling) hedgerows. We determine whether there are change points in the assembly of the communities where the network undergoes significant reorganization. We also ask how are the individual species changing their interaction patterns? What does this mean for the topology/resilience of the network? We also attempted to adapt a financial model to mutualistic networks. Our biggest difficulty with this approach was to translate the price term to mutualistic systems. We explored a range of approaches, such as number of visits a species performs. However, it seems that financial systems cannot be easily translated to mutualistic systems. In addition, we used a Changing Point Detection Algorithm to assess whether the different communities went through a critical reorganization on their interaction patterns. We were able to identify some changing points in the communities, and also to explore some general patterns commonly used to describe ecological networks. For example, on the network level, networks become increasingly modular and less nested, whereas on the species level, species become more specialized, as resources become more reliable.

Keywords: changing points, temporal networks, hedgerows, species interactions

Introduction

- The structure of networks is related to ability of communities to maintain function in the face of species extinction.
- A key restoration aim is to facilitate assembly of robust networks; thus it is critical to study how restoration influences the assembly of plant-pollinator interactions.
- few theories about how networks assemble, preferential attachment
- To date, only two field studies have examined how networks assemble over time, often using space for time gradients. Studying primary succession along a glacier foreland, Albrecht *et al.* (2010) found that specialization of plants and pollinators decreased while nestedness increased as the community aged. Studying “managed succession” of a clear-cut pine forest, Devoto *et al.* (2012) found changes in network structure were explained by a combination of age, tree density and variation in tree diameter.
- Olesen *et al.* (2008) was investigated day-to-day, temporal assembly of a plant-pollinator network within a season, taking advantage of the extreme seasonality of pollinator communities in Greenland.
- In contrast, in non-successional dynamics, network architecture has been found to remain relatively constant despite high turnover of species and interactions in a community (Olesen *et al.*, 2011; Petanidou *et al.*, 2008).
- core vs. periphery idea (Fang & Huang, 2012; Díaz-Castelazo *et al.*, 2010)

Materials & Methods

Study sites and collection methods

Results

Discussion

Acknowledgments

We would like to thank Leto Peel and Aaron Clauset for their invaluable discussions and for help with the change point analysis. We thank the growers and land owners that allowed us to work on their property. We also appreciate the identification assistance of expert taxonomists Robbin Thorp and Jason Gibbs. This work was supported by funding from the Army Research Office (W911NF-11-1-0361 to CK), the Natural Resources Conservation Service (CIG-69-3A75-12-253, CIG-69-3A75-9-142, CIG-68-9104-6-101 and WLF-69-7482-6-277 to The Xerces Society), the National Science Foundation (DEB-0919128 to CK), The U.S. Department of Agriculture (USDA-NIFA 2012-51181-20105 to Michigan State University). Funding for LCP was provided by an NSF Graduate Research Fellowship and the USDA NIFA Graduate Fellowship. FUNING FOR MARILLIA.

References

Albrecht, M., Riesen, M. & Schmid, B. (2010). Plant–pollinator network assembly along the chronosequence of a glacier foreland. *Oikos*, 119, 1610–1624.

- 61 Devoto, M., Bailey, S., Craze, P. & Memmott, J. (2012). Understanding and planning
62 ecological restoration of plant–pollinator networks. *Ecology letters*, 15, 319–328.
- 63 Díaz-Castelazo, C., Guimaraes Jr, P.R., Jordano, P., Thompson, J.N., Marquis, R.J. & Rico-
64 Gray, V. (2010). Changes of a mutualistic network over time: reanalysis over a 10-year
65 period. *Ecology*, 91, 793–801.
- 66 Fang, Q. & Huang, S.Q. (2012). Relative stability of core groups in pollination networks
67 in a biodiversity hotspot over four years. *PloS one*, 7, e32663.
- 68 Olesen, J.M., Bascompte, J., Elberling, H. & Jordano, P. (2008). Temporal dynamics in a
69 pollination network. *Ecology*, 89, 1573.
- 70 Olesen, J.M., Stefanescu, C. & Traveset, A. (2011). Strong long-term temporal dynamics
71 of an ecological network. *Plos One*, 6.
- 72 Petanidou, T., Kallimanis, S., Tzanopoulos, J., Sgardelis, S. & Pantis, J. (2008). Long-term
73 observation of a pollination network: fluctuation in species and interactions, relative
74 invariance of network structure and implications for estimates of specialization. *Ecol.*
75 *Lett.*, 11, 564–575.