

DISSERTATION

EMERGENT TOPOLOGICAL PHENOMENA IN LOW-D SYSTEMS INDUCED BY GAUGE  
POTENTIALS

Submitted by  
Aidan Winblad  
Department of Physics

In partial fulfillment of the requirements  
For the Degree of Doctor of Philosophy  
Colorado State University  
Fort Collins, Colorado  
Fall 2023

Doctoral Committee:

Advisor: Hua Chen

Richard Eykholt  
Martin Gelfand  
Olivier Pinaud

Copyright by Aidan Winblad 2023

All Rights Reserved

## ABSTRACT

# EMERGENT TOPOLOGICAL PHENOMENA IN LOW-D SYSTEMS INDUCED BY GAUGE POTENTIALS

Abstract goes here

## ACKNOWLEDGEMENTS

I would like to thank the CSU Graduate Student Council and the CSU Graduate School for initiating, commissioning and supporting this project. I would also like to thank Nicole Ramo for her support and ensuring that we followed through with this project to completion. I would like to thank Leif Anderson, who created and supported the previous LaTeX template for a number of years. Although I have never met Leif, his work was invaluable in the creation of this package and has helped many students get their thesis approved by the CSU graduate school. Finally, I would like to thank everyone who helps to contribute to this package. Your work will help many CSU graduate students to create professional, beautiful and compelling theses and dissertations using LaTeX. Last but not least, thank you to the creators and maintainers of  $\text{\LaTeX}$  for creating a fantastic typesetting tool.

## DEDICATION

*I would like to dedicate this dissertation to my dog Zeta.*

# TABLE OF CONTENTS

ABSTRACT . . . . .	ii
ACKNOWLEDGEMENTS . . . . .	iii
DEDICATION . . . . .	iv
LIST OF TABLES . . . . .	vi
LIST OF FIGURES . . . . .	vii
Chapter 1      Introduction . . . . .	1
Chapter 2      Superconducting Triangular Islands as a Platform for Manipulating Majorana Zero Modes . . . . .	2
2.1          Introduction . . . . .	2
2.2          Kitaev Triangle . . . . .	4
2.3          Hollow Triangles . . . . .	7
2.4          Braiding MZM in a small network of triangles . . . . .	14
2.5          Discussion . . . . .	15
Chapter 3      Floquet Landau Levels . . . . .	18
Chapter 4      Conclusion and Discussion . . . . .	19
Appendices . . . . .	20
Appendix Chapter A      Suitable Name . . . . .	20
Bibliography . . . . .	21

## LIST OF TABLES

## LIST OF FIGURES

2.1	Schematics of two triangle structures proposed in this work. (a) Three-site Kitaev triangle with bond-dependent Peierls phases. (b) Hollow triangular island with a uniform vector potential. . . . .	4
2.2	(a) Evolution of the eigenvalues of the 3-site Kitaev triangle along the closed parameter path for $\phi$ on the three edges. (b) MZM wavefunctions at different points of the parameter path. Clockwise from the upper left panel: $\phi_1 \rightarrow \frac{1}{2}(\phi_1 + \phi_2) \rightarrow \phi_2 \rightarrow \phi_3$ . . . . .	7
2.3	(a) Topological phase diagram for a $W = 1$ triangular chain with the Hamiltonian Eq. (2.7) obtained by superimposing the $\mathcal{M}(A, \mu)$ plots of 1D chains with $\mathbf{A} = A\hat{y}$ and $\mathbf{A} = A(\frac{\sqrt{3}}{2}\hat{x} + \frac{1}{2}\hat{y})$ . Color scheme: white— $\mathcal{M} = 1$ , dark blue— $\mathcal{M} = -1$ , light blue— $\mathcal{M} = 0$ (b) Near-gap BdG eigen-energies vs $A$ for a finite triangle with edge length $L = 50$ , $W = 1$ , and $\mu = 1.6$ . . . . .	9
2.4	(a) Spectral flow of a hollow triangle with $W = 1$ , $L = 50$ , $\mu = 1.6$ , and $A = 2.75$ with increasing rotation angle $\varphi$ , defined through $\mathbf{A} = A(-\sin \varphi \hat{x} + \cos \varphi \hat{y})$ . (b-d) BdG eigenfunction $ \Psi ^2$ summed over the two zero modes at $\varphi = 0, \frac{\pi}{6}$ , and $\frac{\pi}{3}$ , respectively. . . . .	11
2.5	(a) Topological phase diagram for a $W = 3$ hollow triangle obtained by overlapping the $\mathcal{M}(A, \mu)$ plots of 1D chains with $\mathbf{A} = A\hat{y}$ and $\mathbf{A} = A(\frac{\sqrt{3}}{2}\hat{x} + \frac{1}{2}\hat{y})$ . Color scheme: white— $\mathcal{M} = 1$ , dark blue— $\mathcal{M} = -1$ , light blue— $\mathcal{M} = 0$ (b) Near-gap BdG eigen-energies vs $A$ for a finite triangle with edge length $L = 50$ , $W = 3$ , and $\mu = 1.6$ . (c) BdG eigenfunction $ \Psi ^2$ summed over the two zero modes at $A = 2.4709$ . . . . .	12
2.6	(a) Spectral flow of a hollow triangle with $W = 3$ , $L = 50$ , $\mu = 1.6$ , and $A = 2.75$ with increasing rotation angle $\varphi$ , defined through $\mathbf{A} = A(-\sin \varphi \hat{x} + \cos \varphi \hat{y})$ . (b-c) BdG eigenfunction $ \Psi ^2$ summed over the two zero modes at $\varphi = 0$ and $\frac{\pi}{3}$ , respectively. . . . .	13
2.7	(a) Spectral flow for the critical step of swapping $\gamma_2$ and $\gamma_3$ in the example of Fig. 5 in the main text, calculated using four corner-sharing triangles of $W = 1$ and $L = 50$ , with $\mu = 1.6$ and $A = 2.6$ . Vector potential for the middle triangle in the bottom row can rotate according to $\mathbf{A} = A(-\sin \varphi \hat{x} + \cos \varphi \hat{y})$ from $\varphi = \frac{\pi}{6}$ to $\frac{\pi}{3}$ , while the other three have fixed $\varphi = 0$ . (b)-(g) BdG eigenfunction $ \Psi ^2$ summed over the four zero modes at equally-spaced points along the rotation path. The black arrow indicates the direction of the vector potential for the bottom middle triangle. . . . .	14
2.8	Representative steps for braiding four MZM in four triangles sharing corners. (a) Initialization of four MZM $\gamma_1, \gamma_2, \gamma_3, \gamma_4$ . All three edges of the bottom-middle and the top triangles are in the trivial phase by e.g. controlling the chemical potential. The bottom-left and bottom-right triangles have $\varphi = 0$ so that their bottom edges are nontrivial. (b) Moving $\gamma_3$ by “switching on” the middle triangle by changing the chemical potential under a fixed vector potential at $\varphi = \frac{\pi}{6}$ , and then turning on the top triangle with similar means except $\varphi = 0$ . (c) Transporting $\gamma_2$ to the right triangle through rotating the vector potential in the middle triangle counterclockwise by $\pi/6$ . (d) Moving $\gamma_3$ to the left triangle by “switching off” the top triangle followed by the middle triangle. . . . .	16





# Chapter 1

## Introduction

EM gauge potential appears in electronic Hamiltonian in CM

1. Review Maxwell theory  $\rightarrow$  gauge potential
2. Minimal coupling  $-i\hbar\nabla \rightarrow -i\hbar\nabla + q\mathbf{A}$  or  $-i\partial_\mu \rightarrow -i\partial_\mu + qA_\mu$
3. TB Hamiltonian and Peierls phase

Topological phenomena in CM considered in thesis

1. (1) Majorana and TSC
  - i Kitaev chain (M— topological invariant). BdG?
  - ii Braiding (Application in TQC)
2. Landau Level and Hofstadter butterfly
  - i solve for LL in 2DEG — why it's topological, chern number, TKNN quantum Hall
  - ii square lattice — hofstadter butterfly ( on other lattices, honeycomb)

## Chapter 2


# Superconducting Triangular Islands as a Platform for Manipulating Majorana Zero Modes

### 2.1 Introduction

For more than twenty years, Majorana zero modes (MZM) in condensed matter systems have been highly sought after due to their potential for serving as building blocks of topological quantum computation, thanks to their inherent robustness against decoherence and non-Abelian exchange statistics [1–5]. MZM were originally proposed to be found in half-quantum vortices of two-dimensional (2D) topological  $p$ -wave superconductors and at the ends of 1D spinless  $p$ -wave superconductors [6, 7]. Whether a pristine  $p$ -wave superconductor [8] has been found is still under debate. However, innovative heterostructures proximate to ordinary  $s$ -wave superconductors have been proposed to behave as effective topological superconductors in both 1D and 2D. These include, for example, semiconductor nanowires subject to magnetic fields [9–11], ferromagnetic atomic spin chains [12–17], 3D topological insulators [18–21], quantum anomalous Hall insulators [22–24], quasi-2D spin-orbit-coupled superconductors with a perpendicular Zeeman field [25–30], and planar Josephson junctions [31–37], etc. It has been a challenging task to decisively confirm the existence of MZM in the various experimental systems due to other competing mechanisms that can potentially result in similar features as MZM do in different probes [34, 35, 38–43]. Other proposals for constructing Kitaev chains through a bottom-up approach, based on, e.g. magnetic tunnel junctions proximate to spin-orbit-coupled superconductors [44], and quantum dots coupled through superconducting links [45–47] are therefore promising. In particular, the recent experiment [47] of a designer minimal Kitaev chain based on two quantum dots coupled through tunable crossed Andreev reflections (CAR) offers a compelling route towards MZM platforms based on exactly solvable building blocks.

In parallel with the above efforts of realizing MZM in different materials systems, scalable architectures for quantum logic circuits based on MZM have also been intensely studied over the past decades. A major proposal among these studies is to build networks of T-junctions, which are minimal units for swapping a pair of MZM hosted at different ends of a junction, that allow braiding-based TQC [5]. Alternatively, networks based on coupled wires forming the so-called tetrons and hexons, aiming at measurement-based logic gate operations [48], have also been extensively investigated. To counter the technical challenges of engineering networks with physical wires or atomic chains, various ideas based on effective Kitaev chains, such as quasi-1D systems in thin films [49], cross Josephson junctions [37], scissor cuts on a quantum anomalous Hall insulator [24], and rings of magnetic atoms [50], etc. have been proposed. However, due to the same difficulty of obtaining or identifying genuine MZM in quasi-1D systems mentioned above, it remains unclear how practical these strategies are in the near future.

In this Letter, we propose an alternative structural unit for manipulating MZM, triangular superconducting islands, motivated by the above challenges associated with wire geometries and by the fact that triangular islands routinely appear spontaneously in epitaxial growth [51] on close-packed atomic surfaces. We first show that a minimal “Kitaev triangle” consisting of three sites hosts MZM at different pairs of vertices controlled by Peierls phases on the three edges [Fig. 2.1 (a)], which can be readily realized using quantum dots. To generalize the minimal model to triangular structures involving more degrees of freedom, we study the topological phase transitions of quasi-1D ribbons driven by Peierls phases, which can be created by magnetic fields or supercurrents [52, 53], and use the resulting phase diagram as a guide to construct finite-size triangles with a hollow interior that host MZM [Fig. 2.1 (b)]. In the end we discuss possible experimental systems that can realize our proposals and scaled-up networks of triangles for implementing braiding operations of MZM.



**Figure 2.1:** Schematics of two triangle structures proposed in this work. (a) Three-site Kitaev triangle with bond-dependent Peierls phases. (b) Hollow triangular island with a uniform vector potential.

## 2.2 Kitaev Triangle

In this section we present an exactly solvable minimal model with three sites forming a “Kitaev triangle” that can host MZM at different pairs of vertices controlled by Peierls phases on the edges. The Bogoliubov-de Gennes (BdG) Hamiltonian includes complex hopping and  $p$ -wave pairing between three spinless fermions forming an equilateral triangle [Fig. 2.1 (a)]:

$$\mathcal{H} = \sum_{\langle jl \rangle} (-te^{i\phi_{jl}} c_j^\dagger c_l + \Delta e^{i\theta_{jl}} c_j c_l + \text{h.c.}) - \sum_j \mu c_j^\dagger c_j, \quad (2.1)$$

where  $t$  is the hopping amplitude,  $\Delta$  is the amplitude of the (2D)  $p$ -wave pairing,  $\mu$  is the chemical potential,  $\theta_{jl}$  is the polar angle of  $\mathbf{r}_{jl} = \mathbf{r}_l - \mathbf{r}_j$  (the  $x$  axis is chosen to be along  $\mathbf{r}_{12}$ ), consistent with  $\{c_l^\dagger, c_j^\dagger\} = 0$ .  $\phi_{jl}$  is the Peierls phase due to a bond-dependent vector potential  $\mathbf{A}$  to be specified below (the nearest neighbor distance  $a$  is chosen to be the length unit hereinbelow):

$$\phi_{jl} = \frac{e}{\hbar} \int_{\mathbf{r}_j}^{\mathbf{r}_l} \mathbf{A} \cdot d\mathbf{l} = -\phi_{lj} \quad (2.2)$$

where  $e > 0$  is the absolute value of the electron charge. Below we use the natural units  $e = \hbar = 1$ . To get the conditions for having MZM in this model we rewrite  $\mathcal{H}$  in the Majorana fermion basis

$$a_j = c_j + c_j^\dagger, b_j = \frac{1}{i}(c_j - c_j^\dagger):$$

$$\begin{aligned} \mathcal{H} = & -\frac{i}{2} \sum_{\langle jl \rangle} \left[ (t \sin \phi_{jl} - \Delta \sin \theta_{jl}) a_j a_l \right. \\ & + (t \sin \phi_{jl} + \Delta \sin \theta_{jl}) b_j b_l \\ & + (t \cos \phi_{jl} - \Delta \cos \theta_{jl}) a_j b_l \\ & \left. - (t \cos \phi_{jl} + \Delta \cos \theta_{jl}) b_j a_l \right] - \frac{i\mu}{2} \sum_j a_j b_j \end{aligned} \quad (2.3)$$

For concreteness we consider the Kitaev limit  $t = \Delta$ ,  $\mu = 0$ , and choose  $\phi_{12} = 0$  so that sites 1 and 2 alone form a minimal Kitaev chain with  $\mathcal{H}_{12} = itb_1a_2$  and hosting MZM  $a_1$  and  $b_2$ . In order for the MZM to persist in the presence of site 3, one can choose  $\phi_{23}$  and  $\phi_{31}$  so that all terms involving these Majorana operators cancel out. For example, consider the 2 – 3 bond, for which  $\theta_{23} = 2\pi/3$ , we require

$$\sin \phi_{23} + \sin \frac{2\pi}{3} = \cos \phi_{23} + \cos \frac{2\pi}{3} = 0 \quad (2.4)$$

which means  $\phi_{23} = -\pi/3$ . Similarly one can find  $\phi_{31} = -\phi_{13} = -\pi/3$ . The three Peierls phases can be realized by the following staggered vector potential

$$\mathbf{A} = [1 - 2\Theta(x)] \frac{2\pi}{3\sqrt{3}} \hat{\mathbf{y}} \quad (2.5)$$

where  $\Theta(x)$  is the Heaviside step function. In fact, using a uniform  $\mathbf{A} = \frac{2\pi}{3\sqrt{3}} \hat{\mathbf{y}}$ , which corresponds to  $\phi_{23} = -\pi/3 = -\phi_{31}$  also works, since the existence of  $a_1$  is unaffected by  $\phi_{23}$ . However, in this case the counterpart of  $b_2$  is not localized on a single site. For the same reason, the above condition for MZM localized at triangle corners can be generalized to Kitaev chains forming a triangular loop, as well as to finite-size triangles of 2D spinless  $p$ -wave superconductors in the Kitaev limit, as the existence of  $a_1$  and  $b_2$  are only dictated by the vector potential near the corresponding corners.


It should be noted that in the latter case, 1D Majorana edge states will arise when the triangle becomes larger, and effectively diminish the gap that protects the corner MZM. On the other hand, for the longer Kitaev chain, due to the potential practical difficulty of controlling further-neighbor hopping and pairing amplitudes, it is better to resort to the approach of controlling the individual topological phases of the three edges which will be detailed in the next section.

We next show that the minimal Kitaev triangle suffices to demonstrate braiding of MZM. To this end we consider a closed parameter path linearly interpolating between the following sets of values of  $\phi_{jl}$ :

$$\begin{aligned}
(\phi_{12}, \phi_{23}, \phi_{31}) &= \left(0, -\frac{\pi}{3}, -\frac{\pi}{3}\right) \equiv \phi_1 \\
&\rightarrow \left(-\frac{\pi}{3}, -\frac{\pi}{3}, 0\right) \equiv \phi_2 \\
&\rightarrow \left(-\frac{\pi}{3}, 0, -\frac{\pi}{3}\right) \equiv \phi_3 \\
&\rightarrow \phi_1
\end{aligned} \tag{2.6}$$

It is straightforward to show that at  $\phi_2$  and  $\phi_3$  there are MZM located at sites 3, 1 and 2, 3, respectively. Therefore the two original MZM at sites 1, 2 should switch their positions at the end of the adiabatic evolution.

Indeed, Fig. 2.2 shows that the MZM stays at zero energy throughout the parameter path that interchanges their positions. To show that such an operation indeed realizes braiding, we explicitly calculated the many-body Berry phase of the evolution [4, 50, 54] and found the two degenerate many-body ground states acquire a  $\frac{\pi}{2}$  difference in their Berry phases as expected [4]. Compared to the minimum T-junction model with four sites [4], our Kitaev triangle model only requires three sites to achieve braiding between two MZM, and is potentially also easier to engineer experimentally. In the next section we will show that a more mesoscopic hollow-triangle structure can achieve similar results and may be preferred in other materials platforms.



**Figure 2.2:** (a) Evolution of the eigenvalues of the 3-site Kitaev triangle along the closed parameter path for  $\phi$  on the three edges. (b) MZM wavefunctions at different points of the parameter path. Clockwise from the upper left panel:  $\phi_1 \rightarrow \frac{1}{2}(\phi_1 + \phi_2) \rightarrow \phi_2 \rightarrow \phi_3$ .

## 2.3 Hollow Triangles

For systems with less fine-tuned Hamiltonians than the minimal model in the previous section, it is more instructive to search for MZM based on topological arguments. In this section we show that MZM generally appear at the corners of a hollow triangle, which can be approximated by joining three finite-width chains or ribbons whose bulk topology is individually tuned by the same uniform vector potential.



To this end, we first show that topological phase transitions can be induced by a vector potential in a spinless  $p$ -wave superconductor ribbon. In comparison with similar previous proposals that mostly focused on vector potentials or supercurrents flowing along the chain [52, 53], we consider in particular the tunability by varying the direction of the vector potential relative to the length direction of the ribbon, which will become instrumental in a triangular structure.

Consider Eq. (2.1) on a triangular lattice defined by unit-length lattice vectors  $(\mathbf{a}_1, \mathbf{a}_2) = (\hat{\mathbf{x}}, \frac{1}{2}\hat{\mathbf{x}} + \frac{\sqrt{3}}{2}\hat{\mathbf{y}})$  with  $W$  unit cells along  $\mathbf{a}_2$  but infinite unit cells along  $\mathbf{a}_1$ , and assume the Peierls phases are due to a uniform vector potential  $\mathbf{A}$  so that  $\phi_{jl} = \mathbf{A} \cdot \mathbf{r}_{jl}$ . We also introduce  $\mathbf{a}_3 \equiv -\mathbf{a}_1 + \mathbf{a}_2$  for later convenience. The Hamiltonian is periodic along  $x$  and can be Fourier transformed through  $c_{m,n}^\dagger = \frac{1}{\sqrt{N}} \sum_k c_{k,n}^\dagger e^{-ikm}$ , where  $m, n$  label the lattice sites as  $\mathbf{r}_{m,n} = m\mathbf{a}_1 + n\mathbf{a}_2$ . The resulting momentum space Hamiltonian can be written as the following block form up to a constant

$$\begin{aligned} \mathcal{H} &= \frac{1}{2} \sum_k \Psi_k^\dagger \begin{pmatrix} h_t(k) & h_\Delta(k) \\ h_\Delta^\dagger(k) & -h_t^*(-k) \end{pmatrix} \Psi_k \\ &\equiv \frac{1}{2} \sum_k \Psi_k^\dagger H(k) \Psi_k \end{aligned} \quad (2.7)$$

where  $\Psi_k \equiv (c_{k,1}, \dots, c_{k,W}, c_{-k,1}^\dagger, \dots, c_{-k,W}^\dagger)^T$ .  $h_t(k)$  is a  $W \times W$  Hermitian tridiagonal matrix with  $(h_t)_{n,n} = -2t \cos(k + \mathbf{A} \cdot \mathbf{a}_1) - \mu$  and  $(h_t)_{n,n+1} = -t (e^{i(-k + \mathbf{A} \cdot \mathbf{a}_3)} + e^{i\mathbf{A} \cdot \mathbf{a}_2})$ .  $h_\Delta(k)$  is a  $W \times W$  tridiagonal matrix with  $(h_\Delta)_{n,n} = -2i\Delta \sin k$  and  $(h_\Delta)_{n,n\pm 1} = \mp \Delta [e^{-i(\pm k + \frac{2\pi}{3})} + e^{-i\frac{\pi}{3}}]$ .


By transforming Eq. (2.7) to the Majorana basis using the unitary transformation:

$$U \equiv \frac{1}{\sqrt{2}} \begin{pmatrix} 1 & 1 \\ -i & i \end{pmatrix} \otimes I \quad (2.8)$$

where  $I$  is a  $W \times W$  identity matrix, and defining  $A_k \equiv -iUH(k)U^\dagger$ , not to be confused with the vector potential, one can calculate the Majorana number [7]  $\mathcal{M}$  of the 1D ribbon as [55]

$$\mathcal{M} = \text{sgn} [\text{Pf}(A_{k=0})\text{Pf}(A_{k=\pi})] \quad (2.9)$$

where Pf stands for the Pfaffian of a skew-symmetric matrix [7]. When  $\mathcal{M} = -1$ , the 1D system is in a nontrivial topological phase with MZM appearing at open ends of semi-infinite ribbons, and otherwise for  $\mathcal{M} = 1$ .




**Figure 2.3:** (a) Topological phase diagram for a  $W = 1$  triangular chain with the Hamiltonian Eq. (2.7) obtained by superimposing the  $\mathcal{M}(A, \mu)$  plots of 1D chains with  $\mathbf{A} = A\hat{y}$  and  $\mathbf{A} = A(\frac{\sqrt{3}}{2}\hat{x} + \frac{1}{2}\hat{y})$ . Color scheme: white— $\mathcal{M} = 1$ , dark blue— $\mathcal{M} = -1$ , light blue— $\mathcal{M} = 0$  (b) Near-gap BdG eigen-energies vs  $A$  for a finite triangle with edge length  $L = 50$ ,  $W = 1$ , and  $\mu = 1.6$ .

In Fig. 2.3 (a) we show the topological phase diagrams for a 1D ribbon with width  $W = 1$ ,  $\mathbf{A} = A\hat{\mathbf{y}}$  and  $\mathbf{A} = A(\frac{\sqrt{3}}{2}\hat{\mathbf{x}} + \frac{1}{2}\hat{\mathbf{y}})$  superimposed (see below). We found that the vector potential component normal to the ribbon length direction has no effect on the Majorana number, nor does the sign of its component along the ribbon length direction. However, topological phase transitions can be induced by varying the size of the vector potential component along the ribbon, consistent with previous results [52, 53]. These properties motivate us to consider the structure of a hollow triangle formed by three finite-width ribbons subject to a uniform vector potential  $\mathbf{A} = A\hat{\mathbf{y}}$  as illustrated in Fig. 2.1 (b). The light blue color on the phase diagram Fig. 2.3 (a) therefore means that the bottom edge and the two upper edges of the hollow triangle have different  $\mathcal{M}$ , which should give rise to MZM localized at the two bottom corners if the triangle is large enough so that bulk-edge correspondence holds, and gap closing does not occur at other places along its edges.

To show that corner MZM indeed appear when the conditions given by the phase diagram Fig. 2.3 (a) are met, we directly diagonalize the BdG Hamiltonian of a finite hollow triangle with edge length  $L = 50$  and width  $W = 1$ . Fig. 2.3 (b) shows the spectral flow (BdG eigen-energies evolving with increasing vector potential  $A$ ) close to zero energy at chemical potential  $\mu = 1.6$ . Indeed, zero-energy modes appear in the regions of  $\mu$  and  $A$  consistent with the phase diagram (except when the bulk band gap is too small; see [54] for some examples.). Hollow triangles with larger  $W$  also have qualitatively similar behavior, although the phase diagrams are more complex [54]. The eigenfunctions for the zero-energy modes at  $A = 2.75$  and  $\mu = 1.6$  in Fig. 2.4 (b) also confirm their spatial localization at the bottom corners of the triangle.


We finally show that rotating the uniform vector potential in-plane can manipulate the positions of the MZM without hybridizing them with bulk states for certain ranges of  $\mu$  and  $A$ . Fig. 2.4 (a) plots the spectral flow versus the in-plane azimuthal angle of  $\mathbf{A}$ , which clearly shows that the zero-energy modes persist throughout the rotation and the bulk gap never closes. Figs. 2.4 (b-d) plot the BdG wavefunctions of the MZM at special values of  $\varphi$ . One can see that the two MZM appear to cycle through the three vertices by following the rotation of  $\mathbf{A}$ . The robustness of the MZM therefore requires the condition of two edges being in a different topological phase from



**Figure 2.4:** (a) Spectral flow of a hollow triangle with  $W = 1$ ,  $L = 50$ ,  $\mu = 1.6$ , and  $A = 2.75$  with increasing rotation angle  $\varphi$ , defined through  $\mathbf{A} = A(-\sin \varphi \hat{x} + \cos \varphi \hat{y})$ . (b-d) BdG eigenfunction  $|\Psi|^2$  summed over the two zero modes at  $\varphi = 0$ ,  $\frac{\pi}{6}$ , and  $\frac{\pi}{3}$ , respectively.

the third one to be satisfied throughout the rotation. Such a criterion combined with the individual phase diagrams of the edges can help isolate the desired parameter regions of  $\mu$  and  $A$ . We also note that the positions of the MZM do not interchange after  $\varphi$  increases from 0 to  $\pi$ , different from the situation of the minimal Kitaev triangle in Fig. 2.2. The reason is that the MZM in the latter case are not due to bulk-boundary correspondence [the values of  $A = \frac{2\pi}{3\sqrt{3}}$  and  $\mu = 0$  are a critical point in the phase diagram Fig. 2.3 (a)]. While the positions of the MZM at special points along the parameter path in the hollow triangle case have to be additionally constrained by the bulk topological phases of the three edges, that for the Kitaev triangle have more flexibility and are also protected by the finite size of the system.


A model that is closer to a realistic hollow triangular island is the finite-width triangular chain or ribbon. An example, illustrated in Figure 2.5 (c), has its edge length  $L = 50$  and width  $W = 3$ . The phase diagram Fig. 2.5 (a) is created in a similar way as that in Fig. 2.3 (a), assuming a



**Figure 2.5:** (a) Topological phase diagram for a  $W = 3$  hollow triangle obtained by overlapping the  $\mathcal{M}(A, \mu)$  plots of 1D chains with  $\mathbf{A} = A\hat{y}$  and  $\mathbf{A} = A(\frac{\sqrt{3}}{2}\hat{x} + \frac{1}{2}\hat{y})$ . Color scheme: white— $\mathcal{M} = 1$ , dark blue— $\mathcal{M} = -1$ , light blue— $\mathcal{M} = 0$  (b) Near-gap BdG eigen-energies vs  $A$  for a finite triangle with edge length  $L = 50$ ,  $W = 3$ , and  $\mu = 1.6$ . (c) BdG eigenfunction  $|\Psi|^2$  summed over the two zero modes at  $A = 2.4709$ .

constant vector potential and infinitely long  $W = 3$  ribbons. The spectral flow for the actual triangle with  $\mu = 1.6$  in Fig. 2.5 (b) shows MZM in the parameter regions in agreement with the phase diagram. Fig. 2.5 (c) plots the MZM wavefunction for  $A = 2.7409$  and  $\mu = 1.6$  that are indeed well localized at the bottom corners.


We next rotate the uniform vector potential to examine how the MZM move on a hollow triangle. Figure 2.6 shows the spectral flow and eigenfunctions as we rotate  $\varphi = 0$  to  $\varphi = \pi$  counterclockwisely. The two MZM cycle through the three vertices in a similar manner as that in



**Figure 2.6:** (a) Spectral flow of a hollow triangle with  $W = 3$ ,  $L = 50$ ,  $\mu = 1.6$ , and  $A = 2.75$  with increasing rotation angle  $\varphi$ , defined through  $\mathbf{A} = A(-\sin \varphi \hat{x} + \cos \varphi \hat{y})$ . (b-c) BdG eigenfunction  $|\Psi|^2$  summed over the two zero modes at  $\varphi = 0$  and  $\frac{\pi}{3}$ , respectively.

Fig. 4 of the main text (only the MZM wavefunctions at  $\varphi = 0$  and  $\frac{\pi}{3}$  are plotted as representatives of the  $\varphi = n\pi/3$  cases). Note that the spectral flow has 3-fold rotation symmetry but not 6-fold, since increasing  $\varphi$  by  $\frac{2\pi}{3}$  is equivalent to rotating the coordinate system clockwise by  $\frac{2\pi}{3}$ . In contrast, rotating the vector potential by  $\frac{\pi}{3}$ , if without an additional sign change of the  $p$ -wave pairing potential, is not an exact symmetry of the finite triangle. Also we did not try to scrutinize the phase diagram to find a parameter path in which the bulk gap does not close, as in the  $W = 1$  case in the main text. Here we just point out that identifying a system-specific parameter path for adiabatic manipulation of MZM is in principle always possible, especially if one is allowed to have more knobs other than  $\varphi$  in real structures, such as tuning the chemical potential of individual edges or the size of the vector potential, etc.

## 2.4 Braiding MZM in a small network of triangles



**Figure 2.7:** (a) Spectral flow for the critical step of swapping  $\gamma_2$  and  $\gamma_3$  in the example of Fig. 5 in the main text, calculated using four corner-sharing triangles of  $W = 1$  and  $L = 50$ , with  $\mu = 1.6$  and  $A = 2.6$ . Vector potential for the middle triangle in the bottom row can rotate according to  $\mathbf{A} = A(-\sin \varphi \hat{x} + \cos \varphi \hat{y})$  from  $\varphi = \frac{\pi}{6}$  to  $\frac{\pi}{3}$ , while the other three have fixed  $\varphi = 0$ . (b)-(g) BdG eigenfunction  $|\Psi|^2$  summed over the four zero modes at equally-spaced points along the rotation path. The black arrow indicates the direction of the vector potential for the bottom middle triangle.

In this section we show that one can braid two out of four MZM, a minimal setting for nontrivial manipulation of the degenerate many-body ground states, by using a small network of corner-sharing triangles. We focus on the critical step of swapping  $\gamma_2$  and  $\gamma_3$  as labeled in Fig. 5 of the main text. This can be done by rotating the vector potential of the triangle in the middle of the bottom row from  $\varphi = \frac{\pi}{6}$  to  $\frac{\pi}{3}$ . More specifically, when  $\varphi = \frac{\pi}{6}$ , with the chosen values of  $\mu$  and

A, only the right edge of the said triangle is topologically nontrivial. The chain that hosts  $\gamma_{3,4}$  thus extends through this nontrivial edge to the top triangle as in Fig. 2.7 (b). On the other hand, when  $\varphi$  increases to  $\frac{\pi}{3}$ , the nontrivial edge of the middle triangle changes from right to left, which leads to  $\gamma_2$  hopping from its left corner to the right through the top corner, while  $\gamma_3$  is unaffected [Figs. 2.7 (c-g)]. As a result the  $\gamma_2, \gamma_3$  swapping is done without closing the bulk gap, as can be seen from the spectral flow in Fig. 2.7 (a).

## 2.5 Discussion


The hollow interior of the triangles considered in this work is needed for two reasons: (1)  $W \ll L$  is required for bulk-edge correspondence based on 1D topology to hold; (2) A finite  $W$  is needed to gap out the chiral edge states of a 2D spinless  $p$ -wave superconductor based on which Eq. (2.7) is written. The latter is not essential if one does not start with a spinless  $p$ -wave superconductor but a more realistic model such as the Rashba+Zeeman+ $s$ -wave pairing model. On the other hand, the former constraint may also be removed if one uses the Kitaev triangle. Nonetheless, an effective 3-site Kitaev triangle may emerge as the effective theory of triangular structures if a three-orbital low-energy Wannier basis can be isolated, similar to the continuum theory of moiré structures. We also note in passing that the corner MZM in our triangles appear due to different reasons from that in higher-order topological superconductors [41, 56].

For possible physical realizations of our triangles, immediate choices are quantum dots forming a Kitaev triangle [47], planar Josephson junctions or cuts on quantum anomalous Hall insulator/superconductor heterostructures [24] that form a hollow triangle, and triangular atomic chains assembled by an STM tip [17] on a close-packed surface. The quantum-dot platform may be advantageous in the convenience of implementing parity readout by turning the third vertex temporarily into a normal quantum dot [57–59]. Looking into the future, it is more intriguing to utilize the spontaneously formed triangular islands in epitaxial growth [51] with the center region removed either physically by lithography/ablation, or electrically by gating. To create a staggered vector potential or supercurrent profile for the Kitaev triangle, one can use a uniform magnetic



field, corresponding to a constant vector potential gradient, plus a uniform supercurrent that controls the position of the zero. It is also possible to use two parallel superconducting wires with counter-propagating supercurrents proximate to the triangle.

A tentative design for braiding more than two MZM, illustrated in Fig. 2.8, consists of four triangles sharing corners with their neighbors. The critical step of transporting  $\gamma_2$  to the left vertex of the rightmost triangle, corresponding to Figs. 2.8 (b,c), can be achieved by rotating the vector potential of the bottom-middle triangle counterclockwisely from  $\varphi = \frac{\pi}{6}$  to  $\frac{\pi}{3}$ , which swaps the topological phases of the two side edges as shown in Fig. 2.4. In [54] we show this operation



**Figure 2.8:** Representative steps for braiding four MZM in four triangles sharing corners. (a) Initialization of four MZM  $\gamma_1, \gamma_2, \gamma_3, \gamma_4$ . All three edges of the bottom-middle and the top triangles are in the trivial phase by e.g. controlling the chemical potential. The bottom-left and bottom-right triangles have  $\varphi = 0$  so that their bottom edges are nontrivial. (b) Moving  $\gamma_3$  by “switching on” the middle triangle by changing the chemical potential under a fixed vector potential at  $\varphi = \frac{\pi}{6}$ , and then turning on the top triangle with similar means except  $\varphi = 0$ . (c) Transporting  $\gamma_2$  to the right triangle through rotating the vector potential in the middle triangle counterclockwise by  $\pi/6$ . (d) Moving  $\gamma_3$  to the left triangle by “switching off” the top triangle followed by the middle triangle.

does not involve gap closing at least for certain parameter regions. Our work provides a versatile platform for manipulating MZM based on currently available candidate MZM systems and for potentially demonstrating the non-Abelian nature of MZM in near-term devices.

# Chapter 3

## Floquet Landau Levels

1. Introduction (Tahir's intro is fine, maybe in my own words, Floquet engineering)
  - i Time dependent, motivation—QAHE gap but not QHE gap
  - ii Floquet Theorem— quasi-energy spectrum
2. Results
  - i square +  $A(t)$  (Tahir's perturbative calc)
  - ii honeycomb +  $A(t)$  (Tahir's perturbative calc)
3. Discussion and future

# Chapter 4

## Conclusion and Discussion

What makes gauge potential unique in creating/tuning/manipulating new topoglical systems

Applications

# Appendix A

## Suitable Name

1. [Majorana Number derivation](#)
2. [Other derivations not included in introduction](#)

# Bibliography

- [1] D. A. Ivanov. Non-Abelian Statistics of Half-Quantum Vortices in p -Wave Superconductors. *Phys. Rev. Lett.*, 86(2):268–271, January 2001.
- [2] A. Yu. Kitaev. Fault-tolerant quantum computation by anyons. *Annals of Physics*, 303(1):2–30, January 2003.
- [3] Chetan Nayak, Steven H. Simon, Ady Stern, Michael Freedman, and Sankar Das Sarma. Non-Abelian anyons and topological quantum computation. *Rev. Mod. Phys.*, 80(3):1083–1159, September 2008.
- [4] Jason Alicea, Yuval Oreg, Gil Refael, Felix von Oppen, and Matthew P. A. Fisher. Non-Abelian statistics and topological quantum information processing in 1D wire networks. *Nature Phys*, 7(5):412–417, May 2011.
- [5] David Aasen, Michael Hell, Ryan V. Mishmash, Andrew Higginbotham, Jeroen Danon, Martin Leijnse, Thomas S. Jespersen, Joshua A. Folk, Charles M. Marcus, Karsten Flensberg, and Jason Alicea. Milestones Toward Majorana-Based Quantum Computing. *Phys. Rev. X*, 6(3):031016, August 2016.
- [6] N. Read and Dmitry Green. Paired states of fermions in two dimensions with breaking of parity and time-reversal symmetries and the fractional quantum Hall effect. *Phys. Rev. B*, 61(15):10267–10297, April 2000.
- [7] A. Yu Kitaev. Unpaired Majorana fermions in quantum wires. *Phys.-Usp.*, 44(10S):131, October 2001.
- [8] Jean-Pascal Brison. P-Wave Superconductivity and d-Vector Representation. In Hervé Bulou, Loïc Joly, Jean-Michel Mariot, and Fabrice Scheurer, editors, *Magnetism and Accelerator-Based Light Sources*, Springer Proceedings in Physics, pages 165–204, Cham, 2021. Springer International Publishing.

- [9] V. Mourik, K. Zuo, S. M. Frolov, S. R. Plissard, E. P. A. M. Bakkers, and L. P. Kouwenhoven. Signatures of Majorana Fermions in Hybrid Superconductor-Semiconductor Nanowire Devices. *Science*, 336(6084):1003–1007, May 2012.
- [10] Leonid P. Rokhinson, Xinyu Liu, and Jacek K. Furdyna. The fractional a.c. Josephson effect in a semiconductor–superconductor nanowire as a signature of Majorana particles. *Nature Phys*, 8(11):795–799, November 2012.
- [11] M. T. Deng, C. L. Yu, G. Y. Huang, M. Larsson, P. Caroff, and H. Q. Xu. Anomalous Zero-Bias Conductance Peak in a Nb–InSb Nanowire–Nb Hybrid Device. *Nano Lett.*, 12(12):6414–6419, December 2012.
- [12] T.-P. Choy, J. M. Edge, A. R. Akhmerov, and C. W. J. Beenakker. Majorana fermions emerging from magnetic nanoparticles on a superconductor without spin-orbit coupling. *Phys. Rev. B*, 84(19):195442, November 2011.
- [13] Bernd Braunecker and Pascal Simon. Interplay between Classical Magnetic Moments and Superconductivity in Quantum One-Dimensional Conductors: Toward a Self-Sustained Topological Majorana Phase. *Phys. Rev. Lett.*, 111(14):147202, October 2013.
- [14] Jelena Klinovaja, Peter Stano, Ali Yazdani, and Daniel Loss. Topological Superconductivity and Majorana Fermions in RKKY Systems. *Phys. Rev. Lett.*, 111(18):186805, November 2013.
- [15] S. Nadj-Perge, I. K. Drozdov, B. A. Bernevig, and Ali Yazdani. Proposal for realizing Majorana fermions in chains of magnetic atoms on a superconductor. *Phys. Rev. B*, 88(2):020407, July 2013.
- [16] Stevan Nadj-Perge, Ilya K. Drozdov, Jian Li, Hua Chen, Sangjun Jeon, Jungpil Seo, Allan H. MacDonald, B. Andrei Bernevig, and Ali Yazdani. Observation of Majorana fermions in ferromagnetic atomic chains on a superconductor. *Science*, 346(6209):602–607, October 2014.

- [17] Lucas Schneider, Philip Beck, Jannis Neuhaus-Steinmetz, Levente Rózsa, Thore Posske, Jens Wiebe, and Roland Wiesendanger. Precursors of Majorana modes and their length-dependent energy oscillations probed at both ends of atomic Shiba chains. *Nat. Nanotechnol.*, 17(4):384–389, April 2022.
- [18] Liang Fu and C. L. Kane. Superconducting proximity effect and Majorana fermions at the surface of a topological insulator. *Phys. Rev. Lett.*, 100(9):096407, March 2008.
- [19] Pavan Hosur, Pouyan Ghaemi, Roger S. K. Mong, and Ashvin Vishwanath. Majorana Modes at the Ends of Superconductor Vortices in Doped Topological Insulators. *Phys. Rev. Lett.*, 107(9):097001, August 2011.
- [20] Andrew C. Potter and Patrick A. Lee. Engineering a  $\mathbb{Z}_2$  superconductor: Comparison of topological insulator and Rashba spin-orbit-coupled materials. *Phys. Rev. B*, 83(18):184520, May 2011.
- [21] M. Veldhorst, M. Snelder, M. Hoek, C. G. Molenaar, D. P. Leusink, A. A. Golubov, H. Hilgenkamp, and A. Brinkman. Magnetotransport and induced superconductivity in Bi based three-dimensional topological insulators. *physica status solidi (RRL) – Rapid Research Letters*, 7(1-2):26–38, 2013.
- [22] Chui-Zhen Chen, Ying-Ming Xie, Jie Liu, Patrick A. Lee, and K. T. Law. Quasi-one-dimensional quantum anomalous Hall systems as new platforms for scalable topological quantum computation. *Phys. Rev. B*, 97(10):104504, March 2018.
- [23] Yongxin Zeng, Chao Lei, Gaurav Chaudhary, and Allan H. MacDonald. Quantum anomalous Hall Majorana platform. *Phys. Rev. B*, 97(8):081102, February 2018.
- [24] Ying-Ming Xie, Xue-Jian Gao, Tai-Kai Ng, and K. T. Law. Creating Localized Majorana Zero Modes in Quantum Anomalous Hall Insulator/Superconductor Heterostructures with a Scissor, June 2021.



- [25] Yuval Oreg, Gil Refael, and Felix von Oppen. Helical Liquids and Majorana Bound States in Quantum Wires. *Phys. Rev. Lett.*, 105(17):177002, October 2010.
- [26] Jay D. Sau, Roman M. Lutchyn, Sumanta Tewari, and S. Das Sarma. Generic New Platform for Topological Quantum Computation Using Semiconductor Heterostructures. *Phys. Rev. Lett.*, 104(4):040502, January 2010.
- [27] Roman M. Lutchyn, Tudor D. Stanescu, and S. Das Sarma. Search for Majorana Fermions in Multiband Semiconducting Nanowires. *Phys. Rev. Lett.*, 106(12):127001, March 2011.
- [28] Andrew C. Potter and Patrick A. Lee. Topological superconductivity and Majorana fermions in metallic surface states. *Phys. Rev. B*, 85(9):094516, March 2012.
- [29] Jian Li, Titus Neupert, Zhijun Wang, A. H. MacDonald, A. Yazdani, and B. Andrei Bernevig. Two-dimensional chiral topological superconductivity in Shiba lattices. *Nat Commun*, 7(1):12297, July 2016.
- [30] Chao Lei, Hua Chen, and Allan H. MacDonald. Ultrathin Films of Superconducting Metals as a Platform for Topological Superconductivity. *Phys. Rev. Lett.*, 121(22):227701, November 2018.
- [31] Annica M. Black-Schaffer and Jacob Linder. Majorana fermions in spin-orbit-coupled ferromagnetic Josephson junctions. *Phys. Rev. B*, 84(18):180509, November 2011.
- [32] Falko Pientka, Alessandro Romito, Mathias Duckheim, Yuval Oreg, and Felix von Oppen. Signatures of topological phase transitions in mesoscopic superconducting rings. *New J. Phys.*, 15(2):025001, February 2013.
- [33] Michael Hell, Martin Leijnse, and Karsten Flensberg. Two-Dimensional Platform for Networks of Majorana Bound States. *Phys. Rev. Lett.*, 118(10):107701, March 2017.
- [34] Antonio Fornieri, Alexander M. Whiticar, F. Setiawan, Elías Portolés, Asbjørn C. C. Drachmann, Anna Keselman, Sergei Gronin, Candice Thomas, Tian Wang, Ray Kallagher, Geof-

- frey C. Gardner, Erez Berg, Michael J. Manfra, Ady Stern, Charles M. Marcus, and Fabrizio Nichele. Evidence of topological superconductivity in planar Josephson junctions. *Nature*, 569(7754):89–92, May 2019.
- [35] Hechen Ren, Falko Pientka, Sean Hart, Andrew T. Pierce, Michael Kosowsky, Lukas Lunczer, Raimund Schlereth, Benedikt Scharf, Ewelina M. Hankiewicz, Laurens W. Molenkamp, Bertrand I. Halperin, and Amir Yacoby. Topological superconductivity in a phase-controlled Josephson junction. *Nature*, 569(7754):93–98, May 2019.
- [36] Benedikt Scharf, Falko Pientka, Hechen Ren, Amir Yacoby, and Ewelina M. Hankiewicz. Tuning topological superconductivity in phase-controlled Josephson junctions with Rashba and Dresselhaus spin-orbit coupling. *Phys. Rev. B*, 99(21):214503, June 2019.
- [37] Tong Zhou, Matthieu C. Dartailh, William Mayer, Jong E. Han, Alex Matos-Abiague, Javad Shabani, and Igor Žutić. Phase Control of Majorana Bound States in a Topological  $\mathbb{X}$  Junction. *Phys. Rev. Lett.*, 124(13):137001, April 2020.
- [38] Jin-Peng Xu, Mei-Xiao Wang, Zhi Long Liu, Jian-Feng Ge, Xiaojun Yang, Canhua Liu, Zhu An Xu, Dandan Guan, Chun Lei Gao, Dong Qian, Ying Liu, Qiang-Hua Wang, Fu-Chun Zhang, Qi-Kun Xue, and Jin-Feng Jia. Experimental Detection of a Majorana Mode in the core of a Magnetic Vortex inside a Topological Insulator-Superconductor  $\text{Bi}_2\text{Te}_3/\text{NbSe}_2$  Heterostructure. *Phys. Rev. Lett.*, 114(1):017001, January 2015.
- [39] S. M. Albrecht, A. P. Higginbotham, M. Madsen, F. Kuemmeth, T. S. Jespersen, J. Nygård, P. Krogstrup, and C. M. Marcus. Exponential protection of zero modes in Majorana islands. *Nature*, 531(7593):206–209, March 2016.
- [40] Hao-Hua Sun, Kai-Wen Zhang, Lun-Hui Hu, Chuang Li, Guan-Yong Wang, Hai-Yang Ma, Zhu-An Xu, Chun-Lei Gao, Dan-Dan Guan, Yao-Yi Li, Canhua Liu, Dong Qian, Yi Zhou, Liang Fu, Shao-Chun Li, Fu-Chun Zhang, and Jin-Feng Jia. Majorana Zero Mode Detected

- with Spin Selective Andreev Reflection in the Vortex of a Topological Superconductor. *Phys. Rev. Lett.*, 116(25):257003, June 2016.
- [41] Dongfei Wang, Lingyuan Kong, Peng Fan, Hui Chen, Shiyu Zhu, Wenyao Liu, Lu Cao, Yujie Sun, Shixuan Du, John Schneeloch, Ruidan Zhong, Genda Gu, Liang Fu, Hong Ding, and Hong-Jun Gao. Evidence for Majorana bound states in an iron-based superconductor. *Science*, 362(6412):333–335, October 2018.
- [42] Berthold Jäck, Yonglong Xie, Jian Li, Sangjun Jeon, B. Andrei Bernevig, and Ali Yazdani. Observation of a Majorana zero mode in a topologically protected edge channel. *Science*, 364(6447):1255–1259, June 2019.
- [43] Sujit Manna, Peng Wei, Yingming Xie, Kam Tuen Law, Patrick A. Lee, and Jagadeesh S. Moodera. Signature of a pair of Majorana zero modes in superconducting gold surface states. *Proceedings of the National Academy of Sciences*, 117(16):8775–8782, April 2020.
- [44] Geoffrey L. Fatin, Alex Matos-Abiague, Benedikt Scharf, and Igor Žutić. Wireless Majorana Bound States: From Magnetic Tunability to Braiding. *Phys. Rev. Lett.*, 117(7):077002, August 2016.
- [45] Jay D. Sau and S. Das Sarma. Realizing a robust practical Majorana chain in a quantum-dot-superconductor linear array. *Nat Commun*, 3(1):964, July 2012.
- [46] Martin Leijnse and Karsten Flensberg. Parity qubits and poor man’s Majorana bound states in double quantum dots. *Phys. Rev. B*, 86(13):134528, October 2012.
- [47] Tom Dvir, Guanzhong Wang, Nick van Loo, Chun-Xiao Liu, Grzegorz P. Mazur, Alberto Bordin, Sebastiaan L. D. ten Haaf, Ji-Yin Wang, David van Driel, Francesco Zatelli, Xiang Li, Filip K. Malinowski, Sasa Gazibegovic, Ghada Badawy, Erik P. A. M. Bakkers, Michael Wimmer, and Leo P. Kouwenhoven. Realization of a minimal Kitaev chain in coupled quantum dots. *Nature*, 614(7948):445–450, February 2023.

- [48] Torsten Karzig, Christina Knapp, Roman M. Lutchyn, Parsa Bonderson, Matthew B. Hastings, Chetan Nayak, Jason Alicea, Karsten Flensberg, Stephan Plugge, Yuval Oreg, Charles M. Marcus, and Michael H. Freedman. Scalable designs for quasiparticle-poisoning-protected topological quantum computation with Majorana zero modes. *Phys. Rev. B*, 95(23):235305, June 2017.
- [49] Andrew C. Potter and Patrick A. Lee. Multichannel Generalization of Kitaev’s Majorana End States and a Practical Route to Realize Them in Thin Films. *Phys. Rev. Lett.*, 105(22):227003, November 2010.
- [50] Jian Li, Titus Neupert, B. Andrei Bernevig, and Ali Yazdani. Manipulating Majorana zero modes on atomic rings with an external magnetic field. *Nat Commun*, 7(1):10395, January 2016.
- [51] O. Pietzsch, S. Okatov, A. Kubetzka, M. Bode, S. Heinze, A. Lichtenstein, and R. Wiesendanger. Spin-Resolved Electronic Structure of Nanoscale Cobalt Islands on Cu(111). *Phys. Rev. Lett.*, 96(23):237203, June 2006.
- [52] Alessandro Romito, Jason Alicea, Gil Refael, and Felix von Oppen. Manipulating Majorana fermions using supercurrents. *Phys. Rev. B*, 85(2):020502, January 2012.
- [53] Kazuaki Takasan, Shuntaro Sumita, and Youichi Yanase. Supercurrent-induced topological phase transitions. *Phys. Rev. B*, 106(1):014508, July 2022.
- [54] *See Supplemental Material at [url].*
- [55] Jian Li, Hua Chen, Ilya K. Drozdov, A. Yazdani, B. Andrei Bernevig, and A. H. MacDonald. Topological superconductivity induced by ferromagnetic metal chains. *Phys. Rev. B*, 90(23):235433, December 2014.
- [56] Tudor E. Pahomi, Manfred Sigrist, and Alexey A. Soluyanov. Braiding Majorana corner modes in a second-order topological superconductor. *Phys. Rev. Res.*, 2(3):032068, September 2020.

- [57] Ryan V. Mishmash, Bela Bauer, Felix von Oppen, and Jason Alicea. Dephasing and leakage dynamics of noisy Majorana-based qubits: Topological versus Andreev. *Phys. Rev. B*, 101(7):075404, February 2020.
- [58] Gábor Széchenyi and András Pályi. Parity-to-charge conversion for readout of topological majorana qubits. *Phys. Rev. B*, 101:235441, Jun 2020.
- [59] Guan-Hao Feng and Hong-Hao Zhang. Probing robust Majorana signatures by crossed Andreev reflection with a quantum dot. *Phys. Rev. B*, 105(3):035148, January 2022.