



Superconducting Triangular Islands as a Platform for Manipulating Majorana Zero Modes

Aidan Winblad
Hua Chen

Department of Physics
Colorado State University

March 8, 2024

Motivation

Kitaev
Triangle

Hollow
Triangle

Braiding

Summary



Motivation

Superconducting
triangular islands
as a platform for
manipulating
Majorana zero
modes

Aidan Winblad

Motivation


Kitaev
Triangle

Hollow
Triangle

Braiding

Summary


- P-wave superconductors contain half-quantum vortices.
 - Majorana fermions located at core of a vortex.
 - Braiding vortices exhibits Non-Abelian statistics.
- 1D p-wave superconductors host Majorana fermions on end points.
 - Measurements in real systems:
 - V. Mourik, *Science* **336**, 1003 (2012)
 - S. Nadj-Perge, *Science* **346**, 602 (2014)
 - L. Schneider, *Nat. Nanotechnol.* **17**, 384 (2022)
- Quasi-1D T-junction
 - Braiding of Majorana fermions is defined for 2D.
 - In practice challenging to make, but still feasible and seriously pursued.



Alicea, *Nature Phys.* **7**, 412 (2011)



- Consider triangular islands, topologically similar to T-junctions.
- Islands of three-fold rotational symmetry occur naturally in epitaxial growth on close-packed metal surfaces.
- Good platform for transition from 2D to 1D topological superconductor.




Triangular Co islands on Cu(111).
Pietzsch et al., *PRL* **96**, 237203 (2006)



Kitaev Triangle

$$\mathcal{H} = \sum_{\langle j,l \rangle} \left[-te^{i\phi_{jl}} c_j^\dagger c_l + \Delta e^{i\theta_{jl}} c_j c_l + h.c. \right] - \sum_j \mu c_j^\dagger c_j$$



$$\begin{aligned} (\phi_{12}, \phi_{23}, \phi_{31}) &= \left(0, -\frac{\pi}{3}, -\frac{\pi}{3}\right) = \phi_1 \\ &\rightarrow \left(-\frac{\pi}{3}, -\frac{\pi}{3}, 0\right) = \phi_2 \\ &\rightarrow \left(-\frac{\pi}{3}, 0, -\frac{\pi}{3}\right) = \phi_3 \\ &\rightarrow \phi_1 \end{aligned}$$



Kitaev Triangle

Aidan Winblad

Motivation


Kitaev
Triangle

Hollow
Triangle



Braiding

Summary

$$\mathcal{H} = \sum_{\langle j,l \rangle} \left[-te^{i\phi_{jl}} c_j^\dagger c_l + \Delta e^{i\theta_{jl}} c_j c_l + h.c. \right] - \sum_j \mu c_j^\dagger c_j$$



$$\begin{aligned} (\phi_{12}, \phi_{23}, \phi_{31}) &= \left(0, -\frac{\pi}{3}, -\frac{\pi}{3}\right) = \phi_1 \\ &\rightarrow \left(-\frac{\pi}{3}, -\frac{\pi}{3}, 0\right) = \phi_2 \\ &\rightarrow \left(-\frac{\pi}{3}, 0, -\frac{\pi}{3}\right) = \phi_3 \\ &\rightarrow \phi_1 \end{aligned}$$





Triangular Ribbon and Topological Phases

Superconducting
triangular islands
as a platform for
manipulating
Majorana zero
modes

Aidan Winblad

$$\phi_{jl} = \frac{e}{\hbar} \int_{\mathbf{r}_j}^{\mathbf{r}_l} \mathbf{A} \cdot d\mathbf{l} = -\phi_{lj}$$



Motivation

Kitaev
Triangle

Hollow
Triangle

Braiding

Summary





Triangular Ribbon and Topological Phases

Superconducting
triangular islands
as a platform for
manipulating
Majorana zero
modes

Aidan Winblad

Motivation




Kitaev
Triangle

Hollow
Triangle

Braiding

Summary

$$\phi_{jl} = \frac{e}{\hbar} \int_{\mathbf{r}_j}^{\mathbf{r}_l} \mathbf{A} \cdot d\mathbf{l} = -\phi_{lj}$$





Triangular Ribbon and Topological Phases

Superconducting
triangular islands
as a platform for
manipulating
Majorana zero
modes

Aidan Winblad

Motivation




Kitaev
Triangle

Hollow
Triangle

Braiding

Summary

$$\phi_{jl} = \frac{e}{\hbar} \int_{\mathbf{r}_j}^{\mathbf{r}_l} \mathbf{A} \cdot d\mathbf{l} = -\phi_{lj}$$





Rotating MZMs on a Triangular Chain ($W=1$)

Superconducting
triangular islands
as a platform for
manipulating
Majorana zero
modes

Aidan Winblad


Motivation

Kitaev
Triangle

Hollow
Triangle

Braiding

Summary





Rotating MZMs on a Triangular Chain ($W=1$)

Superconducting
triangular islands
as a platform for
manipulating
Majorana zero
modes

Aidan Winblad



Motivation

Kitaev
Triangle

Hollow
Triangle

Braiding

Summary



$$L = 50, W = 1, \mu = 1.1, A = 2.35$$



Rotating MZMs on a Triangular Chain ($W=1$)

Superconducting
triangular islands
as a platform for
manipulating
Majorana zero
modes

Aidan Winblad



Motivation

Kitaev
Triangle


Hollow
Triangle

Braiding

Summary



$$L = 50, W = 1, \mu = 1.1, A = 2.35$$





Rotating MZMs on a Triangular Chain ($W=1$)

Superconducting
triangular islands
as a platform for
manipulating
Majorana zero
modes

Aidan Winblad



Motivation

Kitaev
Triangle


Hollow
Triangle

Braiding

Summary



$$L = 50, W = 1, \mu = 1.1, A = 2.35$$





Rotating MZMs on a Triangular Chain ($W=1$)

Superconducting
triangular islands
as a platform for
manipulating
Majorana zero
modes

Aidan Winblad



Motivation

Kitaev
Triangle


Hollow
Triangle

Braiding

Summary



$$L = 50, W = 1, \mu = 1.1, A = 2.35$$





Rotating MZMs on a Triangular Chain ($W=1$)

Superconducting
triangular islands
as a platform for
manipulating
Majorana zero
modes

Aidan Winblad



Motivation

Kitaev
Triangle


Hollow
Triangle

Braiding

Summary



$$L = 50, W = 1, \mu = 1.1, A = 2.35$$





Rotating MZMs on a Hollow Triangle ($W=3$)

Superconducting
triangular islands
as a platform for
manipulating
Majorana zero
modes

Aidan Winblad



Motivation

Kitaev
Triangle

Hollow
Triangle

Braiding

Summary





Rotating MZMs on a Hollow Triangle ($W=3$)

Superconducting
triangular islands
as a platform for
manipulating
Majorana zero
modes

Aidan Winblad




Motivation

Kitaev
Triangle

Hollow
Triangle

Braiding

Summary



$$L = 80, W = 3, \mu = 1.6, (A, \phi) = (0.83, 0) \rightarrow (0.77, \pi/6) \rightarrow (0.83, \pi/3) \rightarrow \dots$$



Rotating MZMs on a Hollow Triangle ($W=3$)

Superconducting
triangular islands
as a platform for
manipulating
Majorana zero
modes

Aidan Winblad




Motivation

Kitaev
Triangle


Hollow
Triangle

Braiding

Summary



$$L = 80, W = 3, \mu = 1.6, (A, \varphi) = (0.83, 0) \rightarrow (0.77, \pi/6) \rightarrow (0.83, \pi/3) \rightarrow \dots$$





Rotating MZMs on a Hollow Triangle ($W=3$)

Superconducting
triangular islands
as a platform for
manipulating
Majorana zero
modes

Aidan Winblad




Motivation

Kitaev
Triangle


Hollow
Triangle

Braiding

Summary



$$L = 80, W = 3, \mu = 1.6, (A, \varphi) = (0.83, 0) \rightarrow (0.77, \pi/6) \rightarrow (0.83, \pi/3) \rightarrow \dots$$





Rotating MZMs on a Hollow Triangle ($W=3$)

Superconducting
triangular islands
as a platform for
manipulating
Majorana zero
modes

Aidan Winblad




Motivation

Kitaev
Triangle


Hollow
Triangle

Braiding

Summary



$$L = 80, W = 3, \mu = 1.6, (A, \varphi) = (0.83, 0) \rightarrow (0.77, \pi/6) \rightarrow (0.83, \pi/3) \rightarrow \dots$$





Rotating MZMs on a Hollow Triangle ($W=3$)

Superconducting
triangular islands
as a platform for
manipulating
Majorana zero
modes

Aidan Winblad




Motivation

Kitaev
Triangle


Hollow
Triangle

Braiding

Summary



$$L = 80, W = 3, \mu = 1.6, (A, \varphi) = (0.83, 0) \rightarrow (0.77, \pi/6) \rightarrow (0.83, \pi/3) \rightarrow \dots$$





Braiding Two of Four MZMs

Superconducting
triangular islands
as a platform for
manipulating
Majorana zero
modes

Aidan Winblad





Motivation

Kitaev
Triangle

Hollow
Triangle

Braiding

Summary





Summary

Superconducting
triangular islands
as a platform for
manipulating
Majorana zero
modes

Aidan Winblad

Motivation

Kitaev
Triangle

Hollow
Triangle

Braiding

Summary

- Introduction of Peierls phase allows for a minimal Kitaev triangle, reducing fermionic sites down to 3.
- Vector potential field and its rotation allows additional tunability of topology.
- MZMs can be hosted and braided on a network of triangular islands.



Majorana fermion notation and coupling isolations

The complex fermion operator can be written as a superposition of two Majorana fermions $c_j = \frac{1}{2}(a_j + ib_j)$. Due to the nature of Majorana fermions, $a_j^\dagger = a_j$, the creation operator is $c_j^\dagger = \frac{1}{2}(a_j - ib_j)$.

$$H = -\frac{i\mu}{2} \sum_j a_j b_j - \frac{i}{2} \sum_{\langle jl \rangle} [(t \sin \phi_{jl} - \Delta \sin \theta_{jl}) a_l a_j + (t \sin \phi_{jl} + \Delta \sin \theta_{jl}) b_l b_j + (t \cos \phi_{jl} - \Delta \cos \theta_{jl}) a_l b_j - (t \cos \phi_{jl} + \Delta \cos \theta_{jl}) b_l a_j].$$

$$(t \sin \phi_{jl} - \Delta \sin \theta_{jl}) a_l a_j, \quad (1)$$

$$(t \sin \phi_{jl} + \Delta \sin \theta_{jl}) b_l b_j, \quad (2)$$



$$(t \cos \phi_{jl} + \Delta \cos \theta_{jl}) a_l b_j, \quad (3)$$

$$(t \cos \phi_{jl} - \Delta \cos \theta_{jl}) b_l a_j \quad (4)$$



Braiding MZM in a Small Network of Triangles

Aidan Winblad





Braiding MZM in a Small Network of Triangles

Aidan Winblad

