

Superconducting triangular islands as a platform for manipulating Majorana fermions

Aidan Winblad¹ and Hua Chen^{1,2}

¹*Department of Physics, Colorado State University, Fort Collins, CO 80523, USA*

²*School of Advanced Materials Discovery, Colorado State University, Fort Collins, CO 80523, USA*

PLACEHOLDERAPS MM 2022 Abstract** We study the possibility of obtaining robust Majorana modes at the corners of triangular islands of different superconductor models, with the goal of finding alternative structures that can serve as building blocks of topological quantum computation. By considering both spinless p-wave and ferromagnetic Rashba s-wave superconductor models on an equilateral triangle subject to inhomogeneous supercurrents, we found that Majorana corner modes can generally appear if the system being considered can be made equivalent to a triangular chain model, which can be understood by calculating the bulk Z2 topological invariant. We also discussed the robustness of the corner modes in possible experimental realizations of the triangular islands.

I. INTRODUCTION

PLACEHOLDER

OUTLINE: (Sec I is ..., Sec II is ...)

II. LINEAR VECTOR POTENTIAL EFFECTS ON 1D CHAINS

PLACEHOLDER

A. Gap closing/reopening (Topological Change) and phase diagram

Recent work has predicted that placing a supercurrent along a superconducting chain can induce a topological phase change. Another way of thinking about this is there is a constant vector potential that induces a constant phase on the hopping amplitudes on the superconducting chain. Here we will show that using a linear vector potential can also induce a topological phase change, i.e. closing and reopening a gap. **Maybe here we can state the Hamiltonian in the supplementary? Explain we can't solve this analytically, thus we proceed with a numerical approach starting with the lattice space Hamiltonian.** Let us start with the Hamiltonian of a spinless or spin-polarized p-wave superconductor chain

$$\mathcal{H}_{ch} = \sum_j (-tc_{j+1}^\dagger c_j + \Delta c_{j+1}^\dagger c_j^\dagger + h.c.) - \mu c_j^\dagger c_j, \quad (1)$$

where t is the hopping amplitude, Δ is the superconducting order parameter, and μ is the chemical potential. Next, we will apply Peierls substitution to our creation(annihilation) operators

$$\begin{aligned} c_{j+1}^\dagger c_j &\rightarrow c_{j+1}^\dagger c_j \exp\left(-\frac{ie}{\hbar} \int_{r_j}^{r_{j+1}} \mathbf{A} \cdot d\mathbf{l}\right) \\ &\rightarrow c_{j+1}^\dagger c_j e^{i\phi_{j+1,j}}. \end{aligned} \quad (2)$$

Here the electron charge, e , and Planck's constant, \hbar , will be treated as natural numbers. The vector potential will be linear, symmetric about the y-axis, and point along the x-axis, $\mathbf{A} = Bx\hat{\mathbf{x}}$ for $-L/2 \leq x \leq L/2$. One finds that t is the only term that picks up a phase, thus our modified Hamiltonian looks like

$$\mathcal{H}_{ch} = \sum_j (-te^{i\phi_{j+1,j}} c_{j+1}^\dagger c_j + \Delta c_{j+1}^\dagger c_j^\dagger + h.c.) - \mu c_j^\dagger c_j. \quad (3)$$

To calculate the Majorana number we need to rewrite our Hamiltonian to be in skew-symmetric form. One such way is to write it in Majorana fermion basis, the transformation matrix is of the form

$$u = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 & 1 \\ -i & i \end{pmatrix}$$

Since we are in lattice space we need to expand for the system size, we simply include a tensor product with an identity matrix of the size of the system, $U = u \otimes I_n$. We can now arrive at the skew-symmetric matrix with the following equation

$$A_{ch} = -iU\mathcal{H}_{ch}U^\dagger. \quad (4)$$

The Majorana number, M , of a 1D chain is defined as

$$\mathcal{M} = \text{sgn}[\text{Pf}(A_{ch})], \quad (5)$$

where Pf stands for the Pfaffian of a skew-symmetric matrix. When $\mathcal{M} = -1$, the chain is in a non-trivial topology and we would expect Majorana zero modes at the edges of a finite chain. As we sweep through a range of B values for our vector potential we expect to see a gap closing, $\mathcal{M} = 1$, and our system take on a trivial topology, thus no Majorana zero modes would exist on the edge of a finite chain. **Taking a look at figure XX we can see for varying μ and B values where our system is trivial and non-trivial.**

B. Majorana Edge modes on a double chain

Need to make this script and plot still

III. TRIANGULAR ISLANDS

A. 3-site problem and kitaev limit for tri.chain, solid triangle

We now expand our analysis to triangular islands. The chain Hamiltonian in Eq. 3 can be generalized as

$$\mathcal{H} = \sum_{\langle j,l \rangle} (-te^{i\phi_{l,j}} c_l^\dagger c_j + \Delta e^{i\theta_{l,j}} c_l^\dagger c_j^\dagger + h.c.) - \sum_j \mu c_j^\dagger c_j, \quad (6)$$

where $\langle j,l \rangle$ are nearest neighbors on a triangular lattice and $\theta_{l,j}$ is the angle between the two sites. It is difficult to come to any conclusion while in the complex fermion basis, but if we write our Hamiltonian in Majorana fermion basis we can get a clearer picture. **Maybe a see supplementary material XX for derivation here.** The triangular Hamiltonian now takes the form

$$\begin{aligned} \mathcal{H} = & -\frac{i\mu}{4} \sum_j (a_j b_j - b_j a_j) \\ & -\frac{i}{4} \sum_{\langle j,l \rangle} [(t \sin \phi_{l,j} - \Delta \sin \theta_{l,j}) a_l a_j \\ & + (t \sin \phi_{l,j} + \Delta \sin \theta_{l,j}) b_l b_j \\ & + (t \cos \phi_{l,j} + \Delta \cos \theta_{l,j}) a_l b_j \\ & - (t \cos \phi_{l,j} - \Delta \cos \theta_{l,j}) b_l a_j]. \end{aligned} \quad (7)$$

To clear things up, $\phi_{l,j} = -\phi_{j,l}$ since the direction of integration is reversed and the angle $\theta_{l,j} = \theta_{j,l} + \pi$, this ensures we have a Hermitian Hamiltonian.

Let us consider a 3-point triangle lattice, where each lattice point is a complex fermion housing two Majorana fermions. The bottom left point will have a_1, b_1 , bottom right has a_2, b_2 , and the top point has a_3, b_3 . Similar to Kitaev we will make the same assumptions $t = \Delta$ and $\mu = 0$. Doing this lets us see a combination of trig terms. We need some of these combination to go to zero. Notice how each individual row looks like a Kitaev chain, we need to look at how the rows interact with each other. Since our goal is to have two Majorana zero modes at the bottom corners, let's aim to have a_1 and b_2 be such modes. Anytime a_1 or b_2 appear in our equation we need its trig terms to cancel, eliminating those particles coupling to the rest of the system. Let us look at the energy going from site 1 to site 3, $\theta = \pi/3$, we notice the first and last term should be

$$a_3 a_1 (\sin \phi_{31} - \sin \pi/3) = 0, \quad (8)$$

$$b_3 a_1 (\cos \phi_{31} - \cos \pi/3) = 0. \quad (9)$$

This is true if $\phi_{31} = \pi/3$. Now let's consider the energy from site 3 to site 2. The phase angle $\theta = -\pi/3$ and the

two simplified equations involving b_2 are

$$(\sin \phi_{23} - \sin(\pi/3)) b_2 b_3 = 0,$$

$$(\cos \phi_{23} - \cos(\pi/3)) b_2 a_3 = 0.$$

Here we see $\phi_{23} = \pi/3$. The linear vector potential one can get with the desired phase is

$$\mathbf{A} = \frac{8\pi}{3\sqrt{3}a^2} x \hat{\mathbf{y}} \quad (10)$$

We can now extrapolate from a 3-point triangle to a full triangular island. The goal being to ensure Majorana zero modes at the bottom two vertices. If instead we have a triangular island with n_r -many rows we find

$$\mathbf{A} = \frac{8\pi}{3\sqrt{3}a^2} \frac{x}{2n_r - 3} \hat{\mathbf{y}} = Bx \hat{\mathbf{y}}. \quad (11)$$

Solving the Hamiltonian numerically we can see how turning on this Peierls phase affects the eigenenergies.




FIG. 1. For increasing vector potential strength MZM's appear at the critical strength, denoted with a dotted black line. Notice the density of edge states makes it unclear the Majorana zero modes live in a clean gap.

We clearly see MZM's appear at the critical, unfortunately it doesn't seem to live within a clean gap. To get around this we instead think of using a triangular chain instead to remove a majority of the edge states.

While we have effectively reduced the number of states living in the gap we see the MZM's are doubly degenerate. This is a consequence of a triangular chain's geometry. The next excited state becomes a zero mode due to the geometry forcing the wave function to be well separated and recombines with the other expected MZM's. This leads us to think we should use a finite width for the sides of the triangle.




FIG. 2. As the strength of the vector potential increases MZM's appear at the critical strength, denoted as the dotted black line. Transparency has been added to look for any degeneracies. While a cleaner gap can be seen for the MZM's they are degenerate.

B. Finite width (hollow triangle)

1. *Majorana number (PBC)*
2. *Corner modes (open BC)*




FIG. 3. As the strength of the vector potential increases MZM's appear at the critical strength, denoted as the dotted black line. Transparency has been added to look for any degeneracies.

IV. CONCLUSIONS AND DISCUSSION

PLACEHOLDER

ACKNOWLEDGMENTS

Supported by XYZ Grant No. XXXXXX etc.