

Background

Motivation

Formulation

Results

Summary



Majorana Corner Modes in Triangular Superconductor Islands

Aidan Winblad
Hua Chen

Department of Physics
Colorado State University

March 18, 2022




Outline

- Background:
 - Majorana fermions in particle physics
 - Majorana fermions in condensed matter
- Motivation:
 - P-wave superconductors and braiding
 - 1D wires and T-junctions
 - Triangular structures for braiding
- Formulation:
 - Peierls substitution in Kitaev's model
- Results:
 - Topological phase
 - Triangular chain
 - Hollow triangles
- Summary
 - Future work
 - Additional projects



Background



Enrico Fermi

Paul Dirac



Ettore Majorana

(Dirac) Fermion

- Fermi statistics
- Particle \neq Antiparticle : $c \neq c^\dagger$
- Charged
- Electron

Majorana Fermion

- Fermi statistics
- Particle = Antiparticle : $c = c^\dagger$
- Neutral
- Neutrino? Dark Matter?



Background


Background

Motivation

Formulation

Results

Summary




MAJORANA project:
neutrinoless double beta ($0\nu\beta\beta$) decay

- Are neutrinos Majorana fermions?
- If yes, the standard needs revision
- Negative results for Majorana particles




Background

- Superconductors
 - Cooper pairs
 - Electron-phonon interaction pairs two electrons with opposite spin and momenta.
 - Bogoliubov quasiparticles
 - Excitation from ground state, pairs an electron to a hole.
 - Zero-energy excitations may be Majorana fermions.
 - If so, they come in pairs.





Background



Kitaev, Phys. Uspekhi **44**, 131 (2001)

Majorana fermion notation

$$c_j = \frac{1}{2}(a_j + ib_j). \quad (1)$$



Background

Hamiltonian for a 1D tight-binding chain with spinless p -wave superconductivity

$$\mathcal{H}_{chain} = -\mu \sum_j^N c_j^\dagger c_j - \sum_j^{N-1} t c_j^\dagger c_{j+1} + |\Delta| c_j c_{j+1} + h.c. \quad (2)$$

Hamiltonian in Majorana basis

$$\mathcal{H}_{chain} = \frac{i}{2} \sum_j -\mu a_j b_j + (t + |\Delta|) b_j a_{j+1} + (-t + |\Delta|) a_j b_{j+1}. \quad (3)$$

$t = |\Delta| = 0$ and $\mu < 0$, trivial phase


$$\mathcal{H} = -\frac{i\mu}{2} \sum_j a_j b_j. \quad (4)$$

$t = |\Delta| > 0$ and $\mu = 0$, non-trivial (topological) phase

$$\mathcal{H} = it \sum_j b_j a_{j+1}. \quad (5)$$



Background



Kitaev, *Phys. Uspekhi* **44**, 131 (2001)

Intersite fermion representation

$$\tilde{c}_j = \frac{1}{2}(a_{j+1} + ib_j). \quad (6)$$




The highly non-local fermion state

$$f = \frac{1}{2}(a_1 + ib_N), \quad (7)$$

corresponds to zero energy. This is still true for $|\mu| < 2t$



Background



Mourik, *Science* **336**, 1003 (2012).



Background



Motivation

Formulation

Results

Summary

Background





Nadj-Perge, *Science* **346**, 602 (2014).



Background

a



Mn atoms (red spheres) on top of superconducting Nb (brown spheres).

Schneider et al., *Nature Nanotechnology* **17**, 384 (2022).



Motivation

Aidan Winblad


Background

Motivation

Formulation

Results


Summary



$$T_i : \begin{cases} \gamma_i \rightarrow \gamma_{i+1} \\ \gamma_{i+1} \rightarrow -\gamma_i \\ \gamma_j \rightarrow \gamma_j \end{cases} \quad \text{for } j \text{ and } j+1$$

$$\tau(T_i)\gamma_j[\tau(T_i)]^{-1} = T_i(\gamma_j)$$


$$\tau(T_i) = \exp\left(\frac{\pi}{4}\gamma_{i+j}\gamma_i\right) = \frac{1}{\sqrt{2}}(1 + \gamma_{i+j}\gamma_i)$$



Ivanov, PRL **86**, 268 (2000).




Motivation



$$\mathcal{H}_T = -\mu \sum_j c_j^\dagger c_j - \sum_j t c_j^\dagger c_{j+1} + |\Delta| e^{i\phi} c_j c_{j+1} + h.c. \quad (8)$$

$$c_j = e^{-i(\phi/2)} (\gamma_{j+1,1} + i\gamma_{j,2})/2 \quad (9)$$


- Take pairing term $|\Delta| e^{i\phi} c_j c_{j+1}$ such that the site indices:
- Increase moving \rightarrow / \uparrow in the horizontal/vertical wires: $\phi = 0$,
- Decrease moving \leftarrow / \downarrow in the horizontal/vertical wires: $\phi = \pi$.





Motivation


- Consider triangular islands, topologically similar to T-junctions.
- Islands of three-fold rotational symmetry occur naturally in epitaxial growth on close-packed metal surfaces.
- Good platform to transition from 2D to 1D topological superconductor.



Triangular Co islands on Cu(111).
Pietzsch et al., *PRL* **96**, 237203 (2006)



Previous Work



Alicea, *PRB* **81**, 125318 (2010).

$$c_j = (c_{j\uparrow}, c_{j\downarrow})^T \quad (10)$$

s-wave SC paring term:

$$\mathcal{H}_{SC} = \sum_j \Delta c_{j\uparrow}^\dagger c_{j\downarrow}^\dagger + h.c. \quad (11)$$

Quantum well:

$$\mathcal{H}_0 = \sum_j (6t - \mu) c_j^\dagger c_j - \sum_{\langle j, l \rangle} (t c_l^\dagger c_j + h.c.) \quad (12)$$

Rashba spin-orbit coupling:

$$\mathcal{H}_R = -it_R \sum_{\langle j, l \rangle \alpha\beta} c_{l\alpha}^\dagger (\boldsymbol{\sigma}_{\alpha\beta} \times \hat{\mathbf{r}}_{lj}) \cdot \hat{\mathbf{z}} c_{j\beta} \quad (13)$$

Zeeman field:

$$\mathcal{H}_Z = \sum_j c_j^\dagger \mathbf{V} \cdot \boldsymbol{\sigma} c_j \quad (14)$$



Background




Motivation

Formulation

Results

Summary

Previous Work





Topological phase transition induced by a supercurrent

Majorana corner modes in triangular superconductor islands

Aidan Winblad


Background

Motivation

Formulation

Results

Summary



Romito, PRB 85, 020502(R) (2012).



Topological phase transition induced by a supercurrent

Majorana corner modes in triangular superconductor islands

Aidan Winblad


Background

Motivation





Formulation

Results

Summary



Takasan, *PRB* **106**, 014508 (2022).





Background

Motivation

Formulation

Results

Summary

Peierls substitution

$$\begin{aligned} c_{j+1}^\dagger c_j &\rightarrow c_{j+1}^\dagger c_j \exp\left(-\frac{ie}{\hbar} \int_{r_j}^{r_{j+1}} \mathbf{A} \cdot d\mathbf{l}\right) \\ &\rightarrow c_{j+1}^\dagger c_j e^{i\phi_{j+1,j}}. \end{aligned} \tag{15}$$

$$\mathcal{H}_{chain} = \sum_j (-te^{i\phi_{j+1,j}} c_{j+1}^\dagger c_j + \Delta c_{j+1}^\dagger c_j^\dagger + h.c.) - \mu c_j^\dagger c_j. \tag{16}$$



Kitaev Limit with Vector Potential on a Triangular Island

Majorana corner
modes in
triangular
superconductor
islands

Aidan Winblad


Background

Motivation

Formulation

Results

Summary



$$\mathcal{H} = \sum_{\langle j,l \rangle} \left[-te^{i\phi_{l,j}} c_l^\dagger c_j + \Delta e^{i\theta_{l,j}} c_l^\dagger c_j^\dagger + h.c. \right] - \sum_j \mu c_j^\dagger c_j$$

$$\phi_{l,j} = -\frac{e}{\hbar} \int_{\mathbf{r}_j}^{\mathbf{r}_l} \mathbf{A} \cdot d\mathbf{l}$$

$$\mathbf{A} = -\frac{2\pi}{3\sqrt{3}a} \hat{\mathbf{y}}$$



Triangular Chain

Majorana corner modes in triangular superconductor islands

Aidan Winblad

Background

Motivation




Formulation

Results

Summary

$$\mathbf{A}(x) = Bx\hat{\mathbf{y}}$$

$$B_0 = \frac{8\pi}{3\sqrt{3}a^2(2n_r - 3)}$$





Hollow Triangle

Aidan Winblad



Background

Motivation

Formulation

Results

Summary





Background

Motivation

Formulation

Results

Summary


- Introduction of vector potential allows for additional tunability of topology.
- Triangular islands with a gapped interior can be a promising platform for hosting and manipulating MZMs.
- Next steps
 - Search for safe MZMs in hollow triangles outside the Kitaev limit.
 - Develop a robust braiding scheme.



Additional results from Schneider et al.

Majorana corner
modes in
triangular
superconductor
islands

Aidan Winblad





Majorana fermion notation and coupling isolations

Aidan Winblad

The complex fermion operator can be written as a superposition of two Majorana fermions $c_j = \frac{1}{2}(a_j + ib_j)$. Due to the nature of Majorana fermions, $a_j^\dagger = a_j$, the creation operator is $c_j^\dagger = \frac{1}{2}(a_j - ib_j)$.

$$H = -\frac{i\mu}{4} \sum_j (a_j b_j - b_j a_j) - \frac{i}{4} \sum_{<j,l>} [(t \sin \phi - \Delta \sin \theta) a_l a_j + (t \sin \phi + \Delta \sin \theta) b_l b_j + (t \cos \phi + \Delta \cos \theta) a_l b_j - (t \cos \phi - \Delta \cos \theta) b_l a_j].$$

$$(t \sin \phi_{j,l} - \Delta \sin \theta_{j,l}) a_l a_j, \quad (17)$$


$$(t \sin \phi_{j,l} + \Delta \sin \theta_{j,l}) b_l b_j, \quad (18)$$

$$(t \cos \phi_{j,l} + \Delta \cos \theta_{j,l}) a_l b_j, \quad (19)$$

$$(t \cos \phi_{j,l} - \Delta \cos \theta_{j,l}) b_l a_j \quad (20)$$



Triangular chain degeneracy





Hollow triangle degeneracy?

Aidan Winblad

