

Majorana Corner Modes in Triangular Superconductor Islands

Aidan Winblad
Hua Chen

Department of Physics
Colorado State University

March 18, 2022

Motivation

Formulation

Results

Summary

Motivation

Majorana corner
modes in
triangular
superconductor
islands

Aidan Winblad

Motivation

Formulation

Results

Summary

- P-wave superconductors contain half-quantum vortices.
 - Majorana fermions located at core of a vortex.
 - Braiding vortices exhibits Non-Abelian statistics.
- 1D p-wave superconductors host Majorana fermions on end points.
 - Possibly measured in real systems: Mourik, *Science* **336**, 1003 (2012)
Nadj-Perge, *Science* **346**, 602 (2014)
- Quasi-1D T-junction
 - Braiding of Majorana fermions is defined for 2D.
 - In practice challenging to make, but still feasible and seriously pursued.



Alicea, *Nature Phys.* **7**, 412 (2011)

Motivation

Motivation

Formulation

Results

Summary

- Consider triangular islands, topologically similar to T-junctions.
- Islands of three-fold rotational symmetry occur naturally in epitaxial growth on close-packed metal surfaces.
- Good platform for transition from 2D to 1D topological superconductor.



../../../../images/triangular-islands.png

Triangular Co islands on Cu(111).

Pietzsch et al., *PRL* **96**, 237203 (2006)

Previous Work

Majorana corner
modes in
triangular
superconductor
islands

Aidan Winblad

Motivation

Formulation

Results

Summary

../../../../images/tunable-semiconductor.pdf

../../../../images/dispersion-in-plane.pdf

../../../../images/energy-spectra-in-plane.pdf

../../../../images/wavefunction-2-in-plane.pdf

Kitaev Limit with Vector Potential on a Triangular Island

Motivation

Formulation

Results

Summary



../../../../images/left-triangle-corner.pdf

$$\mathcal{H} = \sum_{\langle j,l \rangle} \left[-te^{i\phi_{l,j}} c_l^\dagger c_j + \Delta e^{i\theta_{l,j}} c_l^\dagger c_j^\dagger + h.c. \right] - \sum_j \mu c_j^\dagger c_j$$

$$\phi_{l,j} = -\frac{e}{\hbar} \int_{\mathbf{r}_j}^{\mathbf{r}_l} \mathbf{A} \cdot d\mathbf{l}$$

$$\mathbf{A} = -\frac{2\pi}{3\sqrt{3}a} \hat{\mathbf{y}}$$

Majorana Number of 1D Chain with Vector Potential

Majorana corner
modes in
triangular
superconductor
islands

Aidan Winblad

Motivation

Formulation

Results

Summary

μ `../../../../images/Majorana-number.pdf`

ϕ

$\phi = \pi$ `../../../../images/kitaev-chain-mu_pi.pdf`

Triangular Chain

Majorana corner
modes in
triangular
superconductor
islands

Aidan Winblad

Motivation

Formulation



Results

Summary



$$\mathbf{A}(x) = Bx\hat{\mathbf{y}}$$

$$B_0 = \frac{8\pi}{3\sqrt{3}a^2(2n_r - 3)}$$

../../../../images/vector-potential-field.pdf



Hollow Triangle



Summary

Motivation

Formulation

Results

Summary

- Introduction of vector potential allows for additional tunability of topology.
- Triangular islands with a gapped interior can be a promising platform for hosting and manipulating MZMs.
- Next steps
 - Search for safe MZMs in hollow triangles outside the Kitaev limit.
 - Develop a robust braiding scheme.

Majorana fermion notation and coupling isolations

The complex fermion operator can be written as a superposition of two Majorana fermions $c_j = \frac{1}{2}(a_j + ib_j)$. Due to the nature of Majorana fermions, $a_j^\dagger = a_j$, the creation operator is $c_j^\dagger = \frac{1}{2}(a_j - ib_j)$.

$$H = -\frac{i\mu}{4} \sum_j (a_j b_j - b_j a_j) - \frac{i}{4} \sum_{\langle j,l \rangle} [(t \sin \phi - \Delta \sin \theta) a_l a_j + (t \sin \phi + \Delta \sin \theta) b_l b_j + (t \cos \phi + \Delta \cos \theta) a_l b_j - (t \cos \phi - \Delta \cos \theta) b_l a_j].$$


$$(t \sin \phi_{j,l} - \Delta \sin \theta_{j,l}) a_l a_j, \quad (1)$$

$$(t \sin \phi_{j,l} + \Delta \sin \theta_{j,l}) b_l b_j, \quad (2)$$

$$(t \cos \phi_{j,l} + \Delta \cos \theta_{j,l}) a_l b_j, \quad (3)$$

$$(t \cos \phi_{j,l} - \Delta \cos \theta_{j,l}) b_l a_j \quad (4)$$

Triangular chain degeneracy



Hollow triangle degeneracy?

